



MATREYA

Fabry disease

Niemann-Pick Types A and B

Sialidosis

Farber disease

Gaucher disease

Lipids, Biochemicals, and Standards for Life Science Research

Sandoff disease

Tay-Sachs disease

Krabbe disease

Metachromatic leukodystrophy

GM₁ gangliosidosis

2017-2018

www.matreya.com

MATREYA

About Matreya LLC

Matreya strives to develop, manufacture and deliver products of the highest value to our customers. Quality will always be the best achievable by state-of-the-art techniques, typically greater than 98%. We strive for rapid delivery. 95+% of our products are shipped within 24 hours of receipt of an order. Within the area of sphingolipids and glycolipids, we have earned a reputation as the preferred problem solver and technology leader.

When you demand quality and consistency, you may rely on Matreya products.

Matreya Products for Life Science Research.

We offer one of the widest selection of ceramides for intracellular signaling research available. We stock antibodies to glycosphingolipids as well as inhibitors of enzymes involved in glycosphingolipid metabolism.

Our products provide the valuable tools necessary for the study of cell membrane and its structure, growth regulators in the cellular metabolism, and intracellular mediators.

We are able to make our products better and better with the latest technology in Chromatography and Mass Spectrometry.

We are proud to offer our products as a valuable tool for your life science research needs.

Matreya Products for Microbiology Research.

Matreya stocks unusual fatty acid standards produced by bacteria that are useful for culture characterization.

Matreya Products for the Food and Agriculture Industries.

Many of Matreya's fatty acid products have been industry standards for years. Our fatty acids and methyl esters are used as standards in analysis and quality control.

Custom Preparations.

Matreya's experience in chemical synthesis and the extraction and purification of natural products allows us to produce custom preparations with the same high quality and purity as the products listed in our catalog. Depending on the complexity of the molecule, a 30% nonrefundable deposit may be required prior to synthesis. This deposit will be deducted from the final invoice upon completion of the project. Delivery will be 4 to 12 weeks after receipt of an order.

If you can't find a product in the catalog, please check the INDEX, where we also try to list common synonyms for our products.

www.matreya.com

TABLE OF CONTENTS

TABLE OF CONTENTS	ii
Technical Service	iv
Natural Products.....	iv
Storage	iv
Package Weight	iv
Matreya's Mission.....	iv
Lysosomal Storage Disorders Pathways Chart	v
Sphingoid Bases, Sphingolipids and Glycosphingolipids	1
Sphingosines.....	2
Synthetic Sphingosines with C18 Sphingoid Base	2
Synthetic Sphingosines with Sphingoid Bases other than C18.....	2
Synthetic Dihydrosphingosines.....	3
3-Keto-Dihydrosphingosines.....	4
Phytosphingosines.....	5
Other Sphingosine Derivatives and Precursors.....	5
Ceramides	6
Synthetic Ceramides Derived from C18-Sphingosine	7
2-Hydroxy Ceramides	11
Ceramide Made from Sphingosines with Sphingoid Bases Other Than C18.....	11
Dihydroceramides	11
2-Hydroxy Dihydroceramides.....	12
Ceramides From Natural Sources	13
Phytoceramides	14
Fluorescent Ceramides.....	15
Phosphosphingolipids	17
Sphingomyelins.....	17
Sphingosylphosphorylcholines (SPC).....	19
Sphingosine and Ceramide Phosphates	19
Fluorescent Sphingomyelins.....	20
Glycosphingolipids	21
Galactosylceramides.....	21
Glucosylceramides.....	24
Sulfatides	26
Lactosylceramides	29
Ceramide Trihexosides (Globotriaosylceramides).....	31
Globosides	33
Stable Isotope Labeled Glycolipids.....	33
Fluorescent Glycolipids.....	34
Gangliosides	36
Glycosphingolipid Reference Mixtures for TLC	39
Antibodies Directed Against Glycolipids	40
Enzyme Inhibitors.....	41
Glycerolipids	45
Glycerophospholipids.....	45
Natural Phospholipids	45
Synthetic Phospholipids.....	47
Phosphatidic Acid Derivatives	47
Phosphatidylcholines	48
Phosphatidylglycerols	49
Phosphatidylethanolamines.....	50
Phosphatidylinositol Phosphates	51
Bacterial Tetraethers.....	51
Glycosyl Glycerides.....	52

Fatty Acids	52
Simple Fatty Acids.....	52
Saturated Fatty Acids and Methyl Esters.....	52
Unsaturated Fatty Acids and Methyl Esters.....	57
Trans Fatty Acids and Methyl Esters.....	64
Conjugated Linoleic Acid Isomers (CLA).....	66
Hydroxy Fatty Acids	68
2-Hydroxy Fatty Acids and Methyl Esters.....	68
3-Hydroxy Fatty Acids and Methyl Esters.....	71
Omega Hydroxy Fatty Acids.....	74
Branched and Cyclic Fatty Acids	75
Iso-Fatty Acids and Methyl Esters	75
Anteiso-Fatty Acids and Methyl Esters	76
Other Branched Methyl Fatty Acids	77
Cyclopropyl Fatty Acids and Methyl Esters	78
Unusual Fatty Acids and Derivatives	78
Vitamin E	79
Tocopherols	79
Tocotrienols	80
Sterols.....	81
Cholestane Derivatives	81
Plant Sterols and Steryl Glucosides	82
Standards and Reference Compounds	83
Food Industry Mixtures	83
Polyunsaturated Fatty Acid Methyl Ester Mixtures	84
Carbohydrate Mixtures.....	85
Other Fatty Acid Methyl Ester Mixtures	85
AOCS Animal and Vegetable Oil Reference Mixtures (RM Mixtures).....	86
Custom Mixtures	88
GLC Standard Mixtures	89
Water Soluble Fatty Acid Mixtures.....	90
Microbiology Standard Mixtures	91
Biochemical Research Standard Mixtures.....	91
Glycosphingolipid Reference Mixtures for TLC	92
Labeled Standards	93
Stable Isotope Labeled Standards.....	93
Fluorescent Standards.....	95
Biotin Labeled Standards.....	98
Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC	100
Literature References.....	105
Cross Reference for Product Numbers and Catalog Pages	106
Product Name Index	109
Category Index.....	113
 Tables	
Table I. AOCS Oil Reference Mixtures.....	86
Table II. Standards for GC Analysis	89
Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC	100

All chemicals listed in this catalog are for research only. They are not intended for drug or diagnostic use, human consumption or to be used in food or food additives. Matreya assumes no liability for any use of these chemicals by the end user. We believe the information in this catalog, offered in good faith, is accurate.

Limited Warranty: All Matreya Products, except those specifically exempted, are warranted (for 30 days) to be free of defects in materials and workmanship, if properly stored. Any replacements required as a result of such defects will be made without charge provided that such defective products are returned with a written explanation. Please request a Returned Goods Authorization before returning products under this warranty.

Technical Service

Our technical service department may be contacted by telephone at 800-342-3595 (US), 814-355-1030 (world-wide) or by e-mail at techservice@matreya.com.

Natural Products

Some of our glycolipids are extracted from natural sources. These products have a normal heterogeneity in their lipid components, particularly in the fatty acids. Variations include carbon chain length as well as the presence or absence of 2-hydroxy fatty acids. Products based on sphingosine may contain longer chain sphingoid bases as well as chains with multiple double bonds. This heterogeneity may result in additional spots showing on TLC plates or multiple peaks in LC analyses. We have listed the typical fatty acid compositions of our natural products in Table III.

Storage

Catalog items in unopened containers are stable for at least one year when stored under the conditions indicated in the catalog listing. Items containing unsaturated fatty acids are subject to oxidation and should be stored in a solution of organic solvents or under argon. Glycolipids and phospholipids should not be stored in aqueous solutions due to potential hydrolysis.

Package Weight


Unless otherwise specified, the package will contain at least the indicated amount and usually slightly more. The user is cautioned to always measure the required amount from the container.

Matreya's Mission

Matreya is committed to manufacturing high purity lipids to be used as research standards in the biotechnology and pharmaceutical areas. These lipids will be offered world-wide at a fair market price, and at a profit sufficient to assure company growth, for the benefit of its customers, employees, share holders, and community. Matreya will also be committed to fast delivery, excellent technical backup, new product development, safety, and environmentally friendly.

Lysosomal Storage Disorders Pathways Chart

Glycosphingolipid
Enzyme
Sphingolipidosis




Ceramide


Glucosylcerebroside

LacCer


alpha-Galactosidase A
Fabry's disease



Ganglioside neuraminidase
Sialidosis




Monosialoganglioside GM₃




Gb3 (CTH)
lyso-Gb3 (lyso-CTH)

Hexosaminidase A
Tay-Sachs disease
Sandhoffs disease




beta-Hexosaminidase A + B
Sandhoffs disease





Gb4

Monosialoganglioside GM₂



Acid-beta-Galactosidase
GM1 gangliosidosis



Monosialoganglioside GM₁


Spingoid Bases, Spingolipids and Glycosphingolipids

Spingoid bases such as spingosine are the characteristic structural unit of the spingolipids. The bases are long chain aliphatic amines, containing two or three hydroxyl groups, and typically a *trans*-double bond at C4. In animal tissues, the most abundant base is spingosine with a C18 aliphatic chain containing a double bond in position 4. The saturated analogue is dihydrosphingosine or spinganine. In plants, the common long chain base is the 4 hydroxy saturated base phytosphingosine.

Spingolipids are widely distributed in animal tissues, particularly cell membranes. Spingoid bases linked to fatty acids via an amide bond at C2 are ceramides and are present in trace amounts in most tissues. Glycosphingolipids (ceramides having various mono- and oligosaccharides on the OH group at C1) are neutral glycosphingolipids (i.e., cerebrosides and globosides). Those with sialic acid derivatized sugars are acidic glycolipids (i.e., gangliosides). They are amphiphilic and can be solubilized in buffers via sonication and micelle formation.


Gangliosides are present in substantial amounts in nerve cell membranes, and together with globosides are found in the membranes of white and red blood cells. These plus the glycosphingolipids of the lacto- and neolacto-series are involved in cell recognition (e.g. blood group determinants). Glycolipid expression on the surface of cells determines their antigenicity as well as their status, i.e. differentiated vs. undifferentiated (embryonic), normal vs. malignant, etc. (1). The ganglioside GM1 stimulates nerve growth (2,3) and has been reported to have a curative effect on experimental Parkinsonism (4). For an overview, see (5). Gangliosides are also being investigated as potential anti-tumor vaccines (6). Glycosphingolipids are also essential for the correct functioning of cell surface receptors (7). Matreya is your best source for many spingolipids. Most of Matreya's spingosines and ceramides are fully synthetic and as such 98%+ pure. Others, particularly the glycosphingolipids are highly purified natural products (98%+), and can be used either as standards or biochemical reagents without further purification.

Through total synthesis, all four isomers of spingosine are available as well as a number of spingosines with other than 18 carbons and a number of ceramides (for details in using ceramides in cell culture see Hauser et al. [9]). Fluorescent labeled ceramides, glycosphingolipids and spingomyelins are also available for study. D. N. Brindley and his group have been exploring the interaction of ceramides, spingosine and spingosine 1-phosphate in regulating DNA synthesis and phospholipase D activity. **See Literature References on page 106.**



Sphingosines

Synthetic Sphingosines with C18 Sphingoid Base



1802 **D-erythro-Sphingosine** **25 mg** **95.00**
Sphingosine with C18 chain

$C_{18}H_{37}NO_2$ **Mol. Wt.:** 299 **CAS#:** 123-78-4
Source: synthetic **Purity:** 98+% by TLC, GC **Identity:** confirmed by MS
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$
Activity: Selective inhibitor of phosphokinase C

2079 **D-erythro-Sphingosine, D9** **1 mg** **385.00**
15,15,16,16,17,17,18,18,18-D9-2-Amino-octadec-4-ene-1,3-diol

$C_{18}H_{28}D_9NO_2$ **Mol. Wt.:** 309 **Identity:** confirmed by MS
Source: synthetic **Purity:** 98+% by TLC, GC, HPLC
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$

1806 **L-threo-Sphingosine** **10 mg** **325.00**
L-threo-Sphingosine, C18 chain

$C_{18}H_{37}NO_2$ **Mol. Wt.:** 299 **CAS#:** 25695-95-8
Source: synthetic **Purity:** 98+% by TLC, GC
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$

1826 **L-erythro-Sphingosine** **5 mg** **365.00**
L-erythro-Sphingosine, C18 chain

$C_{18}H_{37}NO_2$ **Mol. Wt.:** 299 **CAS#:** 6036-75-5
Source: synthetic **Purity:** 98+% by TLC, GC
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$

Synthetic Sphingosines with Sphingoid Bases other than C18

Varying chain lengths allow the study of translocation effects of sphingosines and ceramides into cells.


1838 **D-erythro-C12-Sphingosine** **5 mg** **325.00**
Sphingosine with C12 chain

$C_{12}H_{25}NO_2$ **Mol. Wt.:** 215 **CAS#:** 128427-86-1
Source: synthetic **Purity:** 98+% by TLC, GC **Identity:** confirmed by MS
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$

1833 **D-erythro-C14-Sphingosine** **5 mg** **325.00**
Sphingosine with C14 chain

$C_{14}H_{29}NO_2$ **Mol. Wt.:** 243 **CAS#:** 24558-60-9
Source: synthetic **Purity:** 98+% by TLC, GC, HPLC **Identity:** confirmed by MS
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$


2082	D-erythro-C17-Sphingosine Sphingosine with C17 chain	5 mg	280.00
	$C_{17}H_{35}NO_2$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 286 Purity: 98+% by TLC, GC, HPLC Solubility: chloroform, ethanol, methanol	CAS#: 6918-48-5 Identity: confirmed by MS



1840	D-erythro-C20-Sphingosine Sphingosine with C20 chain	5 mg	315.00
	$C_{20}H_{41}NO_2$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 328 Purity: 98+% by TLC, GC, HPLC Solubility: chloroform, ethanol, methanol, DMSO	CAS#: 6918-49-6

Synthetic Dihydrosphingosines

D,L-*threo*-Dihydrosphingosine has also been found to be a significant inhibitor of sphingosine kinase (8). The D,L-*erythro*-isomer has been used as an inactive control. We offer all four isomers in pure form making detailed studies possible. Safingol, the L-*threo*-isomer is a potent inhibitor of PKC and as such is capable of reversing multi-drug resistance in cancer cells (9). **See Literature References on page 106.**




1807 1807-025	L-threo-Dihydrosphingosine (Safingol) L- <i>threo</i> -Sphinganine, C18 chain	5 mg 25 mg	240.00 725.00
	$C_{18}H_{39}NO_2$ Source: synthetic Appearance: solid Storage: -20°C Activity: Inhibitor of Protein Kinase C (PKC) and Sphingosine Kinase	Mol. Wt.: 301 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO Melting Point (°C): 103-114	CAS#: 15639-50-6

1831	D-erythro-Dihydrosphingosine D- <i>erythro</i> -Sphinganine, C18 chain	25 mg	200.00
	$C_{18}H_{39}NO_2$ Source: synthetic Appearance: solid Storage: -20°C Activity: Inhibitor of PLA ₂ and PLD	Mol. Wt.: 301 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO	CAS#: 764-22-7 Identity: confirmed by MS

1846	L-erythro-Dihydrosphingosine L- <i>erythro</i> -Sphinganine, C18 chain	1 mg	155.00
	$C_{18}H_{39}NO_2$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 301 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO	

1851	D-threo-Dihydrosphingosine D-threo-Sphinganine, C18 chain	1 mg	235.00
	C ₁₈ H ₃₉ NO ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 301 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO	CAS#: 6036-86-8 Identity: confirmed by MS
1324	D,L-erythro-Dihydrosphingosine D,L-erythro-Sphinganine, C18 chain	25 mg	150.00
	C ₁₈ H ₃₉ NO ₂ Source: synthetic Appearance: solid Storage: -20°C Activity: Inhibitor of sphingosine kinase	Mol. Wt.: 301 Purity: erythro 77%; threo 23% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO	CAS#: 3102-56-5 Identity: confirmed by MS
1326	D,L-C16-Dihydrosphingosine (mixed isomers) D,L-Sphinganine with C16 chain	10 mg	145.00
	C ₁₆ H ₃₅ NO ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 273 Purity: erythro 90%, threo 10% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO	
1845	D-erythro-C20-Dihydrosphingosine D-erythro-Sphinganine, C20 chain	5 mg	205.00
	C ₂₀ H ₄₃ NO ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 330 Purity: 98+% by TLC, GC Solubility: warm ethanol, chloroform/methanol, 5:1	CAS#: 24006-62-0
1839	D,L-erythro-C20-Dihydrosphingosine D,L-erythro-Sphinganine, C20 chain	10 mg	145.00
	C ₂₀ H ₄₃ NO ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 330 Purity: 98+% by TLC, GC Solubility: warm ethanol, chloroform/methanol, 5:1	Identity: confirmed by MS


3-Keto-Dihydrosphingosines



1876	3-keto-Dihydrosphingosine-HCl 3-keto-Sphinganine hydrochloride	10 mg	515.00
	C ₁₈ H ₃₇ NO ₂ •HCl Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 299 + HCl Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	CAS#: 18944-28-0 Identity: confirmed by MS

1891	3-keto-C6-Dihydrosphingosine-HCl 1-Hydroxy-2-amino-3-keto-hexane • HCl	10 mg	495.00
	C ₆ H ₁₃ NO ₂ • HCl Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 168 Purity: 98+% by TLC Solubility: ethanol, methanol, DI water	CAS#: 1314999-30-8
1892	3-keto-C8-Dihydrosphingosine-HCl 1-Hydroxy-2-amino-3-keto-octane • HCl	10 mg	495.00
	C ₈ H ₁₇ NO ₂ • HCl Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 196 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol, DI water	CAS#: 1824382-78-6
1893	3-keto-C12-Dihydrosphingosine-HCl 1-Hydroxy-2-amino-3-keto-dodecane • HCl	10 mg	495.00
	C ₁₂ H ₂₅ NO ₂ • HCl Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 252 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	CAS#: 1823032-02-5


Phytosphingosines



1330	Phytosphingosine 4-Hydroxysphinganine	50 mg	155.00
	C ₁₈ H ₃₉ NO ₃ Source: natural, yeast (<i>Pichia ciferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 318 Purity: 98+% by TLC, GC Solubility: ethanol, methanol, chloroform/methanol, 2:1 (warm)	CAS#: 554-62-1 Identity: confirmed by MS

Other Sphingosine Derivatives and Precursors

1320	N,N-Dimethyl-D-erythro-sphingosine	5 mg/ml, 1 ml	135.00
	C ₂₀ H ₄₁ NO ₂ Source: synthetic Appearance: liquid Storage: -20°C Activity: Inhibitor of phosphokinase C	Mol. Wt.: 328 Purity: 98+% by TLC Solubility: chloroform, ethanol, isopropanol, methanol Solvent: isopropanol	CAS#: 119567-63-4 Identity: confirmed by MS



1896	N,N-Dihexyl-D-erythro-sphingosine Sphingosine with tertiary amine group	5 mg/ml, 1 ml	225.00
	C₃₀H₆₁NO₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 468 Purity: 95% by TLC Solubility: chloroform, ethanol, methanol Solvent: ethanol	

Ceramides

Ceramide is a fatty acid amide of sphingosine. It may be formed by dehydrogenation of dihydroceramide; by hydrolysis of sphingomyelin or glycosphingolipids; or by acylation of free sphingosine. Ceramide functions as a precursor in the synthesis of sphingomyelin (by an exchange reaction with phosphatidylcholine and phosphatidylethanolamine); of glycosphingolipids (by glycosylation with UDP-hexose); and of free sphingosine and fatty acid by hydrolysis. The sphingosine can be phosphorylated by a kinase to form sphingosine-1-phosphate, which may undergo further hydrolysis or cleavage.

Control of sphingolipid metabolism maintains vital balance points in cell physiology. Two of ceramide's metabolites, sphingosine-1-phosphate and glucosylceramide, produce cell proliferation. Sphingosine-1-phosphate is also a highly active regulator of angiogenesis, vascular maturation, cardiac development, immunity, and directed cell movement. Sphingosine, the free base, is a potent inhibitor of protein kinase C and is involved in intracellular calcium regulation.

Sphingolipid enzymes seem to be particularly active in cancers, so modifying their activities by exogenous action may provide alternatives to chemical therapies. These enzymes are controlled by many known agents, such as 1,25-dihydroxy-vitamin D₃, tumor necrosis factor- α , nerve growth factor, interleukin 1, endothelial growth factor, glutathione, arachidonic acid, dexamethasone, many anticancer drugs, therapeutic radiation, and activators of the FAS receptor.


Ceramide exerts numerous biological effects, including induction of cell maturation, cell cycle arrest, terminal cell differentiation, cell senescence, and cell death. Other effects include producing reactive oxygen in mitochondria (followed by apoptosis) and stimulating phosphorylation of certain proteins (especially mitogen activated protein). It also stimulates some protein phosphatases (especially protein phosphatase 2A). Thus ceramide is an important controller of protein activity.

It is apparent from these relationships that ceramide exists at the crux of several enzyme reaction cycles and that experiments involving sphingolipid function call for control of all of the cycles and their branch-off points. Matreya is the major supplier of these lipids, which can be used as standards for analysis of tissues (a much needed part of modern research) and identification of major sphingolipids.

Ceramides with short side chains have been shown to enter easily into cells where they are biologically active. Ceramides with longer side chains, however, also enter cells if dissolved in dodecane-isopropanol first. Fluorescent labeled ceramides and sphingomyelin made from fluorescent labeled acids instead of plain fatty acids are also available for the study of the localization and metabolism of sphingolipids in the cell.

In three major reviews, Radin (10-12) has discussed the biochemistry and chemistry of ceramide and outlined many potential approaches to cancer therapy using ceramides and related compounds as generators of apoptosis. **See Literature References on page 106.**

Synthetic Ceramides Derived from C18-Sphingosine



1901	N-Acetyl-D-erythro-sphingosine N-C2:0-D-erythro-Ceramide	10 mg	145.00
	C₂₀H₃₉NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 342 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO (up to 5 mg/ml)	CAS#: 3102-57-6 Identity: confirmed by MS
1829	N-Acetyl-L-threo-sphingosine N-C2:0-L-threo-Ceramide	1 mg	165.00
	C₂₀H₃₉NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 342 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO, DMF (up to 5 mg/ml)	
1847	N-Acetyl-L-erythro-sphingosine N-C2:0-L-erythro-Ceramide	1 mg	150.00
	C₂₀H₃₉NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 342 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO, DMF (up to 5 mg/ml)	
1900	N-Hexanoyl-D-erythro-sphingosine N-C6:0-D-erythro-Ceramide	10 mg	145.00
	C₂₄H₄₇NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 398 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO (up to 5 mg/ml)	CAS#: 124753-97-5 Identity: confirmed by MS
1828	N-Hexanoyl-L-threo-sphingosine N-C6:0-L-threo-Ceramide	1 mg	190.00
	C₂₄H₄₇NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 398 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml)	Identity: confirmed by MS
1848	N-Hexanoyl-L-erythro-sphingosine N-C6:0-L-erythro-Ceramide	1 mg	170.00
	C₂₄H₄₇NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 398 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml)	Identity: confirmed by MS

1809	N-Hexanoyl-D-threo-sphingosine N-C6:0-D-threo-Ceramide	1 mg	190.00
	$C_{24}H_{47}NO_3$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 398 Purity: 98+% by TLC, GC Solubility: chloroform, methanol, DMSO (up to 5 mg/ml)	Identity: confirmed by MS
1903	N-Octanoyl-D-erythro-sphingosine N-C8:0-D-erythro-Ceramide	10 mg	145.00
	$C_{26}H_{51}NO_3$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 426 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO (up to 5 mg/ml)	CAS#: 74713-59-0 Identity: confirmed by MS
1830	N-Octanoyl-L-threo-sphingosine N-C8:0-L-threo-Ceramide	1 mg	190.00
	$C_{26}H_{51}NO_3$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 426 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, DMF (up to 5 mg/ml)	Identity: confirmed by MS
1810	N-Octanoyl-D-threo-sphingosine N-C8:0-D-threo-Ceramide	1 mg	190.00
	$C_{26}H_{51}NO_3$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 426 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, DMF (up to 5 mg/ml)	
1333	N-Decanoyl-D-erythro-sphingosine N-C10:0-D-erythro-Ceramide	10 mg	125.00
	$C_{28}H_{55}NO_3$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 454 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO (up to 5mg/ml)	CAS#: 111122-57-7 Identity: confirmed by MS
1936	N-Dodecanoyl-D-erythro-sphingosine N-C12:0-D-erythro-Ceramide	10 mg	125.00
	$C_{30}H_{59}NO_3$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 482 Purity: 98+% by TLC Solubility: chloroform, ethanol, DMSO, DMF	CAS#: 74713-60-3 Identity: confirmed by MS
2037	N-Pentadecanoyl-D-erythro-sphingosine N-C15:0-D-erythro-Ceramide	10 mg	140.00
	$C_{33}H_{65}NO_3$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 524 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, warm methanol	Identity: confirmed by MS

1915	N-Hexadecanoyl-D-erythro-sphingosine N-C16:0-D-erythro-Ceramide; N-Palmitoyl-D-erythro-sphingosine	10 mg	135.00
	$C_{34}H_{67}NO_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 538 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, warm methanol	CAS#: 24696-26-2 Identity: confirmed by MS
2038	N-Heptadecanoyl-D-erythro-sphingosine N-C17:0-D-erythro-Ceramide	10 mg	140.00
	$C_{35}H_{69}NO_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 552 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, warm methanol	CAS#: 67492-16-4
1832	N-Octadecanoyl-D-erythro-sphingosine N-C18:0-D-erythro-Ceramide; N-Stearoyl-D-erythro-sphingosine	10 mg	130.00
	$C_{36}H_{71}NO_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 566 Purity: 98+% by TLC, GC Solubility: chloroform, hot ethanol, chloroform/methanol, 2:1 (up to 5mg/ml)	CAS#: 2304-81-6
2201	N-omega-CD₃-Octadecanoyl-D-erythro-sphingosine N-C18:0-CD ₃ -D-erythro-Ceramide; N-Stearoyl-CD ₃ -D-erythro-sphingosine	1 mg	350.00
	$C_{36}H_{68}NO_3D_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 569 Purity: 98+% by TLC, GC Solubility: chloroform, hot ethanol, DMF	Identity: confirmed by MS
1939 1939-25	N-Octadecenoyl-(cis-9)-D-erythro-sphingosine N-C18:1-D-erythro-Ceramide; N-Oleoyl-D-erythro-sphingosine	5 mg 25 mg	110.00 415.00
	$C_{36}H_{69}NO_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 564 Purity: 98+% by TLC Solubility: chloroform, hot ethanol, DMF	CAS#: 5966-28-9 Identity: confirmed by MS
1843	N-Octadecanoyl-L-threo-sphingosine N-C18:0-L-threo-Ceramide; N-Stearoyl-L-threo-sphingosine	1 mg	170.00
	$C_{36}H_{71}NO_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 566 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml)	
1850	N-Octadecanoyl-L-erythro-sphingosine N-C18:0-L-erythro-Ceramide; N-Stearoyl-L-erythro-sphingosine	1 mg	170.00
	$C_{36}H_{71}NO_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 566 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml)	
1855	N-Octadecanoyl-D-threo-sphingosine N-C18:0-D-threo-Ceramide; N-Stearoyl-D-threo-sphingosine	1 mg	190.00
	$C_{36}H_{71}NO_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 566 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml)	CAS#: 2304-81-6

2039	N-Nonadecanoyl-D-erythro-sphingosine N-C19:0-D-erythro-Ceramide	10 mg	160.00
	C₃₇H₇₃NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 580 Purity: 98+% by TLC, GC, HPLC Solubility: chloroform, warm ethanol, warm methanol	Identity: confirmed by MS
1916	N-Tetracosanoyl-D-erythro-sphingosine N-C24:0-D-erythro-Ceramide; N-Lignoceroyl-D-erythro-sphingosine	5 mg	150.00
	C₄₂H₈₃NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 650 Purity: 98+% by TLC, GC Solubility: chloroform	CAS#: 34435-05-7 Identity: confirmed by MS
1930	N-Tetracosenoyl-(cis-15)-D-erythro-sphingosine N-cis-15-C24:1-D-erythro-Ceramide; N-Nervonoyl-D-erythro-sphingosine	5 mg	160.00
	C₄₂H₈₁NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 648 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, warm methanol	CAS#: 54164-50-0
2049	N-Triacontanoyl-D-erythro-sphingosine N-C30:0-D-erythro-Ceramide	1 mg	350.00
	C₄₈H₉₅NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 734 Purity: 98+% by TLC, GC Solubility: chloroform/methanol 2:1, chloroform, ethanol	Identity: confirmed by MS
2080	N-omega-Hydroxytriacontanoyl-D-erythro-sphingosine N-omega-Hydroxy-C30:0-D-erythro-ceramide	5 mg	470.00
	C₄₈H₉₅NO₄ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 750 Purity: 98+% by TLC, GC Solubility: chloroform/methanol 2:1, DMF, hot ethanol	Identity: confirmed by MS
2084	N-(30-Linoleoyloxy-triacontanoyl)-sphingosine Ceramide (EOS); EOS Ceramide 1	1 mg	595.00
	C₆₆H₁₂₅NO₅ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 1013 Purity: 98+% by TLC Solubility: chloroform, methanol, DMF	Identity: confirmed by MS
2208 *NEW*	N-(32-Linoleoyloxy-dotriacontanoyl)-sphingosine-D9 EOS Ceramide, deuterated; O-acylceramide, deuterated	1 mg	550.00
	C₆₈H₁₂₀D₉NO₅ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 1050 Purity: 98+% by TLC Solubility: chloroform, methanol, DMF	Identity: confirmed by MS
2048	N-Dotriacontanoyl-D-erythro-sphingosine N-C32:0-D-erythro-Ceramide	5 mg	220.00
	C₅₀H₉₉NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 762 Purity: 98+% by TLC, GC Solubility: chloroform/methanol 2:1, hot methanol, hot ethanol, hot DMSO	Identity: confirmed by MS

2081	N-Hexanoyl-biotin-D-erythro-sphingosine N-C6:0-biotin-D-erythro-Ceramide	5 mg	540.00
	C ₃₄ H ₆₂ N ₄ O ₅ S Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 639 Purity: 98+% by TLC, HPLC Solubility: chloroform/methanol 2:1, DMF	Identity: confirmed by MS


2-Hydroxy Ceramides

2044	N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-sphingosine N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-ceramide; N-(R,S)-alpha-Hydroxystearoyl-D-erythro-sphingosine	5 mg	240.00
	C ₃₆ H ₇₁ NO ₄ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 582 Purity: 98+% by TLC, GC Solubility: chloroform/methanol/DI water, 2:1:0.5	Identity: confirmed by MS

Ceramide Made from Sphingosines with Sphingoid Bases Other Than C18

1842	N-Acetyl-D-erythro-sphingosine (C14 sphingolipid base) N-C2:0 Ceramide of D-erythro-C14-sphingosine	5 mg	285.00
	C ₁₆ H ₃₁ NO ₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 285 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO, DMF (up to 5 mg/ml)	Identity: confirmed by MS
2077	N-Hexadecanoyl-D-erythro-sphingosine (C16 sphingolipid base) N-Palmitoyl-D-erythro-C16-sphingosine; N-C16:0 Ceramide of D-erythro-C16-sphingosine	1 mg	365.00
	C ₃₂ H ₆₃ NO ₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 510 Purity: 98+% by TLC, HPLC Solubility: chloroform, warm ethanol, warm methanol	Identity: confirmed by MS

Dihydroceramides



1834	N-Acetyl-D-erythro-dihydrosphingosine N-C2:0-D-erythro-Dihydroceramide; N-Acetyl-D-erythro-sphinganine	5 mg	90.00
	C ₂₀ H ₄₁ NO ₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 344 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol	CAS#: 13031-64-6 Identity: confirmed by MS
1910	N-Hexanoyl-D-erythro-dihydrosphingosine N-C6:0-D-erythro-Dihydroceramide; N-Hexanoyl-D-erythro-sphinganine	5 mg	90.00
	C ₂₄ H ₄₉ NO ₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 400 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, DMSO	

1854	N-Octanoyl-D-erythro-dihydrosphingosine N-C8:0-D-erythro-Dihydroceramide; N-Octanoyl-D-erythro-sphinganine	5 mg	90.00
	C₂₆H₅₃NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 428 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, DMSO	Identity: confirmed by MS
2078	N-Hexadecanoyl-D-erythro-dihydrosphingosine N-C16:0-D-erythro-Dihydroceramide; N-Hexadecanoyl-D-erythro-sphinganine	10 mg	150.00
	C₃₄H₆₉NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 540 Purity: 98+% by TLC Solubility: chloroform/methanol 5:1, hot ethanol, DMSO	Identity: confirmed by MS
2083	N-Heptadecanoyl-D-erythro-dihydrosphingosine N-C17:0-D-erythro-Dihydroceramide; N-Heptadecanoyl-D-erythro-sphinganine	5 mg	150.00
	C₃₅H₇₁NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 554 Purity: 98+% by TLC, HPLC Solubility: chloroform/methanol 5:1, hot ethanol, DMSO	Identity: confirmed by MS
2041	N-Octadecanoyl-D-erythro-dihydrosphingosine N-C18:0-D-erythro-Dihydroceramide; N-Octadecanoyl-D-erythro-sphinganine; N-Stearoyl-D-erythro-dihydrosphingosine	10 mg	150.00
	C₃₆H₇₃NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 568 Purity: 98+% by TLC, HPLC Solubility: hot ethanol, DMSO, warm chloroform/methanol, 5:1	Identity: confirmed by MS
2093 *NEW*	N-Tetracosanoyl-D-erythro-dihydrosphingosine N-C24:0-D-erythro-dihydroceramide; N-Tetracosanoyl-D-erythro-sphinganine; N-Lignoceryl-D-erythro-dihydrosphingosine	5 mg	325.00
	C₄₂H₈₅NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 652 Purity: 98+% by TLC, HPLC Solubility: chloroform/methanol, 2:1	Identity: confirmed by MS
2202	N-omega-CD₃-Octadecanoyl-D-erythro-dihydrosphingosine N-C18:0-CD ₃ -D-erythro-dihydroceramide; N-Stearoyl-CD ₃ -D-erythro-sphinganine	1 mg	275.00
	C₃₆H₇₀D₃NO₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 571 Purity: 98% by TLC, GC, HPLC Solubility: hot ethanol, DMF, DMSO, chloroform/methanol, 2:1	Identity: confirmed by MS


2-Hydroxy Dihydroceramides

2043	N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-dihydrosphingosine N-(R,S)-alpha-Hydroxy-C12:0-D-erythro-dihydroceramide	5 mg	260.00
	C₃₀H₆₁NO₄ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 500 Purity: 98+% by TLC, GC Solubility: chloroform/methanol/DI water, 2:1:0.5	

2047	N-(R,S)-<i>alpha</i>-Hydroxyhexadecanoyl-D-erythro-dihydrosphingosine N-(R,S)- <i>alpha</i> -Hydroxy-C16:0-D-erythro-dihydroceramide; N-(R,S)- <i>alpha</i> -Hydroxypalmitoyl-D-erythro-dihydrosphingosine	5 mg	160.00
	C₃₄H₆₉NO₄ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 556 Purity: 98+% by TLC, GC Solubility: chloroform/methanol/DI water, 2:1:0.5	Identity: confirmed by MS
2045	N-(R,S)-<i>alpha</i>-Hydroxyoctadecanoyl-D-erythro-dihydrosphingosine N-(R,S)- <i>alpha</i> -Hydroxy-C18:0-D-erythro-dihydroceramide; N-(R,S)- <i>alpha</i> -Hydroxystearoyl-D-erythro-dihydrosphingosine	5 mg	260.00
	C₃₆H₇₃NO₄ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 584 Purity: 98+% by TLC, GC Solubility: chloroform/methanol/DI water, 2:1:0.5	Identity: confirmed by MS


Ceramides From Natural Sources

General ceramide structure



1056	Ceramides (mixture) Ceramides with hydroxy and non-hydroxy acyl groups	25 mg	145.00
	C₄₂H₈₃NO₄ Source: natural, bovine Appearance: solid Storage: -20°C See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 666 (2-hydroxylignoceroyl) Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	CAS#: 104404-17-13
1322	Ceramides (non-hydroxy) Ceramides with mostly non-hydroxy acyl groups	10 mg	110.00
1322-05		50 mg	360.00
	C₃₆H₇₁NO₃ Source: natural, bovine Appearance: solid Storage: -20°C See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 566 (stearoyl) Purity: 98+% by TLC Solubility: ethanol, chloroform/methanol, 2:1	
1323	Ceramides (hydroxy) Ceramides with mostly hydroxy acyl groups	10 mg	110.00
1323-05		50 mg	360.00
	C₃₆H₇₁NO₄ Source: natural, bovine Appearance: solid Storage: -20°C See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 582 (2-hydroxystearoyl) Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	Identity: confirmed by MS

Phytoceramides




1897	N-Acetyl-phytosphingosine N-C2:0-Phytoceramide	5 mg	175.00
	C₂₀H₄₁NO₄ Source: semisynthetic, yeast (<i>Pichia ciferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 360 Purity: 98+% by TLC, HPLC Solubility: ethanol, methanol, warm DMSO, chloroform/methanol, 1:1 (warm)	CAS#: 475995-69-8 Identity: confirmed by MS
1895	N-Hexanoyl-phytosphingosine N-C6:0-Phytoceramide	5 mg	175.00
	C₂₄H₄₉NO₄ Source: semisynthetic, yeast (<i>Pichia ciferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 416 Purity: 98+% by TLC, HPLC Solubility: ethanol, methanol, chloroform/methanol, 1:1 (warm)	Identity: confirmed by MS
1894	N-Octanoyl-phytosphingosine N-C8:0-Phytoceramide	5 mg	175.00
	C₂₆H₅₃NO₄ Source: semisynthetic, yeast (<i>Pichia ciferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 444 Purity: 98+% by TLC, HPLC Solubility: ethanol, methanol, chloroform/methanol, 1:1 (warm)	CAS#: 475995-74-5 Identity: confirmed by MS
2035	N-Hexadecanoyl-phytosphingosine N-C16:0-Phytoceramide; N-Palmitoyl-phytosphingosine	5 mg	165.00
	C₃₄H₆₉NO₄ Source: semisynthetic, yeast (<i>Pichia ciferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 556 Purity: 98+% by TLC, HPLC Solubility: chloroform/methanol, 5:1	Identity: confirmed by MS
2034	N-Octadecanoyl-phytosphingosine N-C18:0-Phytoceramide; N-Stearoyl-phytosphingosine	5 mg	165.00
	C₃₆H₇₃NO₄ Source: semisynthetic, yeast (<i>Pichia ciferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 584 Purity: 98+% by TLC, HPLC Solubility: chloroform/methanol, 1:1 (warm)	CAS#: 34354-88-6 Identity: confirmed by MS
2036	N-Tetracosanoyl-phytosphingosine N-C24:0-Phytoceramide; N-Lignoceroyl-phytosphingosine	5 mg	175.00
	C₄₂H₈₅NO₄ Source: semisynthetic, yeast (<i>Pichia ciferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 668 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1	Identity: confirmed by MS

2210 *NEW*	N-<i>omega</i>-CD₃-Octadecanoyl-phytosphingosine N-C18:0-CD ₃ -Phytoceramide; N-Stearoyl-CD ₃ -phytosphingosine	1 mg	275.00
	C₃₆H₇₀D₃NO₄ Source: semisynthetic, yeast (<i>Pichia cifferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 587 Purity: 98+% by TLC, HPLC Solubility: chloroform, DMF, DMSO	Identity: confirmed by MS

Fluorescent Ceramides

Absorption: 460 nm Emission: 535 nm

1841 1841-001	N-Hexanoyl-NBD-D-erythro-sphingosine N-C6:0-NBD-Ceramide; N-C6:0-NBD-D-erythro-Sphingosine	100 µg 1 mg	155.00 390.00
	C₃₀H₄₉N₅O₆ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 576 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol Melting Point (°C): 85.7-87.9	CAS#: 86701-10-2



1618 1618-001	N-Dodecanoyl-NBD-D-erythro-sphingosine N-C12:0-NBD-Ceramide; N-C12:0-NBD-D-erythro-Sphingosine	100 µg 1 mg	150.00 375.00
	C₃₆H₆₁N₅O₆ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 660 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	Identity: confirmed by MS

1857 1857-001	N-Hexanoyl-NBD-L-threo-sphingosine N-C6:0-NBD-Ceramide; N-C6:0-NBD-L-threo-Sphingosine	100 µg 1 mg	225.00 650.00
	C₃₀H₄₉N₅O₆ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 576 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	

1620	N-Dodecanoyl-NBD-L-threo-sphingosine N-C12:0-NBD-Ceramide; N-C12:0-NBD-L-threo-Sphingosine	100 µg	225.00
	C₃₆H₆₁N₅O₆ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 660 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	

1624	N-Hexanoyl-NBD-L-threo-dihydrosphingosine N-C6:0-NBD-Dihydroceramide; N-C6:0-NBD-L-threo-Dihydrosphingosine	100 µg	200.00
	C₃₀H₅₁N₅O₆ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 578 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	


1623	N-Dodecanoyl-NBD-L-<i>threo</i>-dihydrosphingosine N-C12:0-NBD-Dihydroceramide; N-C12:0-NBD-L- <i>threo</i> -Dihydrosphingosine	100 µg	200.00
	$C_{36}H_{63}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 662 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	
1626	N-Hexanoyl-NBD-D-<i>erythro</i>-dihydrosphingosine N-C6:0-NBD-Dihydroceramide; N-C6:0-NBD-D- <i>erythro</i> -Dihydrosphingosine	100 µg	200.00
	$C_{30}H_{51}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 578 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	Identity: confirmed by MS
1625	N-Dodecanoyl-NBD-D-<i>erythro</i>-dihydrosphingosine N-C12:0-NBD-Dihydroceramide; N-C12:0-NBD-D- <i>erythro</i> -Dihydrosphingosine	100 µg	200.00
	$C_{36}H_{63}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 662 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	
1628	N-Hexanoyl-NBD-phytosphingosine N-C6:0-NBD-Phytoceramide; N-C6:0-NBD-Phytosphingosine	100 µg	200.00
	$C_{30}H_{51}N_5O_7$ Source: semisynthetic, bacteria Appearance: solid Storage: -20°C	Mol. Wt.: 594 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	CAS#: 477239-93-3
1627	N-Dodecanoyl-NBD-phytosphingosine N-C12:0-NBD-Phytoceramide; N-C12:0-NBD-Phytosphingosine	100 µg	185.00
	$C_{36}H_{63}N_5O_7$ Source: semisynthetic, bacteria Appearance: solid Storage: -20°C	Mol. Wt.: 678 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	CAS#: 388566-94-7

See Labeled Standards section (page 96) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical Service for more information. 814-355-1030

Phosphosphingolipids


Sphingomyelins



1051 1051-1	Sphingomyelin, bovine SPM; Ceramide-1-phosphorylcholine	25 mg 1 g	70.00 625.00
	C₄₁H₈₃N₂O₆P Source: natural, bovine Appearance: solid Storage: -20°C See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 731 (stearoyl) Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	CAS#: 85187-10-6 Identity: confirmed by MS
1328	Sphingomyelin, porcine RBC SPM; Ceramide-1-phosphorylcholine	25 mg	85.00
	C₄₇H₉₅N₂O₆P Source: natural, porcine RBC Appearance: solid Storage: -20°C See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 815 (lignoceroyl) Purity: 98+% by TLC Solubility: chloroform, ethanol	CAS#: 85187-10-6 Identity: confirmed by MS
1329	Sphingomyelin, bovine buttermilk SPM; Ceramide-1-phosphorylcholine	25 mg	70.00
	C₄₆H₉₃N₂O₆P Source: natural, bovine buttermilk Appearance: solid Storage: -20°C See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 801 (tricosanoyl) Purity: 98+% by TLC Solubility: chloroform, ethanol	CAS#: 85187-10-6 Identity: confirmed by MS
1332	Sphingomyelin, egg SPM; Ceramide-1-phosphorylcholine	25 mg	90.00
	C₃₉H₇₉N₂O₆P Source: natural, chicken egg Appearance: solid Storage: -20°C See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 703 (palmitoyl) Purity: 98+% by TLC Solubility: chloroform, methanol, warm ethanol	Identity: confirmed by MS
1907	N-Acetyl-sphingosylphosphorylcholine N-C2:0-Sphingomyelin	5 mg	250.00
	C₂₅H₅₁N₂O₆P Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C Note: Mixture of D-erythro and L-threo isomers	Mol. Wt.: 506 Purity: 98+% by TLC Solubility: ethanol, chloroform/methanol, 2:1	


1890	N-Heptadecanoyl-sphingosylphosphorylcholine N-C17:0-Sphingomyelin	5 mg	250.00
	$C_{40}H_{81}N_2O_6P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: $-20^{\circ}C$ Note: Mixture of <i>D-erythro</i> and <i>L-threo</i> isomers	Mol. Wt.: 717 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	
1911	N-Octadecanoyl-sphingosylphosphorylcholine N-C18:0-Sphingomyelin; N-Stearoyl-sphingosylphosphorylcholine	5 mg	250.00
	$C_{41}H_{83}N_2O_6P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: $-20^{\circ}C$ Note: Mixture of <i>D-erythro</i> and <i>L-threo</i> isomers	Mol. Wt.: 731 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	Identity: confirmed by MS
1917	N-Eicosanoyl-D-erythro-sphingosylphosphorylcholine N-C20:0-Sphingomyelin	500 μ g	260.00
	$C_{43}H_{87}N_2O_6P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 759 Purity: 98+% by TLC Solubility: ethanol, methanol, chloroform/methanol, 14:1	
1918	N-Docosanoyl-D-erythro-sphingosylphosphorylcholine N-C22:0-Sphingomyelin	500 μ g	260.00
	$C_{45}H_{91}N_2O_6P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 787 Purity: 98+% by TLC Solubility: ethanol, methanol, chloroform/methanol, 14:1	Identity: confirmed by MS
2200	N-1-¹³C-Hexadecanoyl-D-erythro-sphingosylphosphorylcholine <i>D-erythro</i> -Sphingomyelin with 1- ¹³ C-palmitic acid; N-1- ¹³ C-Palmitoyl-sphingosylphosphorylcholine	1 mg	260.00
	$^{12}C_{38}^{13}CH_{79}N_2O_6P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 703 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	
1327	N-Acyl-D-erythro-sphingosylphosphorylethanolamine Ceramide phosphorylethanolamine	5 mg	210.00
	$C_{43}H_{87}N_2O_6P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 773 (tricosanoyl) Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	Identity: confirmed by MS

Sphingosylphosphorylcholines (SPC)



1318	D-erythro-Sphingosylphosphorylcholine <i>D-erythro</i> -SPC	5 mg	330.00
	C₂₃H₄₉N₂O₅P Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 465 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	CAS#: 1670-26-4 Identity: confirmed by MS
1319	L-threo-Sphingosylphosphorylcholine <i>L-threo</i> -SPC	5 mg	350.00
	C₂₃H₄₉N₂O₅P Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 465 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	Identity: confirmed by MS
1321 1321-05	Sphingosylphosphorylcholine <i>lyso</i> -Sphingomyelin; SPC (mixture of <i>D-erythro</i> and <i>L-threo</i> isomers)	10 mg 50 mg	320.00 1,275.00
	C₂₃H₄₉N₂O₅P Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 465 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	CAS#: 82970-80-7
1913	<i>lyso</i>-Dihydro sphingomyelin Dihydro sphingosylphosphorylcholine (mixture of <i>D-erythro</i> and <i>L-threo</i> isomers)	1 mg	185.00
	C₂₃H₅₁N₂O₅P Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 467 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	

Sphingosine and Ceramide Phosphates




1803	D-erythro-Sphingosine-1-phosphate S-1-P	5 mg	385.00
	C₁₈H₃₈NO₅P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 380 Purity: 98+% by TLC Solubility: chloroform plus a few drops of TFA, chloroform/methanol/40% dimethylamine, 5:15:3, 1mg/ml	CAS#: 26993-30-6 Identity: confirmed by MS

1852	D-erythro-Dihydrosphingosine-1-phosphate	5 mg	390.00
	$C_{18}H_{40}NO_5P$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 382 Purity: 98+% by TLC Solubility: chloroform/methanol/40% dimethylamine, 5:15:3, 1mg/ml	CAS#: 19794-97-9
2046	N-Hexadecanoyl-D-erythro-sphingosine-1-phosphate (NH₄⁺ salt) N-C16:0-Ceramide-1-phosphate	5 mg	520.00
	$C_{34}H_{68}NO_6P \cdot NH_3$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 618 Purity: 98+% by TLC Solubility: chloroform/methanol/acetic acid, 60:15:25 chloroform/methanol/7.5M ammonium hydroxide 80:20:4	Identity: confirmed by MS
2206	N-omega-CD₃-Octadecanoyl-D-erythro-sphingosine-1-phosphate C18:0-CD ₃ -Ceramide-1-phosphate; N-Stearoyl-CD ₃ -C1P	1 mg	365.00
	$C_{36}H_{69}D_3NO_6P$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 649 Purity: 98+% by TLC Solubility: chloroform/methanol/acetic acid, 60:15:25 chloroform/methanol/7.5M ammonium hydroxide 80:20:4	Identity: confirmed by MS

Fluorescent Sphingomyelins

Absorption: 460 nm Emission: 535 nm



1912	N-Hexanoyl-NBD-sphingosylphosphorylcholine	100 µg	160.00
1912-001	N-C6:0-NBD-Sphingomyelin; N-C6:0-NBD-Sphingosylphosphorylcholine	1 mg	375.00
	$C_{35}H_{61}N_6O_9P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: $-20^{\circ}C$ Note: Mixture of <i>D-erythro</i> and <i>L-threo</i> isomers	Mol. Wt.: 740 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	CAS#: 94885-04-8
1619	N-Dodecanoyl-NBD-sphingosylphosphorylcholine	100 µg	200.00
1619-001	N-C12:0-NBD-Sphingomyelin; N-C12:0-NBD-Sphingosylphosphorylcholine	1 mg	395.00
	$C_{41}H_{73}N_6O_9P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: $-20^{\circ}C$ Note: Mixture of <i>D-erythro</i> and <i>L-threo</i> isomers	Mol. Wt.: 825 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 2:1	CAS#: 254117-01-6 Identity: confirmed by MS

See Labeled Standards section (page 96) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical Service for more information. 814-355-1030


Glycosphingolipids

Glycosphingolipids are widely distributed in animal and plant tissues. They consist of a ceramide (Cer) bound in glycosidic linkage through the primary hydroxyl to a mono- or oligosaccharide, which may contain substituents such as a sulfate, acetate, or phosphate group. They are amphiphilic and the less glycosylated compounds can be dispersed in buffers by dissolving them in a detergent or organic solvent (EtOH, DMSO, isoPrOH) and mixing by sonication.

Galactosphingolipids, mainly GalCer (cerebrosides) and its sulfate ester, occur in large amounts in the nervous system. Glucosphingolipids, the simplest of which is GlcCer (glucocerebrosides), are very widely distributed, particularly in nerve cell membranes. GlcCer is isolated from a variety of natural sources including human, bovine, and plant. Each of these sources has a heterogeneity in the fatty acid content of the ceramide as well as an occasional variation in the sphingoid chain. Globosides (containing both glucose and galactose) are a prominent group of glycosphingolipids, they contain an α -linked galactose moiety and are typically located in blood cell membranes. Gangliosides are another prominent group of glycosphingolipids; they are acidic because of substitution with sialic (neuraminic) acid. The glycosphingolipids function in a wide range of enzyme and structural interactions, such as immunological or membrane recognition phenomena, binding of microbial pathogens, hormone and growth factor actions, cancer cell growth and malignancy, atherosclerosis, genetic disease errors, blood group determinants, etc. Tissues change in glycosphingolipid composition during embryogenesis, maturation, aging, and other vital physiological processes. Some glycosphingolipids stimulate cell proliferation, others induce apoptosis, effects of great significance to cancer therapy and maturational development. Marked differences in glycosphingolipid composition are seen in normal and cancerous cells. See references (13-25).

See Literature References on page 106.

Galactosylceramides




1050	Cerebrosides, bovine Galactosylceramide; Ceramide <i>beta</i> -D-galactoside	50 mg	125.00
	<p>$C_{48}H_{93}NO_9$ Source: natural, bovine Appearance: solid Storage: $-20^{\circ}C$ See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.</p>	<p>Mol. Wt.: 828 (2-hydroxytetraacosanoyl) Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1</p>	<p>CAS#: 85305-88-0 Identity: confirmed by MS</p>
1066	Cerebroside; Kerasin (top spot) Galactosylceramide with mostly non-hydroxy fatty acid side chain	10 mg	170.00
	<p>$C_{48}H_{91}NO_8$ Source: natural, bovine Appearance: solid Storage: $-20^{\circ}C$ See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.</p>	<p>Mol. Wt.: 810 (nervonyl, [24:1]) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.5</p>	<p>CAS#: 536-13-0 Identity: confirmed by MS</p>
1138	Cerebroside; Phrenosin (bottom spot) Galactosylceramide with mostly 2-hydroxy fatty acid side chains	10 mg	170.00
	<p>$C_{42}H_{81}NO_9$ Source: natural, bovine Appearance: solid Storage: $-20^{\circ}C$ See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.</p>	<p>Mol. Wt.: 744 (2-hydroxystearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.5</p>	<p>CAS#: 37211-11-3</p>

1305	Psychosine (free amine form) <i>lyso</i> -Cerebroside; 1- <i>beta</i> -D-Galactosylsphingosine	10 mg	275.00
	$C_{24}H_{47}NO_7$ Source: semisynthetic, bovine Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 461 Purity: 98+% by TLC Solubility: ethanol, chloroform/methanol/DI water, 5:1:0.1	CAS#: 2238-90-6 Identity: confirmed by MS
2087	Psychosine, synthetic <i>lyso</i> -Cerebroside; 1- <i>beta</i> -D-Galactosylsphingosine	5 mg	250.00
	$C_{24}H_{47}NO_7$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 461 Purity: 98+% by TLC Solubility: ethanol, chloroform/methanol/DI water, 5:1:0.1	CAS#: 2238-90-6 Identity: confirmed by MS
2091 *NEW*	N-Glycinated galactosylsphingosine N-Glycinated cerebroside; N-Glycinated galactosylceramide; N-Glycinated psychosine	1 mg	325.00
	$C_{26}H_{50}N_2O_8$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 518 Purity: 98+% by TLC Solubility: chloroform/methanol, 80:20, ethanol	Identity: confirmed by MS
1325	N-Acetyl-psychosine N-C2:0-Cerebroside	10 mg	215.00
	$C_{26}H_{49}NO_8$ Source: semisynthetic, bovine Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 503 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	CAS#: 35823-61-1
1334	N-Octanoyl-<i>beta</i>-D-galactosylceramide N-C8:0-Galactosylceramide	10 mg	375.00
	$C_{32}H_{61}NO_8$ Source: semisynthetic, bovine Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 588 Purity: 98+% by TLC Solubility: ethanol, methanol, chloroform/methanol, 9:1	CAS#: 41613-16-5
1937 1937-50	N-Dodecanoyl-<i>beta</i>-D-galactosylceramide N-C12:0-Galactosylceramide; N-Dodecanoyl- <i>beta</i> -D-galactosylsphingosine	10 mg 50 mg	375.00 1,200.00
	$C_{36}H_{69}NO_8$ Source: semisynthetic, bovine Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 644 Purity: 98+% by TLC, HPLC Solubility: ethanol, methanol, chloroform/methanol, 9:1	Identity: confirmed by MS
1335	N-Pentadecanoyl-psychosine N-C15:0-Cerebroside	5 mg	155.00
	$C_{39}H_{75}NO_8$ Source: semisynthetic, bovine Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 686 Purity: 98+% by TLC Solubility: chloroform/ methanol, 2:1	

1914	N-Octadecanoyl-D₃₅-psychosine, (perdeuterated, C18:0 fatty acid) N-C18:0-D ₃₅ -Cerebroside, perdeuterated; N-Stearoyl-D ₃₅ -psychosine, perdeuterated	5 mg	420.00
	C₄₂H₄₆D₃₅NO₈ Source: semisynthetic, bovine Appearance: solid Storage: -20°C Activity: Deuterium labeled stearoyl side chain	Mol. Wt.: 763 Purity: 98+% by TLC Solubility: chloroform, hot ethanol, chloroform/methanol, 2:1	Identity: confirmed by MS
1621	N-Hexanoyl-NBD-galactosylceramide N-C6:0-NBD- <i>beta</i> -D-Galactosylsphingosine; N-C6:0-NBD-Cerebroside	100 µg	250.00
	C₃₆H₅₉N₅O₁₁ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 738 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 5:1	
1633 1633-001	N-Dodecanoyl-NBD-galactosylceramide N-C12:0-NBD- <i>beta</i> -D-Galactosylsphingosine; N-C12:0-NBD-Cerebroside	100 µg 1 mg	210.00 450.00
	C₄₂H₇₁N₅O₁₁ Source: semisynthetic, bovine Appearance: solid Storage: -20°C Absorption: 460 nm	Mol. Wt.: Purity: 98+% by TLC Solubility: chloroform, DMSO, chloroform/methanol, 2:1 Emission: 535 nm	CAS#: 474942-98-8 Identity: confirmed by MS
2204	Lissamine-rhodamine B-dodecanoyl-galactosylceramide Sulforhodamine B-C12:0 cerebroside	500 µg	390.00
	C₆₃H₉₉N₄O₁₄S₂ Source: semisynthetic, bovine Appearance: solid Storage: -20°C Absorption: 540 nm	Mol. Wt.: 1201 Purity: 98+% by TLC Solubility: chloroform/methanol 8:2, DMSO, DMF Emission: 565 nm	Identity: confirmed by MS
2203	N-Hexanoyl-biotin-galactosylceramide N-C6:0-Biotin- <i>beta</i> -D-galactosylsphingosine; N-C6:0-Biotin-cerebroside	5 mg	500.00
	C₄₀H₇₂N₄O₁₀S Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 801 Purity: 98+% by TLC, HPLC Solubility: chloroform/methanol 2:1, methanol, DMF	Identity: confirmed by MS
1946	N-(1-Adamantaneacetyl)-galactosylceramide N-(1-Adamantaneacetyl)-galactocerebroside	5 mg	420.00
	C₃₆H₆₃NO₈ Source: semisynthetic, bovine Appearance: solid Storage: -20°C Activity: Inhibitor of glucosylceramide, sulfatide, and globotriaosylceramide (Gb ₃) synthesis	Mol. Wt.: 638 Purity: 98+% by TLC Solubility: chloroform, methanol, chloroform/methanol 9:1	Identity: confirmed by MS

Glucosylceramides



1057	Glucocerebrosides, Gaucher's spleen	5 mg	285.00
1057-25	Glucosylceramide; Ceramide <i>beta</i> -D-glucoside	25 mg	1,200.00

$C_{48}H_{93}NO_8$
Source: natural, human
Appearance: solid
Storage: $-20^{\circ}C$
 See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.

Mol. Wt.: 812 (lignoceryl)
Purity: 98+% by TLC, GC, MS
Solubility: chloroform/methanol, 2:1


CAS#: 85305-87-9
Identity: confirmed by MS

1521	Glucocerebrosides, buttermilk	5 mg	200.00
1521-50	Glucosylceramide; Ceramide <i>beta</i> -D-glucoside	50 mg	1,500.00

$C_{46}H_{89}NO_8$
Source: natural, bovine buttermilk
Appearance: solid
Storage: $-20^{\circ}C$
 See Table III (pg. 101) for typical fatty acid content of products prepared from natural sources.

Mol. Wt.: 784 (docosanoyl)
Purity: 98+% by TLC
Solubility: chloroform/methanol, 2:1

Identity: confirmed by MS



1522	Glucocerebrosides, plant	5 mg	90.00
1522-100	Glucosylceramide; Ceramide <i>beta</i> -D-glucoside	100 mg	1,250.00

$C_{40}H_{75}NO_9$
Source: natural, plant
Appearance: solid
Storage: $-20^{\circ}C$
 Sphingoid backbone is >95% 4,8-sphingadiene (C18:2 t,t-4,8) and most of the fatty acids are of the 2-hydroxy type.
 See Table III (pg. 101) for typical fatty acid content of products prepared from natural sources.

Mol. Wt.: 714 (2-hydroxyhexadecanoyl)
Purity: 98+% by TLC
Solubility: chloroform/methanol, 2:1

CAS#: 497155-61-0
Identity: confirmed by MS

2086	Glucosylsphingosine, synthetic	5 mg	250.00
	<i>lyso</i> -Glucocerebroside; 1- <i>beta</i> -D-Glucosylsphingosine; Glucosylpsychosine		

$C_{24}H_{47}NO_7$
Source: synthetic
Appearance: solid
Storage: $-20^{\circ}C$

Mol. Wt.: 461
Purity: 98+% by TLC, HPLC
Solubility: ethanol, methanol, chloroform/methanol, 2:1

CAS#: 52050-17-6
Identity: confirmed by MS

2209	$^{13}C_6$-Glucosylsphingosine	1 mg	550.00
NEW	1-(<i>beta</i> -D-Glucosyl-1,2,3,4,5,6- $^{13}C_6$)-sphingosine; $^{13}C_6$ - <i>lyso</i> -glucocerebroside		


$C_{18}^{13}C_6H_{47}NO_7$
Source: synthetic
Appearance: solid
Storage: $-20^{\circ}C$

Mol. Wt.: 468
Purity: 98+% by TLC, HPLC
Solubility: ethanol, methanol, chloroform/methanol, 2:1

Identity: confirmed by MS

1622	N-Hexanoyl-NBD-glucosylceramide	100 µg	250.00
1622-001	N-C6:0-NBD- <i>beta</i> -D-Glucosylsphingosine; N-C6:0-NBD-Glucosylceramide	1 mg	400.00
	$C_{36}H_{59}N_5O_{11}$	Mol. Wt.: 738	
	Source: semisynthetic, bovine	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: methanol, chloroform/methanol, 5:1	
	Storage: -20°C		
	Absorption: 460 nm	Emission: 535 nm	
2085	N-Hexanoyl-biotin-glucosylceramide	5 mg	500.00
	N-C6:0-Biotin- <i>beta</i> -D-glucosylsphingosine; N-C6:0-Biotin-glucosylceramide		
	$C_{40}H_{72}N_4O_{10}S$	Mol. Wt.: 801	
	Source: semisynthetic, plant	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform/methanol 2:1, methanol, DMF	
	Storage: -20°C		
1945	N-(1-Adamantaneacetyl)-glucosylceramide	5 mg	420.00
	N-(1-Adamantaneacetyl)-glucocerebroside		
	$C_{36}H_{61}NO_8$	Mol. Wt.: 636	
	Source: semisynthetic, plant	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform, methanol, chloroform/methanol, 9:1	
	Storage: -20°C		
	Activity: Inhibitor of glucocerebrosidase and lactosylceramide synthase		

Sulfatides




1049	Sulfatides (Na⁺ salt), bovine	50 mg	520.00
	Ceramide-galactoside-3-sulfate; Cerebroside sulfate		
	$C_{42}H_{80}NNaO_{11}S$	Mol. Wt.: 830 (stearoyl) Na ⁺ salt	CAS#: 85496-63-5
	Source: natural, bovine	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: DMSO, chloroform/methanol/DI water, 2:1:0.1 (if needed, a few drops of acetic acid)	
	Storage: -20°C		
	See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.		
1904	<i>lyso</i>-Sulfatide (NH₄⁺ salt)	1 mg	565.00
	Sphingosine-1-galactoside-3-sulfate		
	$C_{24}H_{47}NO_{10}S \cdot NH_3$	Mol. Wt.: 542	CAS#: 38621-58-8
	Source: semisynthetic, bovine	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform/methanol, 2:1	
	Storage: -20°C		
2092	N-Glycinated <i>lyso</i>-sulfatide	1 mg	450.00
NEW	N-Glycinated sphingosine-1-galactoside-3-sulfate		
	$C_{26}H_{50}N_2O_{11}S$	Mol. Wt.: 599	
	Source: semisynthetic, bovine	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform/methanol/DI water, 70:30:4; ethanol	
	Storage: -20°C		

2076	N-Acetyl-sulfatide N-C2:0-Sulfatide; N-Acetyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfatide	1 mg	400.00
	$C_{26}H_{49}NO_{11}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 584 Purity: 98+% by TLC Solubility: ethanol, methanol, chloroform/methanol, 1:1	Identity: confirmed by MS
1938	N-Dodecanoyl-sulfatide N-C12:0-Sulfatide; N-Dodecanoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	400.00
	$C_{36}H_{69}NO_{11}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 724 Purity: 98+% by TLC Solubility: chloroform/methanol, 9:1, DMF	Identity: confirmed by MS
1875	N-Hexadecanoyl-sulfatide N-C16:0-Sulfatide; N-Palmitoyl-sulfatide; N-Palmitoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	400.00
	$C_{40}H_{77}NO_{11}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 780 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	Identity: confirmed by MS
1934	N-Heptadecanoyl-sulfatide N-C17:0-Sulfatide; N-Heptadecanoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	400.00
	$C_{41}H_{79}NO_{11}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 794 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1, DMSO, DMF	Identity: confirmed by MS
1932	N-Octadecanoyl-sulfatide N-C18:0-Sulfatide; N-Octadecanoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	400.00
	$C_{42}H_{81}NO_{11}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 808 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1	CAS#: 244215-65-4
1933	N-Octadecenoyl-(<i>cis</i>-9)-sulfatide N-C18:1-Sulfatide; N-Octadecenoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	400.00
	$C_{42}H_{79}NO_{11}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 806 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1	
1935	N-Nonadecanoyl-sulfatide N-C19:0-Sulfatide; N-Nonadecanoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	400.00
	$C_{43}H_{83}NO_{11}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 822 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1, DMSO, DMF	Identity: confirmed by MS

1888	N-Tetracosanoyl-sulfatide N-C24:0-Sulfatide; N-Tetracosanoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate; N-Lignoceroyl-sulfatide	1 mg	400.00
	C₄₈H₉₃NO₁₁S Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 892 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1	Identity: confirmed by MS
1931	N-Tetracosenoyl-(<i>cis</i>-15)-sulfatide N-Nervonyl-sulfatide; N-C24:1-Sulfatide; N-Tetracosenoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	400.00
	C₄₈H₉₁NO₁₁S Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 890 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1	Identity: confirmed by MS
1536	N-<i>omega</i>-CD₃-Octadecanoyl-sulfatide N-C18:0-CD ₃ -Sulfatide; N-Stearoyl-CD ₃ -sulfatide	1 mg	550.00
	C₄₂H₇₈D₃NO₁₁S Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 811 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS
1632	N-Dodecanoyl-NBD-sulfatide	100 µg	170.00
1632-001	N-C12:0-NBD-Sulfatide; N-Dodecanoyl-NBD- <i>lyso</i> -sulfatide; N-Dodecanoyl-NBD-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	450.00
	C₄₂H₇₁N₅O₁₄S Source: semisynthetic, bovine Appearance: solid Storage: -20°C Absorption: 460 nm	Mol. Wt.: 901 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1 Emission: 535 nm	
2207	N-Hexanoyl-biotin-sulfatide N-C6:0-Biotin-sulfatide; N-Hexanoyl-biotin-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	450.00
	C₄₀H₇₂N₄O₁₃S₂ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 881 Purity: 98+% by TLC Solubility: chloroform/methanol/water 2:1:0.1; methanol/water 9:1; DMF	Identity: confirmed by MS
1540 *NEW*	N-Octadecanoyl-sulfated-lactosylceramide SM3; N-Octadecanoyl-lactosylceramide-3'-sulfate; N-Octadecanoyl-lactosylceramide sulfatide	1 mg	350.00
	C₄₈H₉₁NO₁₆S Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 970 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS


Lactosylceramides



1500	Lactosylceramides LC; Lactocerebrosides; CDH; Ceramide <i>beta</i> -lactoside	1 mg	270.00
	C₄₈H₉₁NO₁₃ Source: natural, porcine RBC Appearance: solid Storage: -20°C See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 890 (stearoyl) Purity: 98+% by TLC Solubility: DMSO, chloroform/methanol/DI water, 5:1:0.1	CAS#: 4682-48-8 Identity: confirmed by MS
1507 1507-50	Lactosylceramides, bovine buttermilk LC; Lactocerebrosides; CDH; Ceramide <i>beta</i> -lactoside	5 mg 50 mg	300.00 2,050.00
	C₅₃H₁₀₁NO₁₃ Source: natural, bovine buttermilk Appearance: solid Storage: -20°C See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 960 (tricosanoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 5:1:0.1	CAS#: 4682-48-8 Identity: confirmed by MS
2088	<i>lyso</i>-Lactosylceramide, synthetic Lactosylsphingosine; 1- <i>beta</i> -lactosyl-sphing-4-enine; <i>lyso</i> -LC	1 mg	250.00
	C₃₀H₅₇NO₁₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 623 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1, DI water, DMSO	Identity: confirmed by MS
1517	<i>lyso</i>-Lactosylceramide Lactosylsphingosine; <i>lyso</i> -LC	1 mg	280.00
	C₃₀H₅₇NO₁₂ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 623 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS
2090 *NEW*	N-Glycinated lactosylsphingosine N-Glycinated <i>lyso</i> -lactosylceramide; N-Glycine 1- <i>beta</i> -lactosyl-sphing-4-enine	1 mg	415.00
	C₃₂H₆₀N₂O₁₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 681 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; DI water; DMSO	Identity: confirmed by MS
1532	N-Hexadecanoyl-lactosylceramide N-C16:0-Lactosylceramide; N-Palmitoyl-lactosylceramide	1 mg	235.00
	C₄₆H₈₇NO₁₃ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 862 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS

1534	N-<i>omega</i>-CD₃-Hexadecanoyl-lactosylceramide N-C16:0-CD ₃ -Lactosylceramide; N-Palmitoyl-CD ₃ -lactosylceramide	1 mg	460.00
	C₄₆H₈₄D₃NO₁₃ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 865 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 5:1:0.1	Identity: confirmed by MS
1538	N-Heptadecanoyl-lactosylceramide N-C17:0-Lactosylceramide; Lactosylceramide with C17:0 fatty acid side chain	1 mg	235.00
	C₄₇H₈₉NO₁₃ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 876 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	CAS#: 1354699-26-5 Identity: confirmed by MS
1629 1629-001	N-Hexanoyl-NBD-lactosylceramide N-Hexanoyl-NBD- <i>beta</i> -D-lactosylsphingosine; N-C6:0-NBD- <i>beta</i> -D-Lactosylsphingosine; N-C6:0-NBD-Lactosylceramide	50 ug 1 mg	265.00 500.00
	C₄₂H₆₉N₅O₁₆ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C Absorption: 460 nm	Mol. Wt.: 900 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1 Emission: 535 nm	
1630 1630-001	N-Dodecanoyl-NBD-lactosylceramide N-Dodecanoyl-NBD- <i>beta</i> -D-lactosylsphingosine; N-C12:0-NBD- <i>beta</i> -D-Lactosylsphingosine; N-C12:0-NBD-Lactosylceramide	50 ug 1 mg	295.00 500.00
	C₄₈H₈₁N₅O₁₆ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C Absorption: 460 nm	Mol. Wt.: 984 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1 Emission: 535 nm	
2205	N-Hexanoyl-biotin-lactosylceramide N-C6:0-Biotin- <i>beta</i> -D-lactosylceramide	1 mg	280.00
	C₄₆H₈₂N₄O₁₅S Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 963 Purity: 98+% by TLC Solubility: chloroform/methanol, 9:1, DMSO, DMF	Identity: confirmed by MS
1540 *NEW*	N-Octadecanoyl-sulfated-lactosylceramide SM3; N-Octadecanoyl-lactosylceramide-3'-sulfate; N-Octadecanoyl-lactosylceramide sulfatide	1 mg	350.00
	C₄₈H₉₁NO₁₆S Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 970 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS


Ceramide Trihexosides (Globotriaosylceramides)



1067 1067-10	Ceramide trihexosides CTH; Gb3; Globotriaosylceramide	1 mg 10 mg	375.00 2,850.00
	<p>$C_{60}H_{113}NO_{18}$ Mol. Wt.: 1137 (tetracosanoyl) CAS#: 71965-57-6</p> <p>Source: natural, porcine RBC Purity: 98+% by TLC Identity: confirmed by MS</p> <p>Appearance: solid Solubility: DMSO, hot methanol, chloroform/methanol, 2:1</p> <p>Storage: $-20^{\circ}C$</p> <p>See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.</p>		
1513	Ceramide trihexosides (top spot) CTH with non-hydroxy fatty acid side chain	500 μg	320.00
	<p>$C_{54}H_{101}NO_{18}$ Mol. Wt.: 1052 (stearoyl) Identity: confirmed by MS</p> <p>Source: natural, porcine RBC Purity: 98+% by TLC</p> <p>Appearance: solid Solubility: chloroform/methanol, 2:1</p> <p>Storage: $-20^{\circ}C$</p> <p>See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.</p>		
1514	Ceramide trihexosides (bottom spot) CTH with hydroxy fatty acid side chain	500 μg	350.00
	<p>$C_{60}H_{113}NO_{19}$ Mol. Wt.: 1153 (2-hydroxytetracosanoyl) Identity: confirmed by MS</p> <p>Source: natural, porcine RBC Purity: 98+% by TLC</p> <p>Appearance: solid Solubility: DMSO, hot methanol, chloroform/methanol, 1:1</p> <p>Storage: $-20^{\circ}C$</p> <p>See Table III (pg. 101) for typical fatty acid content of products prepared from natural sources.</p>		
1520	lyso-Ceramide trihexoside <i>lyso</i> -CTH; <i>lyso</i> -Globotriaosylsphingosine	1 mg	550.00
	<p>$C_{36}H_{67}NO_{17}$ Mol. Wt.: 786 CAS#: 126550-86-5</p> <p>Source: semisynthetic, porcine RBC Purity: 98+% by TLC Identity: confirmed by MS</p> <p>Appearance: solid Solubility: chloroform/methanol/DI water, 2:1:0.1</p> <p>Storage: $-20^{\circ}C$</p>		
1530	N-Glycinated lyso-ceramide trihexoside N-Glycinated globotriaosylsphingosine	1 mg	525.00
	<p>$C_{38}H_{70}N_2O_{18}$ Mol. Wt.: 843 Identity: confirmed by MS</p> <p>Source: semisynthetic, porcine RBC Purity: 98+% by TLC</p> <p>Appearance: solid Solubility: chloroform/methanol/DI water, 2:1:0.1</p> <p>Storage: $-20^{\circ}C$</p>		
1528	N-Hexadecanoyl-ceramide trihexoside N-C16:0-Ceramide trihexoside; N-Hexadecanoyl globotriaosylceramide	500 μg	425.00
	<p>$C_{52}H_{97}NO_{18}$ Mol. Wt.: 1024 Identity: confirmed by MS</p> <p>Source: semisynthetic, porcine RBC Purity: 98+% by TLC</p> <p>Appearance: solid Solubility: DMSO, hot methanol, chloroform/methanol, 2:1</p> <p>Storage: $-20^{\circ}C$</p>		

1523	N-Heptadecanoyl-ceramide trihexoside N-C17:0-Ceramide trihexoside; N-Heptadecanoyl globotriaosylceramide	500 µg	425.00
	$C_{53}H_{99}NO_{18}$ Source: semisynthetic, porcine RBC Appearance: solid Storage: -20°C	Mol. Wt.: 1038 Purity: 98+% by TLC Solubility: DMSO, hot methanol, chloroform/methanol, 2:1	Identity: confirmed by MS
1529	N-Octadecanoyl-ceramide trihexoside N-C18:0-Ceramide trihexoside; N-Octadecanoyl globotriaosylceramide	500 µg	425.00
	$C_{54}H_{101}NO_{18}$ Source: semisynthetic, porcine RBC Appearance: solid Storage: -20°C	Mol. Wt.: 1052 Purity: 98+% by TLC Solubility: DMSO, hot methanol, chloroform/methanol, 2:1	Identity: confirmed by MS
1524	N-Tricosanoyl-ceramide trihexoside N-C23:0-Ceramide trihexoside; N-Tricosanoyl globotriaosylceramide	500 µg	425.00
	$C_{59}H_{111}NO_{18}$ Source: semisynthetic, porcine RBC Appearance: solid Storage: -20°C	Mol. Wt.: 1122 Purity: 98+% by TLC Solubility: DMSO, hot methanol, chloroform/methanol, 2:1	Identity: confirmed by MS
1631 1631-001	N-Dodecanoyl-NBD-ceramide trihexoside N-C12:0-NBD-CTH; N-C12:0-NBD-Globotriaosylceramide	100 µg 1 mg	265.00 650.00
	$C_{54}H_{91}N_5O_{21}$ Source: semisynthetic, porcine RBC Appearance: solid Storage: -20°C Absorption: 460 nm	Mol. Wt.: 1145 Purity: 98+% by TLC Solubility: DMSO, hot methanol, chloroform/methanol, 2:1 Emission: 535 nm	Identity: confirmed by MS
1537	N-<i>omega</i>-CD₃-Octadecanoyl-ceramide trihexoside N-C18:0-CD ₃ -CTH; N-C18:0-CD ₃ -Gb ₃ ; N-Octadecanoyl-CD ₃ -globotriaosylceramide; N-Stearoyl-CD ₃ -ceramide trihexoside	500 µg	495.00
	$C_{54}H_{98}D_3NO_{18}$ Source: semisynthetic, porcine RBC Appearance: solid Storage: -20°C	Mol. Wt.: 1055 Purity: 98+% by TLC, HPLC Solubility: DMSO, chloroform/methanol, 2:1	Identity: confirmed by MS
1947	N-(1-Adamantaneacetyl)-ceramide trihexoside N-Adamantyl-globotriaosylceramide; AdaGb ₃	1 mg	420.00
	$C_{48}H_{83}N_4O_{18}$ Source: semisynthetic, porcine Appearance: solid Storage: -20°C	Mol. Wt.: 962 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1, methanol, DMF	Identity: confirmed by MS

Globosides



1068	Globosides Gb ₄ ; Globotetrahexosylceramide	5 mg	475.00
	<p>C₆₈H₁₂₆N₂O₂₃ Mol. Wt.: 1340 (tetracosanoyl) CAS#: 11034-93-8</p> <p>Source: natural, porcine RBC Purity: 98+% by TLC Identity: confirmed by MS</p> <p>Appearance: solid Solubility: DMSO, hot methanol, chloroform/methanol, 2:1</p> <p>Storage: -20°C</p> <p>See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.</p>		

Stable Isotope Labeled Glycolipids

1914	N-Octadecanoyl-D₃₅-psychosine, (perdeuterated, C18:0 fatty acid) N-C18:0-D ₃₅ -Cerebroside, perdeuterated; N-Stearoyl-D ₃₅ -psychosine, perdeuterated	5 mg	420.00
	<p>C₄₂H₄₆D₃₅NO₈ Mol. Wt.: 763</p> <p>Source: semisynthetic, bovine Purity: 98+% by TLC Identity: confirmed by MS</p> <p>Appearance: solid Solubility: chloroform, hot ethanol, chloroform/methanol, 2:1</p> <p>Storage: -20°C</p> <p>Activity: Deuterium labeled stearoyl side chain</p>		
2209 *NEW*	¹³C₆-Glucosylsphingosine 1-(beta-D-Glucosyl-1,2,3,4,5,6- ¹³ C ₆)-sphingosine; ¹³ C ₆ -lyso-glucocerebroside	1 mg	550.00
	<p>C₁₈¹³C₆H₄₇NO₇ Mol. Wt.: 468</p> <p>Source: synthetic Purity: 98+% by TLC, HPLC Identity: confirmed by MS</p> <p>Appearance: solid Solubility: ethanol, methanol, chloroform/methanol, 2:1</p> <p>Storage: -20°C</p>		
1533	N-omega-CD₃-Hexadecanoyl-glucopsychosine N-C16:0-CD ₃ -Glucopsychosine; N-C16:0-CD ₃ -Glucocerebroside; N-Palmitoyl-CD ₃ -glucopsychosine	1 mg	400.00
	<p>C₄₀H₇₄D₃NO₈ Mol. Wt.: 703</p> <p>Source: semisynthetic, bovine buttermilk Purity: 98+% by TLC Identity: confirmed by MS</p> <p>Appearance: solid Solubility: chloroform/methanol, 2:1</p> <p>Storage: -20°C</p>		
1536	N-omega-CD₃-Octadecanoyl-sulfatide N-C18:0-CD ₃ -Sulfatide; N-Stearoyl-CD ₃ -sulfatide	1 mg	550.00
	<p>C₄₂H₇₈D₃NO₁₁S Mol. Wt.: 811</p> <p>Source: semisynthetic, bovine Purity: 98+% by TLC Identity: confirmed by MS</p> <p>Appearance: solid Solubility: chloroform/methanol/DI water, 2:1:0.1</p> <p>Storage: -20°C</p>		

1534	N-<i>omega</i>-CD₃-Hexadecanoyl-lactosylceramide N-C16:0-CD ₃ -Lactosylceramide; N-Palmitoyl-CD ₃ -lactosylceramide	1 mg	460.00
	C₄₆H₈₄D₃NO₁₃ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 865 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 5:1:0.1	Identity: confirmed by MS

1537	N-<i>omega</i>-CD₃-Octadecanoyl-ceramide trihexoside N-C18:0-CD ₃ -CTH; N-C18:0-CD ₃ -Gb ₃ ; N-Octadecanoyl-CD ₃ -globotriaosylceramide; N-Stearoyl-CD ₃ -ceramide trihexoside	500 µg	495.00
	C₅₄H₉₈D₃NO₁₈ Source: semisynthetic, porcine RBC Appearance: solid Storage: -20°C	Mol. Wt.: 1055 Purity: 98+% by TLC , HPLC Solubility: DMSO, chloroform/methanol, 2:1	Identity: confirmed by MS

Fluorescent Glycolipids

Absorption: 460 nm Emission: 535 nm

1621	N-Hexanoyl-NBD-galactosylceramide N-C6:0-NBD- <i>beta</i> -D-Galactosylsphingosine; N-C6:0-NBD-Cerebroside	100 µg	250.00
	C₃₆H₅₉N₅O₁₁ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 738 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 5:1	

1633 1633-001	N-Dodecanoyl-NBD-galactosylceramide N-C12:0-NBD- <i>beta</i> -D-Galactosylsphingosine; N-C12:0-NBD-Cerebroside	100 µg 1 mg	210.00 450.00
	C₄₂H₇₁N₅O₁₁ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 822 Purity: 98+% by TLC Solubility: chloroform, DMSO, chloroform/methanol, 2:1	CAS#: 474942-98-8 Identity: confirmed by MS

1622 1622-001	N-Hexanoyl-NBD-glucosylceramide N-C6:0-NBD- <i>beta</i> -D-Glucosylsphingosine; N-C6:0-NBD-Glucosylceramide	100 µg 1 mg	250.00 400.00
	C₃₆H₅₉N₅O₁₁ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 738 Purity: 98+% by TLC Solubility: methanol, chloroform/methanol, 5:1	Identity: confirmed by MS

1632 1632-001	N-Dodecanoyl-NBD-sulfatide N-C12:0-NBD-Sulfatide; N-Dodecanoyl-NBD- <i>lyso</i> -sulfatide; N-Dodecanoyl-NBD-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	100 µg 1 mg	170.00 450.00
	C₄₂H₇₁N₅O₁₄S Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 901 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	


1629	N-Hexanoyl-NBD-lactosylceramide	50 ug	265.00
1629-001	N-Hexanoyl-NBD- <i>beta</i> -D-lactosylsphingosine; N-C6:0-NBD- <i>beta</i> -D-Lactosylsphingosine; N-C6:0-NBD-Lactosylceramide	1 mg	500.00
	$C_{42}H_{69}N_5O_{16}$	Mol. Wt.: 900	
	Source: semisynthetic, bovine buttermilk	Purity: 98+% by TLC	
	Appearance: solid	Solubility: chloroform/methanol, 2:1	
	Storage: -20°C		
1630	N-Dodecanoyl-NBD-lactosylceramide	50 ug	295.00
1630-001	N-Dodecanoyl-NBD- <i>beta</i> -D-lactosylsphingosine; N-C12:0-NBD- <i>beta</i> -D-Lactosylsphingosine; N-C12:0-NBD-Lactosylceramide	1 mg	500.00
	$C_{48}H_{81}N_5O_{16}$	Mol. Wt.: 984	
	Source: semisynthetic, bovine buttermilk	Purity: 98+% by TLC	
	Appearance: solid	Solubility: chloroform/methanol, 2:1	
	Storage: -20°C		
1631	N-Dodecanoyl-NBD-ceramide trihexoside	100 µg	265.00
1631-001	N-C12:0-NBD-CTH; N-C12:0-NBD-Globotriaosylceramide	1 mg	650.00
	$C_{54}H_{91}N_5O_{21}$	Mol. Wt.: 1145	
	Source: semisynthetic, porcine RBC	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: DMSO, hot methanol, chloroform/methanol, 2:1	
	Storage: -20°C		
2204	Lissamine-rhodamine B-dodecanoyl-galactosylceramide	500 µg	390.00
	Sulforhodamine B-C12:0 cerebroside		
	$C_{63}H_{99}N_4O_{14}S_2$	Mol. Wt.: 1201	
	Source: semisynthetic, bovine	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform/methanol 8:2, DMSO, DMF	
	Storage: -20°C		
	Absorption: 540 nm	Emission: 565 nm	

See Labeled Standards section (page 96) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical service for more information. 814-355-1030

Gangliosides

The diagram below can be used with the general formulas given in the ganglioside descriptions to construct the individual structures.



1064	Gangliotetraosylceramide Asialo GM ₁ ; Gg4	1 mg	390.00
	$C_{62}H_{114}N_2O_{23}$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C Note: General formula: 1,2,3,4 See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1256 (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 71012-19-6 Identity: confirmed by MS
1512	Gangliotriaosylceramide Asialo GM ₂ ; Gg3	100 µg	630.00
	$C_{56}H_{104}N_2O_{18}$ Source: semisynthetic, human Appearance: solid Storage: -20°C Note: General formula: 1,2,3	Mol. Wt.: 1093 (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 35960-33-9 Identity: confirmed by MS
1061 1061-50	Monosialoganglioside GM₁ (NH₄⁺ salt) GM ₁	5 mg 50 mg	315.00 2,450.00
	$C_{73}H_{131}N_3O_{31} \cdot NH_3$ Source: natural, bovine Appearance: solid Storage: -20°C Note: General formula: 1,2,3,4,5 See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1547 + NH ₃ (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 37758-47-7 Identity: confirmed by MS
1518	lyso-Monosialoganglioside GM₁ (NH₄⁺ salt) <i>lyso</i> -GM ₁	500 µg	390.00
	$C_{55}H_{97}N_3O_{30} \cdot NH_3$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 1280 + NH ₃ Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.2	CAS#: 171483-40-2 Identity: confirmed by MS

1526	Fucosylated monosialoganglioside GM₁ (NH₄⁺ salt) Fucosyl-GM ₁	500 µg	595.00
	C₇₉H₁₄₁N₃O₃₅•NH₃ Source: natural, porcine Appearance: solid Storage: -20°C See Table III (pg. 101) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1693 + NH ₃ (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 71812-11-8 Identity: confirmed by MS
2050	N-<i>omega</i>-CD₃-Octadecanoyl monosialoganglioside GM₁ (NH₄⁺ salt) N-CD ₃ -Stearoyl GM ₁	500 µg	550.00
	C₇₃H₁₂₈N₃O₃₁D₃•NH₃ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 1550 + NH ₃ Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	Identity: confirmed by MS
2053	N-Hexanoyl-biotin-monosialoganglioside GM₁ Biotin-C6:0-GM ₁	500 µg	575.00
	C₇₁H₁₂₂N₆O₃₃S Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 1620 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS
1502	Monosialoganglioside GM₂ (NH₄⁺ salt) GM ₂	500 µg	335.00
	C₆₇H₁₂₁N₃O₂₆•NH₃ Source: natural, human Tay-Sachs Appearance: solid Storage: -20°C Note: General formula: 1,2,3,5 See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1385+ NH ₃ (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 19600-01-2
2051	N-<i>omega</i>-CD₃-Octadecanoyl monosialoganglioside GM₂ (NH₄⁺ salt) N-CD ₃ -Stearoyl GM ₂	250 µg	490.00
	C₆₇H₁₁₈D₃N₃O₂₆•NH₃ Source: semisynthetic, human Tay-Sachs Appearance: solid Storage: -20°C	Mol. Wt.: 1388 + NH ₃ Purity: 98+% by TLC, MS Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	Identity: confirmed by MS
1503	Monosialoganglioside GM₃ (NH₄⁺ salt) GM ₃	1 mg	175.00
	C₆₄H₁₁₈N₂O₂₁•NH₃ Source: natural, bovine buttermilk Appearance: solid Storage: -20°C Note: General formula: 1,2,5 See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1252+ NH ₃ (tricosanoyl) Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1; forms micellar solution in water	CAS#: 54827-14-4 Identity: confirmed by MS
2052	N-<i>omega</i>-CD₃-Octadecanoyl monosialoganglioside GM₃ (NH₄⁺ salt) N-CD ₃ -Stearoyl GM ₃	250 µg	410.00
	C₅₉H₁₀₅D₃N₂O₂₁•NH₃ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 1185 + NH ₃ Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	Identity: confirmed by MS

1535	Monosialoganglioside GM₄ (NH₄⁺ salt) GM ₄	500 µg	450.00
	C₅₇H₁₀₆N₂O₁₇•NH₃ Source: natural, egg, chicken Appearance: solid Storage: -20°C Note: General formula: 2,5 See Table III (pg. 101) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1091+NH ₃ (2-hydroxydocosanoyl) Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1; forms micellar solution in water	CAS#: 66456-69-7 Identity: confirmed by MS
1062	Disialoganglioside GD_{1a} (NH₄⁺ salt) GD _{1a}	5 mg	295.00
	C₈₄H₁₄₈N₄O₃₉•2NH₃ Source: natural, bovine Appearance: solid Storage: -20°C Note: General formula: 1,2,3,4,5,7 See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1838 + 2NH ₃ (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 12707-58-3 Identity: confirmed by MS
1501	Disialoganglioside GD_{1b} (NH₄⁺ salt) GD _{1b}	1 mg	285.00
	C₈₄H₁₄₈N₄O₃₉•2NH₃ Source: natural, bovine Appearance: solid Storage: -20°C Note: General formula: 1,2,3,4,5,6 See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1838 + 2NH ₃ (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 19553-76-5 Identity: confirmed by MS
1527	Disialoganglioside GD₂ (NH₄⁺ salt) GD ₂	500 µg	665.00
	C₇₈H₁₃₈N₄O₃₄•2NH₃ Source: semisynthetic, rabbit Appearance: solid Storage: -20°C Note: General formula: 1,2,3,5,6 See Table III (pg. 101) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1676 + 2NH ₃ (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 65988-71-8 Identity: confirmed by MS
1504 1504-25	Disialoganglioside GD₃ (NH₄⁺ salt) GD ₃	5 mg 25 mg	315.00 925.00
	C₇₅H₁₃₅N₃O₂₉•2NH₃ Source: natural, bovine buttermilk Appearance: solid Storage: -20°C Note: General formula: 1,2,5,6 See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 1543+2NH ₃ (tricosanoyl) Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1; forms micellar solution in water	CAS#: 62010-37-1 Identity: confirmed by MS
2054 *NEW*	N-omega-CD₃-Octadecanoyl disialoganglioside GD₃ N-CD ₃ -Stearoyl GD ₃	500 µg	550.00
	C₇₀H₁₂₂D₃N₃O₂₉ Source: semisynthetic bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 1476 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1, water	Identity: confirmed by MS

2055 *NEW*	N-Hexanoyl-biotin-disialoganglioside GD₃ Biotin-C6:0-GD ₃	500 µg	575.00
	C₆₈H₁₁₆N₆O₃₁S Source: semisynthetic bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 1546 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS
1063	Trisialoganglioside GT_{1b} (NH₄⁺ salt) GT _{1b}	5 mg	370.00
	C₉₅H₁₆₅N₅O₄₇•3NH₃ Source: natural, bovine Appearance: solid Storage: -20°C Note: General Formula: 1,2,3,4,5,6,7 See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 2129 + 3NH ₃ (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 59247-13-1 Identity: confirmed by MS
1516	Tetrasialoganglioside GQ_{1b} (NH₄⁺ salt) GQ _{1b}	100 µg	245.00
	C₁₀₆H₁₈₂N₆O₅₅•4NH₃ Source: natural, bovine Appearance: solid Storage: -20°C Note: General formula: 1,2,3,4,5,6,7,8 See Table III (pg. 101) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 2421+4NH ₃ (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	CAS#: 68652-37-9 Identity: confirmed by MS
1065	Mixed Gangliosides, purified (NH₄⁺ salt), bovine Mixed Gangliosides	25 mg	270.00
	Source: natural, bovine Appearance: solid Storage: -20°C See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.	Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water Approximately 98% GM ₁ , GD _{1a} , GD _{1b} and GT _{1b} , remaining 2% other gangliosides	
1525	Mixed Gangliosides, purified (NH₄⁺ salt), porcine	25 mg	280.00
	Source: natural, porcine Appearance: solid Storage: -20°C See Table III (pg. 101) for typical fatty acid content of products prepared from natural sources.	Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water Approximately 98% GM ₁ , GD _{1a} , GD _{1b} and GT _{1b} , remaining 2% other gangliosides	

Glycosphingolipid Reference Mixtures for TLC

These mixtures are qualitative standards prepared from our purified glycosphingolipids.

1505	Neutral Glycosphingolipid Mixture	1 mg/ml, 1 ml	210.00
	Source: natural, bovine and porcine Appearance: liquid Storage: -20°C Contains: cerebrosides, lactosylceramides, ceramide trihexosides, globosides.	Solubility: chloroform/methanol, 2:1 Solvent: chloroform/methanol, 2:1	
1508	Monosialoganglioside Mixture	0.5 mg/ml, 1 ml	210.00
	Source: natural, bovine, human Appearance: liquid Storage: -20°C Contains: GM ₃ , GM ₂ , GM ₁	Solubility: chloroform/methanol/DI water, 2:1:0.1 Solvent: chloroform/methanol/DI water, 2:1:0.1	

1509	Disialoganglioside Mixture	0.5 mg/ml, 1 ml	210.00
	Source: natural, bovine Appearance: liquid Storage: -20°C Contains: GD ₃ , GD _{1a} , GD _{1b}	Solubility: chloroform/methanol/DI water, 2:1:0.1 Solvent: chloroform/methanol/DI water, 2:1:0.1	
1510	Lactosylceramide and Sialosyl Derivatives Mixture	0.5 mg/ml, 1 ml	225.00
	Source: natural, bovine buttermilk Appearance: liquid Storage: -20°C Contains: LC, GM ₃ , GD ₃	Solubility: chloroform/methanol/DI water, 2:1:0.1 Solvent: chloroform/methanol/DI water, 2:1:0.1	
1511	Gangliotetraosylceramide and Sialosyl Derivatives Mixture	0.5 mg/ml, 1 ml	210.00
	Source: natural, bovine Appearance: liquid Storage: -20°C Contains: asialo GM ₁ , GM ₁ , GD _{1a} , GD _{1b} , GT _{1b}	Solubility: chloroform/methanol/DI water, 2:1:0.1 Solvent: chloroform/methanol/DI water, 2:1:0.1	

Antibodies Directed Against Glycolipids

These monoclonal and polyclonal antibodies are directed against the carbohydrate chains of Matreya's glycolipids. The same carbohydrate moieties are found on many glycoproteins. The antibodies are for use in ELISA or TLC immunoblotting applications (9). All antibodies are quality tested by actual performance in ELISA and TLC immunoblotting. The antibodies contain no preservatives and are shipped on dry ice. See Literature References on page 106.

1977	Anti-ganglioside GD₃	50 µl	375.00
	Monoclonal antibody to GD ₃ , isotype IgG/IgM Source: natural, mouse hybridoma R-24 cell line Appearance: liquid Storage: -20°C Suitable for TLC immunoblotting, ELISA	Solubility: DI water Dry Ice Charge Applies	
1950	Anti-ganglioside asialo GM₁	100 µl	470.00
	Polyclonal antibody to asialo GM ₁ , isotype IgG/IgM Source: natural, rabbit Appearance: liquid Storage: -20°C Suitable for ELISA, TLC-immunoblotting. Slight cross reaction to GM ₁	Solubility: DI water Dry Ice Charge Applies	
1951	Anti-ganglioside asialo GM₂	50 µl	375.00
	Polyclonal antibody to asialo GM ₂ , isotype IgG/IgM Source: natural, rabbit Appearance: liquid Storage: -20°C Suitable for ELISA, TLC-immunoblotting	Solubility: DI water Dry Ice Charge Applies	
1954	Anti-ganglioside GM₁	100 µl	375.00
	Polyclonal antibody to GM ₁ , isotype IgG/IgM Source: natural, rabbit Appearance: liquid Storage: -20°C Suitable for ELISA, TLC-immunoblotting. Slight cross reaction to asialo-GM ₁	Solubility: DI water Dry Ice Charge Applies	

1964	Anti-ganglioside GD_{1b} Polyclonal antibody to GD _{1b} , isotype IgG/IgM	50 µl	390.00
	Source: natural, rabbit Appearance: liquid Storage: -20°C Suitable for ELISA, TLC-immunoblotting	Solubility: DI water	Dry Ice Charge Applies
1960	Anti-globoside GL-4 Polyclonal antibody to GL-4, isotype IgG/IgM	50 µl	375.00
	Source: natural, rabbit Appearance: liquid Storage: -20°C Suitable for ELISA, TLC-immunoblotting	Solubility: DI water	Dry Ice Charge Applies

Enzyme Inhibitors


Ceramide: UDPglucose Transferase. PDMP (D,L-*threo*-1-phenyl-2-decanoylamino-3-morpholino-1-propanol-HCl) closely resembles the natural sphingolipid substrate of brain glucosyl transferase and is a very potent and competitive inhibitor of the enzyme (26). It has been shown to block outgrowth of neurites in cultured retina and to block glucolipid synthesis in cultured 3T3 cells (27). N.S. Radin and co-workers have shown (28) that PPMP has activity equivalent to that of PDMP when cell homogenates and brain and liver microsomes are used, but it is about 20 times more potent when used with intact cells. In another paper (29), Radin's group has shown that PDMP has substantial activity against Ehrlich ascites tumors in mice. Recent publications from the laboratory of Myles Cabot (30, 31) show that PPMP can reverse multi-drug resistance in cancer cells by causing a build-up of ceramide and preventing the synthesis of glycosylated ceramides. **See Literature References on page 106.**

Matreya also offers the resolved D- and L-*threo*-isomers of PDMP and PPMP.

Protein Kinase C Inhibitor. Sphingosine is a potent and reversible inhibitor of protein kinase C (32); it also has been shown at low concentrations to stimulate DNA synthesis and act synergistically with known growth factors (33). Note that Safingol (our L-*threo*-dihydrosphingosine) has also been shown to partially reverse multi-drug resistance in cancer cells (31) *via* inhibition of protein kinase C.


Dihydroceramide desaturase Inhibitor. Cyclopropenylceramide is the first known inhibitor of this enzyme and may allow significant studies on the role of ceramide in apoptosis. Matreya is the only source for this inhibitor. (34)

Ceramidase Inhibitors. N-Oleoylethanolamine has been shown to be an efficacious inhibitor of the ceramidase found in human kidney and cerebellum (35). It is specifically an inhibitor of acid ceramidase (36) with an IC₅₀ of ca. 500 µM. N-Hexadecanoylethanolamine can be used as an inactive control. D-MAPP is a potent (IC₅₀ approximately 5 µM) inhibitor of alkaline ceramidase. Its enantiomer L-MAPP is inactive as an inhibitor and acts as a substrate for this enzyme (36, 37). **See Literature References on page 106.**




1751	N-Oleoylethanolamine NOE	100 mg	230.00
	C₂₀H₃₉NO₂ Source: synthetic Appearance: solid Storage: -20°C Activity: acid ceramidase inhibitor	Mol. Wt.: 326 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, ethyl ether, DMSO Melting Point (°C): 63-66	CAS#: 111-58-0 Identity: confirmed by MS

1807	L-threo-Dihydrosphingosine (Safingol)	5 mg	240.00
1807-025	L-threo-Sphinganine, C18 chain	25 mg	725.00
	C ₁₈ H ₃₉ NO ₂	Mol. Wt.: 301	CAS#: 15639-50-6
	Source: synthetic	Purity: 98+% by TLC, GC	
	Appearance: solid	Solubility: chloroform, ethanol, methanol	
	Storage: -20°C	Melting Point (°C): 103-114	
	Activity: Protein Kinase C (PKC) and Sphingosine Kinase inhibitor		



1859	D-MAPP	100 mg	210.00
	D-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol		
	C ₂₃ H ₃₉ NO ₂	Mol. Wt.: 361	CAS#: 143492-39-1
	Source: synthetic	Purity: 98+% by TLC	
	Appearance: solid	Solubility: ethanol	
	Storage: -20°C		
	Activity: alkaline ceramidase inhibitor		

1860	L-MAPP	100 mg	210.00
	L-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol		
	C ₂₃ H ₃₉ NO ₂	Mol. Wt.: 361	CAS#: 143492-38-0
	Source: synthetic	Purity: 98+% by TLC	
	Appearance: solid	Solubility: ethanol	
	Storage: -20°C		
	Activity: inactive as alkaline ceramidase inhibitor		



1889	Conduritol B Epoxide	25 mg	270.00
	D,L-1,2-Anhydro-myo-inositol		
	C ₆ H ₁₀ O ₅	Mol. Wt.: 162	CAS#: 6090-95-5
	Source: synthetic	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: DI water, DMSO, methanol (slightly)	
	Storage: -20°C	Melting Point (°C): 164-166	
	Activity: Inhibits α- and β-glucosidase activity; specific inhibitor of glucocerebrosidase in cultured cells.		


PDMP's can be solubilized in 5% tween-80 in saline. S. Chatterjee et al. PLoS One. 2013; 8(5): e63726

1719 **D,L-threo-PDMP** **100 mg** **250.00**
D,L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl

$C_{23}H_{38}N_2O_3 \cdot HCl$ **Mol. Wt.:** 427 **CAS#:** 80938-69-8
Source: synthetic **Purity:** 98+% by TLC **Identity:** confirmed by MS
Appearance: solid **Solubility:** ethanol, methanol, chloroform, DMSO
Storage: -20°C
Activity: glucosylceramide synthase inhibitor

1720 **D,L-threo-PPMP** **100 mg** **250.00**
D,L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl

$C_{29}H_{50}N_2O_3 \cdot HCl$ **Mol. Wt.:** 511 **CAS#:** 149022-18-4
Source: synthetic **Purity:** 98+% by TLC **Identity:** confirmed by MS
Appearance: solid **Solubility:** alcohols, chloroform
Storage: -20°C
Activity: glucosylceramide synthase inhibitor



1749 **L-threo-PDMP** **10 mg** **390.00**
L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl

$C_{23}H_{38}N_2O_3 \cdot HCl$ **Mol. Wt.:** 427 **CAS#:** 109836-81-9
Source: synthetic **Purity:** 98+% by TLC
Appearance: solid **Solubility:** ethanol, methanol
Storage: -20°C

1753 **D,L-erythro-PPMP** **100 mg** **235.00**
D,L-erythro-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl


$C_{29}H_{50}N_2O_3 \cdot HCl$ **Mol. Wt.:** 511 **Identity:** confirmed by MS
Source: synthetic **Purity:** 98+% by TLC **Solubility:** chloroform, ethanol, methanol, DMSO
Appearance: solid
Storage: -20°C

1755 **D,L-erythro-PDMP** **100 mg** **235.00**
D,L-erythro-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl

$C_{23}H_{38}N_2O_3 \cdot HCl$ **Mol. Wt.:** 427 **CAS#:** 109760-77-2
Source: synthetic **Purity:** 98+% by TLC **Identity:** confirmed by MS
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: -20°C

1756 **D-threo-PDMP** **10 mg** **345.00**
D-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl

$C_{23}H_{38}N_2O_3 \cdot HCl$ **Mol. Wt.:** 427 **CAS#:** 109836-82-0
Source: synthetic **Purity:** 98+% by TLC **Identity:** confirmed by MS
Appearance: solid **Solubility:** ethanol, methanol
Storage: -20°C
Activity: glucosylceramide synthase inhibitor




1865 **D-threo-PPMP** **10 mg** **325.00**
D-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl

$C_{29}H_{50}N_2O_3 \cdot HCl$ **Mol. Wt.:** 511
Source: synthetic **Purity:** 98+% by TLC
Appearance: solid **Solubility:** ethanol, methanol
Storage: $-20^{\circ}C$
Activity: glucosylceramide synthase inhibitor

1868 **L-threo-PPMP** **10 mg** **325.00**
L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl

$C_{29}H_{50}N_2O_3 \cdot HCl$ **Mol. Wt.:** 511 **CAS#:** 207278-87-3
Source: synthetic **Purity:** 98+% by TLC
Appearance: solid **Solubility:** ethanol, methanol
Storage: $-20^{\circ}C$



1717 **2-Fluoropalmitic acid** **25 mg** **220.00**

$C_{16}H_{31}FO_2$ **Mol. Wt.:** 274 **CAS#:** 89270-22-4
Source: synthetic **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol, methanol
Storage: $-20^{\circ}C$ **Melting Point ($^{\circ}C$):** 83-85
Activity: Acyl-CoA synthase inhibitor

1718 **Methyl 2-fluoropalmitate** **10 mg** **220.00**

$C_{17}H_{33}FO_2$ **Mol. Wt.:** 288
Source: synthetic **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol, methanol
Storage: $-20^{\circ}C$ **Melting Point ($^{\circ}C$):** 36-38
Activity: inactive ester of 2-fluoropalmitic acid


1858 **2-Acetyl-4-(1R, 2S, 3R, 4-tetrahydroxybutyl)-imidazole** **1 mg** **150.00**
 THI

$C_9H_{14}N_2O_5$ **Mol. Wt.:** 230 **CAS#:** 94944-70-4
Source: synthetic **Purity:** 99% by HPLC, MS, NMR
Appearance: solid **Solubility:** DI water
Storage: $-20^{\circ}C$
Activity: Sphingosine-1-phosphate lyase inhibitor

1945 **N-(1-Adamantaneacetyl)-glucosylceramide** **5 mg** **420.00**
 N-(1-Adamantaneacetyl)-glucocerebroside

$C_{36}H_{61}NO_8$ **Mol. Wt.:** 636
Source: semisynthetic, plant **Purity:** 98+% by TLC **Identity:** confirmed by MS
Appearance: solid **Solubility:** chloroform, methanol, chloroform/methanol, 9:1
Storage: $-20^{\circ}C$
Activity: inhibitor of glucocerebrosidase and lactosylceramide synthase

1946	N-(1-Adamantaneacetyl)-galactosylceramide N-(1-Adamantaneacetyl)-galactocerebroside	5 mg	420.00
	$C_{36}H_{63}NO_8$ Source: semisynthetic, bovine Appearance: solid Storage: $-20^{\circ}C$ Activity: inhibitor of glucosylceramide, sulfatide, and globotriaosylceramide (Gb ₃) synthase	Mol. Wt.: 638 Purity: 98+% by TLC Solubility: chloroform, methanol, chloroform/methanol, 9:1	Identity: confirmed by MS
1320	N,N-Dimethyl-D-erythro-sphingosine	5 mg/ml, 1 ml	135.00
	$C_{20}H_{41}NO_2$ Source: synthetic Appearance: liquid Storage: $-20^{\circ}C$ Activity: Inhibitor of phosphokinase C	Mol. Wt.: 328 Purity: 98+% by TLC Solubility: chloroform, ethanol, isopropanol, methanol Solvent: isopropanol	CAS#: 119567-63-4 Identity: confirmed by MS




1896	N,N-Dihexyl-D-erythro-sphingosine Sphingosine with tertiary amine group	5 mg/ml, 1 ml	225.00
	$C_{30}H_{61}NO_2$ Source: synthetic Appearance: liquid Storage: $-20^{\circ}C$	Mol. Wt.: 468 Purity: 95% by TLC Solubility: chloroform, ethanol, methanol Solvent: ethanol	


Glycerolipids

Glycerophospholipids

Natural Phospholipids



1044	Lecithin, egg Phosphatidylcholine; PC	50 mg/ml, 1 ml	75.00
	$C_{44}H_{84}NO_8P$ Source: natural, chicken, egg Appearance: liquid Storage: $-20^{\circ}C$ See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 787 (oleoyl) Purity: 98+% by TLC Solubility: chloroform, ethyl ether, ethanol Solvent: chloroform	CAS#: 8002-43-5


1070	Lecithin, bovine Phosphatidylcholine; PC	50 mg/ml, 1 ml	285.00
	$C_{44}H_{84}NO_8P$ Source: natural, bovine Appearance: liquid Storage: $-20^{\circ}C$ See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 787 (oleoyl) Purity: 98+% by TLC Solubility: chloroform, ethyl ether Solvent: chloroform	CAS#: 8002-43-5
			
1046	lyso-Lecithin, egg <i>lyso</i> -Phosphatidylcholine	50 mg	120.00
	$C_{24}H_{50}NO_7P$ Source: semisynthetic, chicken, egg Appearance: solid Storage: $-20^{\circ}C$ See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 496 (palmitoyl) Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	CAS#: 9008-30-4
1047	Phosphatidylserine, bovine PS	50 mg/ml, 1 ml	275.00
	$C_{42}H_{78}NO_{10}P$ Source: natural, bovine Appearance: liquid Storage: $-20^{\circ}C$ See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 788 (oleoyl) Purity: 98+% by TLC Solubility: chloroform, toluene Solvent: chloroform	CAS#: 51446-62-9 Identity: confirmed by MS
1048	Phosphatidylinositol (Na⁺ salt), plant, wheat germ PI	10 mg/ml, 1 ml	180.00
	$C_{45}H_{78}O_{13}P \cdot Na$ Source: natural, plant Appearance: liquid Storage: $-20^{\circ}C$ See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 858 + Na (linoleoyl) Purity: 98+% by TLC Solubility: chloroform, ethyl ether Solvent: chloroform	CAS#: 383907-36-6 Identity: confirmed by MS
1336	Phosphatidylinositol (Na⁺ salt), plant, soy	50 mg/ml, 1ml	455.00
	$C_{46}H_{80}O_{13}P \cdot Na$ Source: natural, plant, soy Appearance: liquid Storage: $-20^{\circ}C$ See Table III (pg. 100) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 858 + Na (linoleoyl) Purity: 98+% by TLC Solubility: chloroform, ethyl ether Solvent: chloroform	CAS#: 383907-36-6
1053	Phosphatidic acid (NH₄⁺ salt), egg PA	50 mg	155.00
	$C_{39}H_{72}O_8P \cdot NH_4^+$ Source: semisynthetic, Chicken, egg Appearance: solid Storage: $-20^{\circ}C$ See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 718 (oleoyl) Purity: 98+% by TLC Solubility: chloroform, ethyl ether	Identity: confirmed by MS

1045	Phosphatidylethanolamine, egg PE	50 mg/ml, 1 ml	250.00
	<p>$C_{41}H_{78}NO_8P$ Source: natural, chicken, egg Appearance: liquid Storage: $-20^{\circ}C$ See Table III (pg. 97) for typical fatty acid content of products prepared from natural sources.</p>	<p>Mol. Wt.: 744 (oleoyl) Purity: 98+% by TLC Solubility: chloroform Solvent: chloroform</p>	<p>CAS#: 39382-08-6 Identity: confirmed by MS</p>
1069	Phosphatidylethanolamine, bovine PE	50 mg/ml, 1 ml	300.00
	<p>$C_{41}H_{78}NO_8P$ Source: natural, bovine Appearance: liquid Storage: $-20^{\circ}C$</p>	<p>Mol. Wt.: 744 (oleoyl) Purity: 98+% by TLC Solubility: chloroform Solvent: chloroform</p>	<p>CAS#: 90989-93-8</p>
1301	Phosphatidylethanolamine, plant PE	50 mg/ml, 1 ml	270.00
	<p>$C_{41}H_{74}NO_8P$ Source: natural, plant Appearance: liquid Storage: $-20^{\circ}C$ See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.</p>	<p>Mol. Wt.: 740 (oleoyl) Purity: 98+% by TLC Solubility: chloroform Solvent: chloroform</p>	<p>CAS#: 90989-93-8</p>

Synthetic Phospholipids

These phospholipids have 98+% chemical purity except where stated and 99% fatty acid chain purity. Store at $-20^{\circ}C$.

Phosphatidic Acid Derivatives




1428	1,2-Dimyristoyl-sn-glycero-3-phosphatidic acid (Na⁺ salt) DMPA	100 mg	90.00
	<p>$C_{31}H_{60}O_8P\cdot Na$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$</p>	<p>Mol. Wt.: 615 Purity: 98+% by TLC Solubility: chloroform/methanol/acetic acid, 4:1:0.1</p>	<p>CAS#: 80724-31-8</p>

1429	1,2-Dipalmitoyl-sn-glycero-3-phosphatidic acid (Na⁺ salt) DPPA	100 mg	90.00
	C ₃₅ H ₆₈ O ₈ P•Na Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 671 Purity: 98+% by TLC Solubility: chloroform/methanol/acetic acid, 4:1:0.1	CAS#: 71065-87-7

1430	1,2-Distearoyl-sn-glycero-3-phosphatidic acid (Na⁺ salt) DSPA	100 mg	90.00
	C ₃₉ H ₇₆ O ₈ P•Na Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 727 Purity: 98+% by TLC Solubility: chloroform/methanol/acetic acid, 4:1:0.1	CAS#: 108321-18-2

Phosphatidylcholines

1442	1,2-Dilauroyl-sn-glycero-3-phosphorylcholine DLPC	100 mg	90.00
	C ₃₂ H ₆₄ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 622 Purity: 98+% by TLC Solubility: methylene chloride, methanol	CAS#: 18194-25-7




1425	1,2-Dimyristoyl-sn-glycero-3-phosphorylcholine DMPC	100 mg	90.00
	C ₃₆ H ₇₂ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 678 Purity: 98+% by TLC Solubility: methylene chloride, methanol Melting Point: 130-139°C	CAS#: 18194-24-6

1426	1,2-Dipalmitoyl-sn-glycero-3-phosphorylcholine DPPC	100 mg	90.00
	C ₄₀ H ₈₀ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 734 Purity: 98+% by TLC Solubility: methylene chloride, methanol	CAS#: 63-89-8

1400	1,2-Diheptadecanoyl-sn-glycero-3-phosphorylcholine DHDPC	50 mg	90.00
	C ₄₂ H ₈₄ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 762 Purity: 98+% by TLC Solubility: methylene chloride, methanol	CAS#: 70897-27-7 Identity: confirmed by MS

1427	1,2-Distearoyl-sn-glycero-3-phosphorylcholine DSPC	100 mg	90.00
	C ₄₄ H ₈₈ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 790 Purity: 98+% by TLC Solubility: methylene chloride, methanol	CAS#: 816-94-4
1437	1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylcholine POPC	100 mg	120.00
	C ₄₂ H ₈₂ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 760 Purity: 98+% by TLC Solubility: methylene chloride, methanol	CAS#: 26853-31-6
1445	1-Palmitoyl-sn-glycero-3-phosphorylcholine <i>lyso</i> -PPC	100 mg	120.00
	C ₂₄ H ₅₀ NO ₇ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 496 Purity: 98+% by TLC Solubility: methylene chloride, methanol	CAS#: 17364-16-8

Phosphatidylglycerols




1443	1,2-Dilauroyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt) DLPG	100 mg	90.00
	C ₃₀ H ₅₈ O ₁₀ P•Na Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 632 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1	CAS#: 73548-69-3
1431	1,2-Dimyristoyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt) DMPG	100 mg	90.00
	C ₃₄ H ₆₆ O ₁₀ P•Na Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 689 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1 Melting Point: 120-129°C	CAS#: 200880-40-6
1432	1,2-Dipalmitoyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt) DPPG	100 mg	90.00
	C ₃₈ H ₇₄ O ₁₀ P•Na Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 745 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1 Melting Point: 122-127°C	CAS#: 200880-41-7

1433	1,2-Distearoyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt) DSPG	100 mg	90.00
	C ₄₂ H ₈₂ O ₁₀ P•Na Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 801 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1	CAS#: 4537-78-4

1438	1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt) POPG	100 mg	180.00
	C ₄₀ H ₇₆ O ₁₀ P•Na Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 771 Purity: 98+% by TLC Solubility: chloroform/methanol, 5:1	CAS#: 202070-86-8

Phosphatidylethanolamines



1444	1,2-Dilauroyl-sn-glycero-3-phosphorylethanolamine DLPE	100 mg	100.00
	C ₂₉ H ₅₈ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 580 Purity: 98+% by TLC Solubility: chloroform + methanol mixture	CAS#: 42436-56-6

1434	1,2-Dimyristoyl-sn-glycero-3-phosphorylethanolamine DMPE	100 mg	100.00
	C ₃₃ H ₆₆ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 636 Purity: 98+% by TLC Solubility: chloroform/acetic acid, 95:5; chloroform/methanol/DI water/acetic acid, 100:30:10:2.5	CAS#: 998-07-2

1435	1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine DPPE	100 mg	100.00
	C ₃₇ H ₇₄ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 692 Purity: 98+% by TLC Solubility: chloroform/acetic acid, 95:5; chloroform/methanol/DI water/acetic acid, 100:30:10:2.5	CAS#: 923-61-5

1436	1,2-Distearoyl-sn-glycero-3-phosphorylethanolamine DSPE	100 mg	100.00
	C ₄₁ H ₈₂ NO ₈ P Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 748 Purity: 98+% by TLC Solubility: chloroform/acetic acid, 95:5; chloroform/methanol/DI water/acetic acid, 100:30:10:2.5	CAS#: 1069-79-0

1439	1,2-Distearoyl-phosphatidylethanolamine-methyl-polyethyleneglycol conjugate-2000 (Na⁺ salt) DSPE-MPEG-2000	100 mg	150.00
	$C_{43}H_{83}NO_{10}P(C_2H_4O)_n \cdot Na$	Mol. Wt.: 2807	CAS#: 147867-65-0
	Source: synthetic	Purity: 98+% by TLC	
	Appearance: solid	Solubility: chloroform	
	Storage: -20°C		
	Note: n ≈ 45		

Phosphatidylinositol Phosphates

1773	Phosphatidylinositol 3-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	95.00
1773-1	DPPI-3-P; PI-3-P dipalmitoyl (NH ₄ ⁺ salt)	1 mg	550.00
	$C_{41}H_{77}O_{16}P_2 \cdot 3NH_4$	Mol. Wt.: 942	CAS#: 165689-81-6
	Source: synthetic	Purity: 98+% by ¹ H NMR, ³¹ P NMR	
	Appearance: solid	Solubility: chloroform/methanol/DI water, 1:1:0.3	
	Storage: -20°C		


1775	Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl, (Na⁺ salt)	100 µg	60.00
1775-1	DPPI-3,4,5-P3; PI-3,4,5-P3 dipalmitoyl (Na ⁺ salt)	1 mg	375.00
	$C_{41}H_{75}O_{22}P_4 \cdot 7Na$	Mol. Wt.: 1205	CAS#: 165689-82-7
	Source: synthetic	Purity: 98+% by ¹ H NMR, ³¹ P NMR	
	Appearance: solid	Solubility: DI water	
	Storage: -20°C		

Bacterial Tetraethers

1303	Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i>, (>95% pure) <i>beta</i> -L-gulopyranosyl-caldarchaetidyl-glycerol	5 mg	1,150.00
	$C_{95}H_{188}O_{16}P$	Mol. Wt.: 1618	
	Source: natural, Archaeobacteria	Purity: >95% by TLC, HPLC	
	Appearance: solid	Solubility: chloroform/methanol, 2:1; hexane/2-propanol/DI water, 30:40:5	
	Storage: 4-8°C		

1303-2	Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i>, (>50% pure) <i>beta</i> -L-gulopyranosyl-caldarchaetidyl-glycerol	50 mg	3,025.00
	$C_{95}H_{188}O_{16}P$	Mol. Wt.: 1618	
	Source: natural, Archaeobacteria	Purity: >50% by TLC	
	Appearance: solid	Solubility: chloroform/methanol, 2:1; hexane/2-propanol/DI water, 30:40:5	
	Storage: 4-8°C	highly hygroscopic	

Glycosyl Glycerides



1058	Monogalactosyldiglyceride (hydrogenated), plant MGDG (hydrogenated)	10 mg	275.00
	$C_{45}H_{86}O_{10}$ Source: natural, plant Appearance: solid Storage: $-20^{\circ}C$ See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 787 (stearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 4:1:0.1	CAS#: 41670-62-6 Identity: confirmed by MS
1059	Digalactosyldiglyceride (hydrogenated), plant DGDG (hydrogenated)	5 mg	275.00
	$C_{51}H_{96}O_{15}$ Source: natural, plant Appearance: solid Storage: $-20^{\circ}C$ See Table III (pg. 98) for typical fatty acid content of products prepared from natural sources.	Mol. Wt.: 949 (distearoyl) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 4:1:0.1	CAS#: 92457-02-8 Identity: confirmed by MS

Fatty Acids

Simple Fatty Acids

Saturated Fatty Acids and Methyl Esters

These products are 99% pure by GC. They are stable at room temperature and are supplied neat.

1200	Methyl hexanoate Methyl caproate; C6:0 Methyl ester	1 g	60.00
	$C_7H_{14}O_2$ Source: natural, plant Appearance: liquid Storage: room temperature	Mol. Wt.: 130 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 106-70-7
1196	Heptanoic acid C7:0 Fatty acid	1 g	95.00
	$C_7H_{14}O_2$ Source: natural, plant Appearance: liquid Storage: room temperature	Mol. Wt.: 130 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 111-14-8
1197	Methyl heptanoate C7:0 Methyl ester	1 g	95.00
	$C_8H_{16}O_2$ Source: natural, plant Appearance: liquid Storage: room temperature	Mol. Wt.: 144 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 106-73-0

1198	Octanoic acid Caprylic acid; C8:0 Fatty acid		1 g	60.00
	$C_8H_{16}O_2$ Source: natural, plant Appearance: liquid Storage: room temperature	Mol. Wt.: 144 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 124-07-2
1199	Methyl octanoate Methyl caprylate; C8:0 Methyl ester		1 g	60.00
	$C_9H_{18}O_2$ Source: natural, plant Appearance: liquid Storage: room temperature	Mol. Wt.: 158 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 111-11-5
1163	Nonanoic acid C9:0 Fatty acid; Pelargonic acid		100 mg	50.00
	$C_9H_{18}O_2$ Source: synthetic Appearance: liquid Storage: room temperature	Mol. Wt.: 158 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 112-05-0
1164	Methyl nonanoate C9:0 Methyl ester		100 mg	50.00
	$C_{10}H_{20}O_2$ Source: synthetic Appearance: liquid Storage: room temperature	Mol. Wt.: 172 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 1731-84-6
1261	Methyl decanoate Methyl caprate; C10:0 Methyl ester		500 mg	50.00
	$C_{11}H_{22}O_2$ Source: natural, plant Appearance: liquid Storage: room temperature	Mol. Wt.: 186 Purity: 99% by TLC, GC Solubility: chloroform, hexane		CAS#: 110-42-9
1165	Undecanoic acid C11:0 Fatty acid		100 mg	50.00
	$C_{11}H_{22}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 186 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 112-37-8
1166	Methyl undecanoate C11:0 Methyl ester		100 mg	50.00
	$C_{12}H_{24}O_2$ Source: synthetic Appearance: liquid Storage: room temperature	Mol. Wt.: 200 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 1731-86-8
1008	Dodecanoic acid Lauric acid; C12:0 Fatty acid		1 g	70.00
	$C_{12}H_{24}O_2$ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 200 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 143-07-7

1009	Methyl dodecanoate Methyl laurate; C12:0 Methyl ester		1 g	70.00
	$C_{13}H_{26}O_2$ Source: natural, plant Appearance: liquid Storage: room temperature	Mol. Wt.: 214 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 111-82-0
1161	Tridecanoic acid C13:0 Fatty acid		100 mg	85.00
	$C_{13}H_{26}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 214 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 638-53-9
1162	Methyl tridecanoate C13:0 Methyl ester		100 mg	85.00
	$C_{14}H_{28}O_2$ Source: synthetic Appearance: liquid Storage: room temperature	Mol. Wt.: 228 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 1731-88-0
1010	Tetradecanoic acid Myristic acid; C14:0 Fatty acid		1 g	70.00
	$C_{14}H_{28}O_2$ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 228 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 544-63-8
1011	Methyl tetradecanoate Methyl myristate; C14:0 Methyl ester		1 g	70.00
	$C_{15}H_{30}O_2$ Source: natural, plant Appearance: liquid Storage: room temperature	Mol. Wt.: 242 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 124-10-7
1012	Pentadecanoic acid C15:0 Fatty acid		1 g	145.00
	$C_{15}H_{30}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 242 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 1002-84-2
1013	Methyl pentadecanoate C15:0 Methyl ester		1 g	145.00
	$C_{16}H_{32}O_2$ Source: synthetic Appearance: liquid Storage: room temperature	Mol. Wt.: 256 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 7132-64-1
1014	Hexadecanoic acid Palmitic acid; C16:0 Fatty acid		1 g	60.00
	$C_{16}H_{32}O_2$ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 256 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether		CAS#: 57-10-3

1015	Methyl hexadecanoate Methyl palmitate; C16:0 Methyl ester	1 g	60.00
	$C_{17}H_{34}O_2$ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 270 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 112-39-0
1018	Heptadecanoic acid Margaric acid; C17:0 Fatty acid	1 g	200.00
	$C_{17}H_{34}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 270 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 506-12-7
1019	Methyl heptadecanoate Methyl margarate; C17:0 Methyl ester	1 g	200.00
	$C_{18}H_{36}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 284 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 1731-92-6
1020	Octadecanoic acid Stearic acid; C18:0 Fatty acid	1 g	75.00
	$C_{18}H_{36}O_2$ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 284 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 57-11-4
1021	Methyl octadecanoate Methyl stearate; C18:0 Methyl ester	1 g	75.00
	$C_{19}H_{38}O_2$ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 298 Purity: 99% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 112-61-8
1028	Nonadecanoic acid C19:0 Fatty acid	100 mg	75.00
	$C_{19}H_{38}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 298 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 646-30-0
1029	Methyl nonadecanoate C19:0 Methyl ester	100 mg	75.00
	$C_{20}H_{40}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 312 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 1731-94-8
1030	Eicosanoic acid Arachidic acid; C20:0 Fatty acid	500 mg	100.00
	$C_{20}H_{40}O_2$ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 312 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 506-30-9

1031	Methyl eicosanoate Methyl arachidate; C20:0 Methyl ester	500 mg	100.00
	C ₂₁ H ₄₂ O ₂ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 327 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 1120-28-1
1241	Heneicosanoic acid C21:0 Fatty acid	100 mg	95.00
	C ₂₁ H ₄₂ O ₂ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 326 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 2363-71-5
1242	Methyl heneicosanoate C21:0 Methyl ester	100 mg	95.00
	C ₂₂ H ₄₄ O ₂ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 341 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 6064-90-0
1035	Docosanoic acid Behenic acid; C22:0 Fatty acid	500 mg	95.00
	C ₂₂ H ₄₄ O ₂ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 341 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 112-85-6
1036	Methyl docosanoate Methyl behenate; C22:0 Methyl ester	500 mg	95.00
	C ₂₃ H ₄₆ O ₂ Source: natural, plant Appearance: solid Storage: room temperature	Mol. Wt.: 354 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 929-77-1
1186	Tricosanoic acid C23:0 Fatty acid	100 mg	115.00
	C ₂₃ H ₄₆ O ₂ Source: semisynthetic, plant Appearance: solid Storage: room temperature	Mol. Wt.: 355 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 2433-96-7
1187	Methyl tricosanoate C23:0 Methyl ester	100 mg	115.00
	C ₂₄ H ₄₈ O ₂ Source: semisynthetic, plant Appearance: solid Storage: room temperature	Mol. Wt.: 368 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 2433-97-8
1037	Tetracosanoic acid Lignoceric acid; C24:0 Fatty acid	100 mg	90.00
	C ₂₄ H ₄₈ O ₂ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 369 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 557-59-5

1038	Methyl tetracosanoate Methyl lignocerate; C24:0 Methyl ester	100 mg	90.00
	$C_{25}H_{50}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 382 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 2442-49-1
1251	Hexacosanoic acid Cerotic acid; C26:0 Fatty acid	25 mg	130.00
	$C_{26}H_{52}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 397 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 506-46-7
1252	Methyl hexacosanoate Methyl cerotate; C26:0 Methyl ester	25 mg	130.00
	$C_{27}H_{54}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 411 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether	CAS#: 5802-82-4
1271	Methyl octacosanoate Methyl montanate; C28:0 Methyl ester	50 mg	155.00
	$C_{29}H_{58}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 439 Purity: 98+% by TLC, GC Solubility: chloroform, methylene chloride	CAS#: 55682-92-3
1273	Methyl triacontanoate Methyl melissate; C30:0 Methyl ester	50 mg	170.00
	$C_{31}H_{62}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 467 Purity: 98+% by TLC, GC Solubility: chloroform, methylene chloride	CAS#: 629-83-4
1275	Methyl dotriacontanoate Methyl lacceroate; C32:0 Methyl ester	50 mg	170.00
	$C_{33}H_{66}O_2$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 495 Purity: 98+% by TLC, GC Solubility: chloroform, methylene chloride	CAS#: 41755-79-7

Unsaturated Fatty Acids and Methyl Esters

Unsaturated fatty acids are easily oxidized. Flush open containers with argon or nitrogen and store at -20°C, in dark.

1157	Tetradecenoic acid (<i>cis</i>-9) Myristoleic acid; C14:1 (<i>cis</i> -9) Fatty acid	100 mg	130.00
	$C_{14}H_{26}O_2$ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 226 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 544-64-9

1040	Methyl tetradecenoate (<i>cis</i>-9) Methyl myristoleate; C14:1 (<i>cis</i> -9) Methyl ester	100 mg	130.00
	C ₁₅ H ₂₈ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 240 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 56219-06-8
1209	Pentadecenoic acid (<i>cis</i>-10) C15:1 (<i>cis</i> -10) Fatty acid	50 mg	100.00
	C ₁₅ H ₂₈ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 240 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 84743-29-3
1210	Methyl pentadecenoate (<i>cis</i>-10) C15:1 (<i>cis</i> -10) Methyl ester	50 mg	100.00
	C ₁₆ H ₃₀ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 254 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 90176-52-6
1243	Hexadecenoic acid (<i>cis</i>-6) Sapienic acid, C16:1 (<i>cis</i> -6) Fatty acid	25 mg	240.00
	C ₁₆ H ₃₀ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 254 Purity: 99% by TLC, GC Solubility: ethanol, methanol, chloroform, ethyl ether	CAS#: 17004-51-2 Identity: confirmed by MS
1016	Hexadecenoic acid (<i>cis</i>-9) Palmitoleic acid; C16:1 (<i>cis</i> -9) Fatty acid	100 mg	80.00
	C ₁₆ H ₃₀ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 254 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 373-49-9
1017	Methyl hexadecenoate (<i>cis</i>-9) Methyl palmitoleate; C16:1 (<i>cis</i> -9) Methyl ester	100 mg	80.00
	C ₁₇ H ₃₂ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 268 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 1120-25-8
1147	Hexadecenoic acid (<i>trans</i>-9) Palmitelaidic acid; C16:1 (<i>trans</i> -9) Fatty acid	100 mg	105.00
	C ₁₆ H ₃₀ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 254 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 10030-73-6
1148	Methyl hexadecenoate (<i>trans</i>-9) Methyl palmitelaidate; C16:1 (<i>trans</i> -9) Methyl ester	100 mg	105.00
	C ₁₇ H ₃₂ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 268 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 10030-74-7

1208	11-Hexadecenoic acid (92% <i>cis</i>, 8% <i>trans</i>) C16:1 (<i>cis</i> -11/ <i>trans</i> -11) Fatty acid	50 mg	185.00
	C ₁₆ H ₃₀ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 254 Purity: >98%, by TLC Solubility: chloroform, ethanol, hexane, methanol	CAS#: 2271-34-3 Identity: confirmed by MS
1204	Heptadecenoic acid (<i>cis</i>-10) C17:1 (<i>cis</i> -10) Fatty acid	100 mg	100.00
	C ₁₇ H ₃₂ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 268 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 29743-97-3
1203	Methyl heptadecenoate (<i>cis</i>-10) C17:1 (<i>cis</i> -10) Methyl ester	100 mg	100.00
	C ₁₈ H ₃₄ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 282 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 75190-82-8
1022	Octadecenoic acid (<i>cis</i>-9) Oleic acid; C18:1 (<i>cis</i> -9) Fatty acid	1 g	75.00
	C ₁₈ H ₃₄ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 282 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 112-80-1
1023	Methyl octadecenoate (<i>cis</i>-9) Methyl oleate; C18:1 (<i>cis</i> -9) Methyl ester	1 g	75.00
	C ₁₉ H ₃₆ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 296 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 112-62-9
1149	Octadecenoic acid (<i>trans</i>-9) Elaidic acid; C18:1 (<i>trans</i> -9) Fatty acid	1 g	145.00
	C ₁₈ H ₃₄ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 282 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 112-79-8
1150	Methyl octadecenoate (<i>trans</i>-9) Methyl elaidate; C18:1 (<i>trans</i> -9) Methyl ester	1 g	145.00
	C ₁₉ H ₃₆ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 296 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 1937-62-8
1266	Octadecenoic acid (<i>cis</i>-11) <i>cis</i> -Vaccenic acid; C18:1(<i>cis</i> -11) Fatty acid	100 mg	130.00
	C ₁₈ H ₃₄ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 282 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 506-17-2

1267	Methyl octadecenoate (<i>cis</i>-11) Methyl <i>cis</i> -vaccenate; C18:1(<i>cis</i> -11) Methyl ester	100 mg	130.00
	C ₁₉ H ₃₆ O ₂ Source: semisynthetic, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 296 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 1937-63-9
1262	Octadecenoic acid (<i>trans</i>-11) <i>trans</i> -Vaccenic acid; C18:1 (<i>trans</i> -11) Fatty acid	100 mg	130.00
	C ₁₈ H ₃₄ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 282 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 693-72-1
1263	Methyl octadecenoate (<i>trans</i>-11) Methyl <i>trans</i> -vaccenate; C18:1 (<i>trans</i> -11) Methyl ester	100 mg	130.00
	C ₁₉ H ₃₆ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 296 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 6198-58-9
1024	Octadecadienoic acid (all <i>cis</i>-9,12) Linoleic acid; C18:2 (all <i>cis</i> -9,12) Fatty acid	1 g	75.00
	C ₁₈ H ₃₂ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 280 Purity: 99% by TLC, GC Solubility: ethyl ether, ethanol, hexane	CAS#: 60-33-3
1025	Methyl octadecadienoate (all <i>cis</i>-9,12) Methyl linoleate; C18:2 (all <i>cis</i> -9,12) Methyl ester	1 g	75.00
	C ₁₉ H ₃₄ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 294 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 112-63-0
1151	Octadecadienoic acid (all <i>trans</i>-9,12) C18:2 (all <i>trans</i> -9,12) Fatty acid; Linoelaidic acid (all <i>trans</i> -9,12)	100 mg	80.00
	C ₁₈ H ₃₂ O ₂ Source: semisynthetic, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 280 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 506-21-8
1152	Methyl octadecadienoate (all <i>trans</i>-9,12) Methyl linoelaidate; C18:2 (all <i>trans</i> -9,12) Methyl ester	100 mg	80.00
	C ₁₉ H ₃₄ O ₂ Source: semisynthetic, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 294 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 2566-97-4
1026	Octadecatrienoic acid (all <i>cis</i>-9,12,15) Linolenic acid; C18:3 (all <i>cis</i> -9,12,15) Fatty acid	500 mg	155.00
	C ₁₈ H ₃₀ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 278 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 463-40-1

1027	Methyl octadecatrienoate (all <i>cis</i>-9,12,15) Methyl linolenate; C18:3 (all <i>cis</i> -9,12,15) Methyl ester	500 mg	155.00
	C ₁₉ H ₃₂ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 292 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 301-00-8
1153	Octadecatrienoic acid (all <i>cis</i>-6,9,12) <i>gamma</i> -Linolenic acid; C18:3 (all <i>cis</i> -6,9,12) Fatty acid	100 mg	90.00
	C ₁₈ H ₃₀ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 278 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 506-26-3
1154	Methyl octadecatrienoate (all <i>cis</i>-6,9,12) Methyl <i>gamma</i> -linolenate; C18:3 (all <i>cis</i> -6,9,12) Methyl ester	100 mg	90.00
	C ₁₉ H ₃₂ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 292 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 16326-32-2
1240 *NEW*	Methyl punicate Methyl 9(Z),11(E),13(Z)-Octadecatrienoate; Conjugated linolenic acid methyl ester; CLnA	25 mg	185.00
	C ₁₉ H ₃₂ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 292 Purity: 97+% by TLC, GC Solubility: hexane, ethanol, methanol, chloroform	Identity: confirmed by MS
1234 *NEW*	Methyl jacarate Methyl 8(Z),10(E),12(Z)-Octadecatrienoate; Jacaric acid methyl ester; Conjugated linolenic acid methyl ester; CLnA	25 mg	325.00
	C ₁₉ H ₃₂ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 293 Purity: 98+% by TLC, 96+% by GC Solubility: hexane, ethanol, methanol, chloroform	Identity: confirmed by MS
1233 *NEW*	Methyl <i>alpha</i>-eleostearate Methyl 9(Z),11(E),13(E)-octadecatrienoate; <i>alpha</i> -Eleostearic acid methyl ester; Conjugated linolenic acid methyl ester; CLnA	25 mg	325.00
	C ₁₉ H ₃₂ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 293 Purity: 98+% by TLC, GC Solubility: hexane, ethanol, methanol, chloroform	Identity: confirmed by MS
1276	Stearidonic acid (all <i>cis</i>-6,9,12,15) Morotic acid; C18:4 (all <i>cis</i> -6,9,12,15) Fatty acid	25 mg	375.00
	C ₁₈ H ₂₈ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 276 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 20290-75-9 Identity: confirmed by MS
1277	Methyl stearidonate (all <i>cis</i>-6,9,12,15) Morotic acid methyl ester; C18:4 (all <i>cis</i> -6,9,12,15) Methyl ester	25 mg	375.00
	C ₁₉ H ₃₀ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 290 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 73097-00-4 Identity: confirmed by MS

1205	Nonadecenoic acid (<i>cis</i>-10) C19:1 (<i>cis</i> -10) Fatty acid	100 mg	120.00
	C ₁₉ H ₃₆ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 296 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 73033-09-7
1206	Methyl nonadecenoate (<i>cis</i>-10) C19:1 (<i>cis</i> -10) Methyl ester	100 mg	120.00
	C ₂₀ H ₃₈ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 310 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 19788-74-0
1032	Eicosenoic acid (<i>cis</i>-11) C20:1 (<i>cis</i> -11) Fatty acid; Gondoic acid	100 mg	90.00
	C ₂₀ H ₃₈ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 310 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 5561-99-9
1033	Methyl eicosenoate (<i>cis</i>-11) Methyl eicosenoate; C20:1 (<i>cis</i> -11) Methyl ester; Methyl gondoate	100 mg	90.00
	C ₂₁ H ₄₀ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 324 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 2390-09-2
1192	Eicosadienoic acid (all <i>cis</i>-11,14) C20:2 (all <i>cis</i> -11,14) Fatty acid	100 mg	90.00
	C ₂₀ H ₃₆ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 309 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 2091-39-6
1193	Methyl eicosadienoate (all <i>cis</i>-11,14) Methyl eicosadienoate; C20:2 (all <i>cis</i> -11,14) Methyl ester	100 mg	90.00
	C ₂₁ H ₃₈ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 322 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 2463-02-7
1179	Methyl eicosatrienoate (all <i>cis</i>-5,8,11) Mead acid methyl ester; C20:3 (all <i>cis</i> -5,8,11) Methyl ester	1 mg/ml, 1 ml	250.00
	C ₂₁ H ₃₆ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 320 Purity: 90% by TLC, GC Solubility: chloroform, hexane, ethyl ether Solvent: hexane	CAS#: 14602-39-2
1269	Methyl eicosatrienoate (all <i>cis</i>-8,11,14) Methyl homogamma linolenate; C20:3 (all <i>cis</i> -8,11,14) Methyl ester	50 mg	140.00
	C ₂₁ H ₃₆ O ₂ Source: semisynthetic, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 321 Purity: 99% by TLC, GC Solubility: hexane, ethyl ether, chloroform	CAS#: 21061-10-9

1042	Eicosatetraenoic acid (all <i>cis</i>-5,8,11,14) Arachidonic acid (all <i>cis</i> -5,8,11,14); C20:4 (all <i>cis</i> -5,8,11,14) Fatty acid	100 mg	125.00
	$C_{20}H_{32}O_2$ Source: natural, fungal Appearance: liquid Storage: -20°C	Mol. Wt.: 304 Purity: 99% by TLC, GC Solubility: ethyl ether, hexane, methylene chloride	CAS#: 506-32-1
1034	Methyl eicosatetraenoate (all <i>cis</i>-5,8,11,14) Methyl arachidonate; C20:4 (all <i>cis</i> -5,8,11,14) Methyl ester	100 mg	125.00
	$C_{21}H_{34}O_2$ Source: natural, fungal Appearance: liquid Storage: -20°C	Mol. Wt.: 318 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 2566-89-4
1167	Eicosapentaenoic acid (all <i>cis</i>-5,8,11,14,17) EPA ; <i>omega</i> -3 Fatty acid; C20:5 (all <i>cis</i> -5,8,11,14,17) fatty acid	25 mg	135.00
	$C_{20}H_{30}O_2$ Source: natural, fish oil Appearance: liquid Storage: -20°C Anti-hyperlipoproteinemic agent; 5-LOX inhibitor	Mol. Wt.: 302 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether, ethanol, DMSO, DMF	CAS#: 10417-94-4
1194	Methyl eicosapentaenoate (all <i>cis</i>-5,8,11,14,17) Methyl ester of <i>omega</i> -3 Fatty acid; C20:5 (all <i>cis</i> -5,8,11,14,17) Methyl ester	25 mg	135.00
	$C_{21}H_{32}O_2$ Source: natural, fish oil Appearance: liquid Storage: -20°C	Mol. Wt.: 316 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether, hexane	CAS#: 2734-47-6
1264	Docosenoic acid (<i>cis</i>-13) Erucic acid; C22:1 (<i>cis</i> -13) Fatty acid	100 mg	70.00
	$C_{22}H_{42}O_2$ Source: natural, plant Appearance: solid Storage: -20°C	Mol. Wt.: 339 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether, hexane	CAS#: 112-86-7
1265	Methyl docosenoate (<i>cis</i>-13) Methyl erucate; C22:1 (<i>cis</i> -13) Methyl ester	100 mg	70.00
	$C_{23}H_{44}O_2$ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 352 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether, hexane	CAS#: 1120-34-9
1175	Docosapentaenoic acid (all <i>cis</i>-7,10,13,16,19) C22:5 (all <i>cis</i> -7,10,13,16,19) Fatty acid	25 mg	135.00
	$C_{22}H_{34}O_2$ Source: semisynthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 330 Purity: 99% by TLC, GC Solubility: chloroform, ethyl ether, hexane	CAS#: 24880-45-3
1244	Methyl docosapentaenoate (all <i>cis</i>-7,10,13,16,19) C22:5 (all <i>cis</i> -7,10,13,16,19) Methyl ester	25 mg	135.00
	$C_{23}H_{36}O_2$ Source: semisynthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 344 Purity: 99% by TLC, GC Solubility: ethyl ether, ethanol, hexane	CAS#: 108698-02-8

1136	Docosahexaenoic acid (all <i>cis</i>-4,7,10,13,16,19) DHA; C22:6 (all <i>cis</i> -4,7,10,13,16,19) <i>omega</i> -3 Fatty acid	100 mg	125.00
	C ₂₂ H ₃₂ O ₂ Source: natural, algae Appearance: liquid Storage: -20°C	Mol. Wt.: 328 Purity: 99% by TLC, GC Solubility: ethyl ether, hexane, methylene chloride, ethanol, DMSO, DMF	CAS#: 6217-54-5
1041	Methyl docosahexaenoate (all <i>cis</i>-4,7,10,13,16,19) Methyl ester of <i>omega</i> -3 fatty acid; C22:6 (all <i>cis</i> -4,7,10,13,16,19) Methyl ester	100 mg	125.00
	C ₂₃ H ₃₄ O ₂ Source: natural, algae Appearance: liquid Storage: -20°C	Mol. Wt.: 342 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 2566-90-7
1155	Tetracosenoic acid (<i>cis</i>-15) Nervonic acid (<i>cis</i> -15); C24:1 (<i>cis</i> -15) Fatty acid	100 mg	125.00
	C ₂₄ H ₄₆ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 367 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 506-37-6
1156	Methyl tetracosenoate (<i>cis</i>-15) Methyl nervonate; C24:1 (<i>cis</i> -15) Methyl ester	100 mg	125.00
	C ₂₅ H ₄₈ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 381 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 2733-88-2

Trans Fatty Acids and Methyl Esters

1147	Hexadecenoic acid (<i>trans</i>-9) Palmitelaidic acid; C16:1 (<i>trans</i> -9) Fatty acid	100 mg	105.00
	C ₁₆ H ₃₀ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 254 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 10030-73-6
1148	Methyl hexadecenoate (<i>trans</i>-9) Methyl palmitelaidate; C16:1 (<i>trans</i> -9) Methyl ester	100 mg	105.00
	C ₁₇ H ₃₂ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 268 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 10030-74-7
1149	Octadecenoic acid (<i>trans</i>-9) Elaidic acid; C18:1 (<i>trans</i> -9) Fatty acid	1 g	145.00
	C ₁₈ H ₃₄ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 282 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 112-79-8

1150	Methyl octadecenoate (<i>trans</i>-9) Methyl elaidate; C18:1 (<i>trans</i> -9) Methyl ester	1 g	145.00
	C ₁₉ H ₃₆ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 296 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 1937-62-8
1262	Octadecenoic acid (<i>trans</i>-11) <i>trans</i> Vaccenic acid; C18:1 (<i>trans</i> -11) Fatty acid	100 mg	130.00
	C ₁₈ H ₃₄ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 282 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 693-72-1
1263	Methyl octadecenoate (<i>trans</i>-11) Methyl <i>trans</i> vaccenate; C18:1 (<i>trans</i> -11) Methyl ester	100 mg	130.00
	C ₁₉ H ₃₆ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 296 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 6198-58-9
1151	Octadecadienoic acid (all <i>trans</i>-9,12) C18:2 (all <i>trans</i> -9,12) Fatty acid; Linoelaidic acid (all <i>trans</i> -9,12)	100 mg	80.00
	C ₁₈ H ₃₂ O ₂ Source: semisynthetic, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 280 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 506-21-8
1152	Methyl octadecadienoate (all <i>trans</i>-9,12) Methyl linoelaidate; C18:2 (all <i>trans</i> -9,12) Methyl ester	100 mg	80.00
	C ₁₉ H ₃₄ O ₂ Source: semisynthetic, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 294 Purity: 99% by TLC, GC Solubility: chloroform, hexane, ethyl ether	CAS#: 2566-97-4
1131	<i>Cis-Trans</i> FAME Isomer Standard Mixture	5.5 mg/ml, 5 ml	170.00
	Source: margarine Appearance: liquid Storage: -20°C	Solubility: methylene chloride, chloroform Solvent: methylene chloride	

Analysis of positional *cis-trans* fatty acid isomers is ever more important in light of the new food industry rules. These isomers can be resolved on Supelco SP-2560 or an equivalent capillary GC column. Use this specially formulated mix to ensure proper operation of your column for this tricky separation. Mix consists of *cis-trans* fatty acid isomers as methyl esters in methylene chloride.

This is a qualitative standard containing in order of elution: C16:0, C18:0, C18:1 *trans* isomers (4 peaks), C18:1 *cis* & *trans* isomers (2 peaks), C18:1 *cis* isomers (4 peaks), C18:2, C20:0, C20:1 and C18:3 (same peak), C22:0

Conjugated Linoleic Acid Isomers (CLA)


Linoleic acid is an essential fatty acid (18:2 ω6) of which several naturally occurring conjugated derivatives have been identified. These derivatives, called “conjugated linoleic acid” or CLA can have the two double bonds mainly in the 9 and 11 or in the 10 and 12 positions, resulting in eight possible geometric isomers. CLA occurs in meat (41) and dairy products (42,43). In both cases, the 9(Z),11(E)-isomer is predominant and is thought to be the biologically active form. CLA assimilated through the diet of animals is found in the intestinal mucosa, liver and adipose tissue (44). See also review article by Parodi (43). CLA has several biological properties. It's anti-carcinogenic activity has been demonstrated by its ability to inhibit chemically induced tumor formation in animal models of carcinogenesis (41,45-47). The addition of CLA to culture medium suppresses the *in vitro* growth of human melanoma, colorectal and breast cancer cells (48). CLA also exhibits anti-atherogenic activity. Addition of CLA to a controlled atherogenic diet significantly reduced the development of atherosclerosis in hamsters and rabbits (49,50). Animals fed a diet containing CLA also had lower levels of low-density lipoprotein (LDL) cholesterol. CLA may be involved in regulating fat and protein metabolism (51,52). Several species of animals fed CLA-supplemented diets showed improved feed efficiency. Lean body mass increased while body fat was reduced. This seems to be due, mainly or exclusively, to the 10(E),12(Z)-isomer (catalog # 1249, see below). CLA competes with linoleate for Δ6 desaturase (53). Dietary CLA normalizes impaired glucose tolerance in the Zucker diabetic fatty *fa/fa* rat (54) *via* activation of PPAR γ, a result which bears on the possible ameliorization or prevention of NIDDM. The 11(Z),13(E)-isomer has been shown to be concentrated in the heart and in mitochondria.

See Literature References on page 106.


CLA Research is Being Redone With Our Highly Pure Isomers

Most studies to date have utilized a mixture of CLA isomers containing less than 30% of the presumed active 9(Z),11(E)-isomer (55,56). In addition to the 9,11- and 10,12-isomers, 8,10- and 11,13-isomers have recently been identified in the widely used mixture (56,57). Matreya offers a highly pure CLA which is 98+% the active 9,11-“cis, trans” isomer. The corresponding “trans,trans” and “cis,cis” isomers are also available. In addition, we now offer the pure 10(E),12(Z)-isomer, which has been widely sought for comparison studies.

See Literature References on page 106.



1245	9(Z),11(E)-Octadecadienoic acid 9- <i>cis</i> , 11- <i>trans</i> CLA		25 mg	155.00
	C ₁₈ H ₃₂ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 280 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, hexane, methanol, DMSO		CAS#: 2540-56-9 Identity: confirmed by MS
1278	9(Z),11(E)-Octadecadienoic acid (Na⁺ salt) 9- <i>cis</i> ,11- <i>trans</i> -CLA (Na ⁺ salt)		25 mg	165.00
	C ₁₈ H ₃₁ NaO ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 302 Purity: 98+% by TLC, GC Solubility: water, methanol, DMSO		Identity: confirmed by MS
1255	Methyl 9(Z),11(E)-octadecadienoate Methyl ester of CLA (9- <i>cis</i> , 11- <i>trans</i>)		25 mg	155.00
	C ₁₉ H ₃₄ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 294 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, hexane, methanol		CAS#: 13058-52-1 Identity: confirmed by MS

1181	9(E),11(E)-Octadecadienoic acid 9- <i>trans</i> , 11- <i>trans</i> CLA	25 mg	155.00
	C ₁₈ H ₃₂ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 280 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, hexane, methanol Melting Point (°C): 55-57	CAS#: 544-71-8 Identity: confirmed by MS
1257	Methyl 9(E),11(E)-octadecadienoate Methyl ester of CLA (9- <i>trans</i> , 11- <i>trans</i>)	25 mg	155.00
	C ₁₉ H ₃₄ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 294 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, hexane, methanol	CAS#: 13038-47-6 Identity: confirmed by MS
1248	9(Z),11(Z)-Octadecadienoic acid 9- <i>cis</i> , 11- <i>cis</i> CLA	25 mg	155.00
	C ₁₈ H ₃₂ O ₂ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 280 Purity: 96+% by TLC, GC Solubility: chloroform, ethanol, methanol, ethyl ether Melting Point (°C): 40-42	CAS#: 544-40-7 Identity: confirmed by MS
1256	Methyl 9(Z),11(Z)-octadecadienoate Methyl ester of CLA (9- <i>cis</i> , 11- <i>cis</i>)	25 mg	155.00
	C ₁₉ H ₃₄ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 294 Purity: 96+% by TLC, GC Solubility: chloroform, ethanol, hexane, methanol	
			
1249	10(E),12(Z)-Octadecadienoic acid 10- <i>trans</i> , 12- <i>cis</i> CLA	25 mg	155.00
	C ₁₈ H ₃₂ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 280 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, hexane, methanol	CAS#: 2420-44-2 Identity: confirmed by MS
1254	Methyl 10(E),12(Z)-octadecadienoate Methyl ester of CLA (10- <i>trans</i> , 12- <i>cis</i>)	25 mg	155.00
	C ₁₉ H ₃₄ O ₂ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 294 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, hexane, methanol	CAS#: 21870-97-3
1240 *NEW*	Methyl punicate Methyl 9(Z),11(E),13(Z)-Octadecatrienoate; Conjugated linolenic acid methyl ester; CLnA	25 mg	185.00
	C ₁₉ H ₃₂ O ₂ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 292 Purity: 97+% by TLC, GC Solubility: hexane, ethanol, methanol, chloroform	Identity: confirmed by MS


1234 *NEW*	Methyl jacarate Methyl 8(Z),10(E),12(Z)-Octadecatrienoate; Jacaric acid methyl ester; Conjugated linolenic acid methyl ester; CLnA	25 mg	325.00
	$C_{19}H_{32}O_2$ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 293 Purity: 98+% by TLC, 96+% by GC Solubility: hexane, ethanol, methanol, chloroform	Identity: confirmed by MS

1233 *NEW*	Methyl <i>alpha</i>-eleostearate Methyl 9(Z),11(E),13(E)-octadecatrienoate; <i>alpha</i> -Eleostearic acid methyl ester; Conjugated linolenic acid methyl ester; CLnA	25 mg	325.00
	$C_{19}H_{32}O_2$ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 293 Purity: 98+% by TLC, GC Solubility: hexane, ethanol, methanol, chloroform	Identity: confirmed by MS

Hydroxy Fatty Acids

2-Hydroxy Fatty Acids and Methyl Esters

These products are racemic and 98+% pure by GC and TLC. The 2-hydroxy fatty acids are components of glycosphingolipids and are involved in fatty acid degradation. They are stable and are supplied neat in vials.



1758	2-Hydroxydecanoic acid 2-Hydroxy C10:0 fatty acid	50 mg	155.00
	$C_{10}H_{20}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 188 Purity: 98+% by TLC, GC Solubility: chloroform, methanol	CAS#: 5393-81-7 Identity: confirmed by MS

1759	Methyl 2-hydroxydecanoate 2-Hydroxy C10:0 methyl ester	50 mg	155.00
	$C_{11}H_{22}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 202 Purity: 98+% by TLC, GC Solubility: chloroform, methanol	CAS#: 71271-24-4


1701	2-Hydroxydodecanoic acid 2-Hydroxy C12:0 fatty acid	50 mg	155.00
	$C_{12}H_{24}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 216 Purity: 98+% by TLC, GC Solubility: chloroform, methanol	CAS#: 2984-55-6

1702	Methyl 2-hydroxydodecanoate 2-Hydroxy C12:0 methyl ester	50 mg	155.00
	$C_{13}H_{26}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 230 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether, methanol	CAS#: 51067-85-7
1703	2-Hydroxytetradecanoic acid 2-Hydroxy C14:0 fatty acid	50 mg	155.00
	$C_{14}H_{28}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 244 Purity: 98+% by TLC, GC Solubility: chloroform, methanol Melting Point (°C): 81-82	CAS#: 2507-55-3 Identity: confirmed by MS
1704	Methyl 2-hydroxytetradecanoate 2-Hydroxy C14:0 methyl ester	50 mg	155.00
	$C_{15}H_{30}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 258 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether, methanol	CAS#: 56009-40-6 Identity: confirmed by MS
1705	2-Hydroxyhexadecanoic acid 2-Hydroxy C16:0 fatty acid	50 mg	155.00
	$C_{16}H_{32}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 272 Purity: 98+% by TLC, GC Solubility: methanol, chloroform/methanol, 2:1 Melting Point (°C): 86-87	CAS#: 764-67-0
1706	Methyl 2-hydroxyhexadecanoate 2-Hydroxy C16:0 methyl ester	50 mg	155.00
	$C_{17}H_{34}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 286 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether, methanol Melting Point (°C): 59-60	CAS#: 16742-51-1
1707	2-Hydroxyoctadecanoic acid 2-Hydroxy C18:0 fatty acid	50 mg	155.00
	$C_{18}H_{36}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 300 Purity: 98+% by TLC, GC Solubility: chloroform/methanol, 5:1 Melting Point (°C): 92-93	CAS#: 629-22-1
1708	Methyl 2-hydroxyoctadecanoate 2-Hydroxy C18:0 methyl ester	50 mg	155.00
	$C_{19}H_{38}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 315 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether, methanol Melting Point (°C): 64-66	CAS#: 2420-35-1
1709	2-Hydroxyeicosanoic acid 2-Hydroxy C20:0 fatty acid	25 mg	155.00
	$C_{20}H_{40}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 329 Purity: 98+% by TLC, GC Solubility: chloroform/methanol, 5:1 Melting Point (°C): 91-92	CAS#: 16742-48-6

1710	Methyl 2-hydroxyeicosanoate 2-Hydroxy C20:0 methyl ester	25 mg	155.00
	C₂₁H₄₂O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 343 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether Melting Point (°C): 62-64	CAS#: 16742-49-7 Identity: confirmed by MS
1711	2-Hydroxydocosanoic acid 2-Hydroxy C22:0 fatty acid	25 mg	155.00
	C₂₂H₄₄O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 356 Purity: 98+% by TLC, GC Solubility: chloroform/methanol, 5:1 Melting Point (°C): 96-97	CAS#: 13980-14-8 Identity: confirmed by MS
1712	Methyl 2-hydroxydocosanoate 2-Hydroxy C22:0 methyl ester	25 mg	155.00
	C₂₃H₄₆O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 371 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether Melting Point (°C): 72-73	CAS#: 13980-17-1
1713	2-Hydroxytricosanoic acid 2-Hydroxy C23:0 fatty acid	10 mg	220.00
	C₂₃H₄₆O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 371 Purity: 98+% by TLC, GC Solubility: chloroform/methanol, 5:1 Melting Point (°C): 98-99	CAS#: 2718-37-8
1714	Methyl 2-hydroxytricosanoate 2-Hydroxy C23:0 methyl ester	10 mg	220.00
	C₂₄H₄₈O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 385 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether Melting Point (°C): 68-70	CAS#: 118745-41-8 Identity: confirmed by MS
1715	2-Hydroxytetracosanoic acid 2-Hydroxylignoceric; 2-Hydroxy C24:0 fatty acid; Cerebronic acid	5 mg	220.00
	C₂₄H₄₈O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 385 Purity: 98+% by TLC, GC Solubility: chloroform/methanol, 5:1 Melting Point (°C): 101-104	CAS#: 544-57-0
1716	Methyl 2-hydroxytetracosanoate Methyl 2-hydroxylignocerate; 2-Hydroxy C24:0 methyl ester	5 mg	220.00
	C₂₅H₅₀O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 399 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether Melting Point (°C): 77-80	CAS#: 2433-95-6
1722	2-Hydroxy Methyl Ester Mixture Quantitative mixture	10 mg/ml, 1 ml	180.00
	Source: synthetic Appearance: liquid Storage: -20°C Contains: 2-OH C14:0, 20.0%; 2-OH C16:0, 20.0%; 2-OH C18:0, 15.0%; 2-OH C20:0, 15.0%; 2-OH C22:0, 10.0%; 2-OH C23:0, 10.0%; 2-OH C24:0, 10.0%	Solubility: chloroform Solvent: chloroform	

3-Hydroxy Fatty Acids and Methyl Esters

These products are racemic and 98+% pure by GC and TLC. 3-Hydroxy fatty acids occur in the lipid fraction of many microorganisms and are useful in the typing of microbial isolates. They are stable and are supplied neat in vials.




1747	3-Hydroxyhexanoic acid 3-Hydroxy C6:0 fatty acid	25 mg	190.00
	C₆H₁₂O₃ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 132 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol	CAS#: 10191-24-9
1748	Methyl 3-hydroxyhexanoate 3-Hydroxy C6:0 methyl ester	25 mg	190.00
	C₇H₁₄O₃ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 146 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol	CAS#: 21188-58-9 Identity: confirmed by MS
1745	3-Hydroxyoctanoic acid 3-Hydroxy C8:0 fatty acid	25 mg	180.00
	C₈H₁₆O₃ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 160 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol	CAS#: 88930-08-9 Identity: confirmed by MS
1746	Methyl 3-hydroxyoctanoate 3-Hydroxy C8:0 methyl ester	25 mg	180.00
	C₉H₁₈O₃ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 174 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 85549-54-8 Identity: confirmed by MS
1725	3-Hydroxynonanoic acid 3-Hydroxy C9:0 fatty acid	25 mg	190.00
	C₉H₁₈O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 174 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol Melting Point (°C): 60-62	CAS#: 88930-09-0
1726	Methyl 3-hydroxynonanoate 3-Hydroxy C9:0 methyl ester	25 mg	190.00
	C₁₀H₂₀O₃ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 188 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 83968-06-3

1727	3-Hydroxydecanoic acid 3-Hydroxy C10:0 fatty acid	25 mg	220.00
	$C_{10}H_{20}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 188 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol Melting Point (°C): 57-60	CAS#: 5561-87-5 Identity: confirmed by MS
1728	Methyl 3-hydroxydecanoate 3-Hydroxy C10:0 methyl ester	25 mg	220.00
	$C_{11}H_{22}O_3$ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 202 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol	CAS#: 62675-82-5
1729	3-Hydroxyundecanoic acid 3-Hydroxy C11:0 fatty acid	25 mg	200.00
	$C_{11}H_{22}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 202 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol Melting Point (°C): 74-76	CAS#: 40165-88-6
1730	Methyl 3-hydroxyundecanoate 3-Hydroxy C11:0 methyl ester	25 mg	200.00
	$C_{12}H_{24}O_3$ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 216 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol	CAS#: 127593-21-9
1731	3-Hydroxydodecanoic acid 3-Hydroxy C12:0 fatty acid	25 mg	190.00
	$C_{12}H_{24}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 216 Purity: 98+% by TLC, GC Solubility: ethanol, methanol Melting Point (°C): 71-72	CAS#: 1883-13-2
1732	Methyl 3-hydroxydodecanoate 3-Hydroxy C12:0 methyl ester	25 mg	200.00
	$C_{13}H_{26}O_3$ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 230 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, ethyl ether	CAS#: 85464-97-7 Identity: confirmed by MS
1733	3-Hydroxytridecanoic acid 3-Hydroxy C13:0 fatty acid	25 mg	200.00
	$C_{13}H_{26}O_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 230 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol Melting Point (°C): 80-83	CAS#: 32602-69-0
1734	Methyl 3-hydroxytridecanoate 3-Hydroxy C13:0 methyl ester	25 mg	200.00
	$C_{14}H_{28}O_3$ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 244 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether	

1735	3-Hydroxytetradecanoic acid 3-Hydroxy C14:0 fatty acid	25 mg	190.00
	C₁₄H₂₈O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 244 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol Melting Point (°C): 80-81	CAS#: 3422-31-9 Identity: confirmed by MS
1736	Methyl 3-hydroxytetradecanoate 3-Hydroxy C14:0 methyl ester	25 mg	190.00
	C₁₅H₃₀O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 258 Purity: 98+% by TLC, GC Solubility: chloroform, ethyl ether, methanol Melting Point (°C): 36-37	CAS#: 55682-83-2 Identity: confirmed by MS
1739	3-Hydroxyhexadecanoic acid 3-Hydroxy C16:0 fatty acid	25 mg	190.00
	C₁₆H₃₂O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 272 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol Melting Point (°C): 85-86	CAS#: 928-17-6 Identity: confirmed by MS
1740	Methyl 3-hydroxyhexadecanoate 3-Hydroxy C16:0 methyl ester	25 mg	190.00
	C₁₇H₃₄O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 286 Purity: 98+% by TLC, GC Solubility: ethanol, methanol Melting Point (°C): 43-45	CAS#: 51883-36-4 Identity: confirmed by MS
1741	3-Hydroxyheptadecanoic acid 3-Hydroxy C17:0 fatty acid	25 mg	180.00
	C₁₇H₃₄O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 286 Purity: 98+% by TLC, GC Solubility: ethanol, methanol Melting Point (°C): 93-95	CAS#: 40165-89-7
1742	Methyl 3-hydroxyheptadecanoate 3-Hydroxy C17:0 methyl ester	25 mg	180.00
	C₁₈H₃₆O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 300 Purity: 98+% by TLC, GC Solubility: ethanol, methanol Melting Point (°C): 53-55	CAS#: 112538-92-8
1743	3-Hydroxyoctadecanoic acid 3-Hydroxy C18:0 fatty acid	25 mg	180.00
	C₁₈H₃₆O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 300 Purity: 98+% by TLC, GC Solubility: ethanol, methanol Melting Point (°C): 52-54	CAS#: 45261-96-9 Identity: confirmed by MS
1744	Methyl 3-hydroxyoctadecanoate 3-Hydroxy C18:0 methyl ester	25 mg	180.00
	C₁₉H₃₈O₃ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 314 Purity: 98+% by TLC, GC Solubility: ethanol, methanol Melting Point (°C): 52-54	CAS#: 14531-40-9

Omega Hydroxy Fatty Acids




1754	Royal Jelly acid 10-Hydroxy-2-(E)-decenoic acid; <i>omega</i> -Hydroxy C10:1 (2- <i>trans</i>) fatty acid; 10-HDA	50 mg	250.00
	<p>$C_{10}H_{18}O_3$ Source: synthetic Appearance: solid Storage: -20°C</p>	<p>Mol. Wt.: 186 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol Melting Point (°C): 63-65</p>	<p>CAS#: 14113-05-4 Identity: confirmed by MS</p>
1881	15-Hydroxypentadecanoic acid <i>omega</i> -Hydroxy C15:0 fatty acid	25 mg	125.00
	<p>$C_{15}H_{30}O_3$ Source: synthetic Appearance: solid Storage: room temperature</p>	<p>Mol. Wt.: 258 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol Melting Point (°C): 84-86</p>	<p>CAS#: 4617-33-8</p>
1882	Methyl 15-hydroxypentadecanoate <i>omega</i> -Hydroxy C15:0 methyl ester	25 mg	125.00
	<p>$C_{16}H_{32}O_3$ Source: synthetic Appearance: solid Storage: room temperature</p>	<p>Mol. Wt.: 272 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, ethyl ether Melting Point (°C): 50-52</p>	<p>CAS#: 76529-42-5 Identity: confirmed by MS</p>
1760	17-Hydroxyheptadecanoic acid <i>omega</i> -Hydroxy C17:0 fatty acid	25 mg	250.00
	<p>$C_{17}H_{34}O_3$ Source: synthetic Appearance: solid Storage: room temperature</p>	<p>Mol. Wt.: 286 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, ethyl ether Melting Point (°C): 93-95</p>	<p>CAS#: 13099-34-8 Identity: confirmed by MS</p>
1761	Methyl 17-hydroxyheptadecanoate <i>omega</i> -Hydroxy C17:0 methyl ester	25 mg	250.00
	<p>$C_{18}H_{36}O_3$ Source: synthetic Appearance: solid Storage: room temperature</p>	<p>Mol. Wt.: 300 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, ethyl ether Melting Point (°C): 59-63</p>	<p>CAS#: 94036-00-7</p>
1877	20-Hydroxyeicosanoic acid <i>omega</i> -Hydroxy C20:0 fatty acid	25 mg	250.00
	<p>$C_{20}H_{40}O_3$ Source: synthetic Appearance: solid Storage: room temperature</p>	<p>Mol. Wt.: 328 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol Melting Point (°C): 96-98</p>	<p>CAS#: 62643-46-3</p>
1878	Methyl 20-hydroxyeicosanoate <i>omega</i> -Hydroxy C20:0 methyl ester	25 mg	250.00
	<p>$C_{21}H_{42}O_3$ Source: synthetic Appearance: solid Storage: room temperature</p>	<p>Mol. Wt.: 342 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, ethyl ether Melting Point (°C): 69-71</p>	<p>CAS#: 37477-29-5 Identity: confirmed by MS</p>

1880	Methyl 21-hydroxyheneicosanoate <i>omega</i> -Hydroxy C21:0 methyl ester	25 mg	275.00
	$C_{22}H_{44}O_3$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 356 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, ethyl ether Melting Point (°C): 73-76	CAS#: 94035-98-0
1818	22-Hydroxydocosanoic acid <i>omega</i> -Hydroxy C22:0 fatty acid	25 mg	275.00
	$C_{22}H_{44}O_3$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 356 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol Melting Point (°C): 100-102	Identity: confirmed by MS
1819	Methyl 22-hydroxydocosanoate <i>omega</i> -Hydroxy C22:0 methyl ester	25 mg	275.00
	$C_{23}H_{46}O_3$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 370 Purity: 98+% by TLC, GC Solubility: chloroform, warm ethanol, ethyl ether Melting Point (°C): 73-75	
1883	Methyl 27-hydroxyheptacosanoate <i>omega</i> -Hydroxy C27:0 methyl ester	25 mg	300.00
	$C_{28}H_{56}O_3$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 440 Purity: 97+% by TLC, GC Solubility: chloroform Melting Point (°C): 85-89	Identity: confirmed by MS
1884	Methyl 30-hydroxytriacontanoate <i>omega</i> -Hydroxy C30:0 methyl ester	25 mg	300.00
	$C_{31}H_{62}O_3$ Source: synthetic Appearance: solid Storage: room temperature	Mol. Wt.: 482 Purity: 97+% by TLC, GC Solubility: chloroform Melting Point (°C): 88-91	CAS#: 79162-70-2 Identity: confirmed by MS

Branched and Cyclic Fatty Acids


Iso-Fatty Acids and Methyl Esters



1656	Methyl 11-methyldodecanoate iso-Tridecanoic methyl ester; iso C13 Methyl ester	20 mg	355.00
	$C_{14}H_{28}O_2$ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 228 Purity: 98+% by GC Solubility: hexane, ethyl ether, methylene chloride	CAS#: 5129-57-7
1657	Methyl 12-methyltridecanoate iso-Tetradecanoic methyl ester; iso C14 Methyl ester	20 mg	355.00
	$C_{15}H_{30}O_2$ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 242 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 5129-58-8

1605	13-Methyltetradecanoic acid iso-Pentadecanoic acid; iso C15 Fatty acid	20 mg	355.00
	$C_{15}H_{30}O_2$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 242 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 2485-71-4
1600	Methyl 13-methyltetradecanoate iso-Pentadecanoic methyl ester; iso C15 Methyl ester	20 mg	355.00
	$C_{16}H_{32}O_2$ Source: synthetic Appearance: liquid Storage: $-20^{\circ}C$	Mol. Wt.: 256 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 5129-59-9
1601	Methyl 14-methylpentadecanoate iso-Palmitic methyl ester; iso C16 Methyl ester	20 mg	355.00
	$C_{17}H_{34}O_2$ Source: synthetic Appearance: liquid Storage: $-20^{\circ}C$	Mol. Wt.: 270 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 5129-60-2
1606	15-Methylhexadecanoic acid iso-Heptadecanoic acid; iso C17 Fatty acid	20 mg	355.00
	$C_{17}H_{34}O_2$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 270 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 1603-03-8
1602	Methyl 15-methylhexadecanoate iso-Heptadecanoic methyl ester; iso C17 Methyl ester	20 mg	355.00
	$C_{18}H_{36}O_2$ Source: synthetic Appearance: liquid Storage: $-20^{\circ}C$	Mol. Wt.: 284 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 6929-04-0
1603	Methyl 17-methyloctadecanoate iso-Nonadecanoic methyl ester; iso C19 Methyl ester	20 mg	355.00
	$C_{20}H_{40}O_2$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 313 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 55124-97-5

Anteiso-Fatty Acids and Methyl Esters



1615	12-Methyltetradecanoic acid anteiso-Pentadecanoic acid; anteiso C15 Fatty acid	20 mg	355.00
	$C_{15}H_{30}O_2$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 242 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 5502-94-3

1612	Methyl 12-methyltetradecanoate anteiso-Pentadecanoic methyl ester; anteiso C15 Methyl ester	20 mg	355.00
	$C_{16}H_{32}O_2$ Source: synthetic Appearance: liquid Storage: $-20^{\circ}C$	Mol. Wt.: 256 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 5129-66-8
1613	Methyl 13-methylpentadecanoate anteiso-Palmitic methyl ester; anteiso C16 Methyl ester	20 mg	355.00
	$C_{17}H_{34}O_2$ Source: synthetic Appearance: liquid Storage: $-20^{\circ}C$	Mol. Wt.: 270 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 5487-50-3
1616	14-Methylhexadecanoic acid anteiso-Heptadecanoic acid; anteiso C17 Fatty acid	20 mg	355.00
	$C_{17}H_{34}O_2$ Source: synthetic Appearance: solid Storage: $-20^{\circ}C$	Mol. Wt.: 270 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 5918-29-6
1614	Methyl 14-methylhexadecanoate anteiso-Heptadecanoic methyl ester; anteiso C17 Methyl ester	20 mg	355.00
	$C_{18}H_{36}O_2$ Source: synthetic Appearance: liquid Storage: $-20^{\circ}C$	Mol. Wt.: 284 Purity: 98+% by GC Solubility: chloroform, ethyl ether, ethanol	CAS#: 2490-49-5
Other Branched Methyl Fatty Acids			
1207	D,L-2,6-Dimethylheptanoic acid 2,6-Dimethyl C7:0 fatty acid	50 mg	170.00
	$C_9H_{18}O_2$ Source: synthetic Appearance: liquid Storage: room temperature	Mol. Wt.: 158 Purity: 98+% by TLC, GC Solubility: chloroform	CAS#: 60148-94-9
1792	Methyl 10-methylhexadecanoate 10-Methyl C16:0 methyl ester	25 mg	245.00
	$C_{18}H_{36}O_2$ Source: synthetic Appearance: liquid Storage: room temperature	Mol. Wt.: 284 Purity: 98+% by TLC, GC Solubility: chloroform	CAS#: 2490-51-9
1195	Phytanic acid 3,7,11,15-Tetramethylhexadecanoic acid	25 mg	365.00
	$C_{20}H_{40}O_2$ Source: semisynthetic Appearance: liquid Storage: $-20^{\circ}C$	Mol. Wt.: 312 Purity: 97+% by GC Solubility: chloroform, methanol	CAS#: 14721-66-5

Cyclopropyl Fatty Acids and Methyl Esters


1822	<i>cis</i>-9,10-Methyleneoctadecanoic acid Dihydrosterculic acid	25 mg	250.00
	$C_{19}H_{36}O_2$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 296 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, hexane Melting Point (°C): 38-42	CAS#: 4675-61-0
1823	Methyl <i>cis</i>-9,10-methyleneoctadecanoate Methyl dihydrosterculate	25 mg	250.00
	$C_{20}H_{38}O_2$ Source: synthetic Appearance: liquid Storage: -20°C	Mol. Wt.: 310 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, hexane	CAS#: 3971-54-8 Identity: confirmed by MS
1238 *NEW*	Methyl malvalate Methyl 8,9-methylene-heptadec-8Z-enoate; Methyl 2-octyl-1-cyclopropene-1-heptanoate	5 mg	375.00
	$C_{19}H_{34}O_2$ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 294 Purity: 95+% by TLC, GC Solubility: hexane, ethyl ether, chloroform, methanol	CAS#: 5026-66-4
1235 *NEW*	Sterculic acid 9,10-Methylene-octadec-9-enoic acid; 2-Octyl-1-cyclopropene-1-octanoic acid	25 mg	325.00
	$C_{19}H_{34}O_2$ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 295 Purity: 98+% by TLC, GC Solubility: chloroform, hexane, ethyl ether, methanol	CAS#: 738-87-4 Identity: confirmed by MS
1236 *NEW*	Methyl sterculate Methyl 9,10-methylene-octadec-9-enoate; Methyl 2-octyl-1-cyclopropene-1-octanoate	25 mg	325.00
	$C_{20}H_{36}O_2$ Source: natural, plant Appearance: liquid Storage: -20°C	Mol. Wt.: 309 Purity: 98+% by TLC, GC Solubility: chloroform, hexane, ethyl ether, methanol	CAS#: 3220-60-8 Identity: confirmed by MS

Unusual Fatty Acids and Derivatives

1751	N-Oleoylethanolamine NOE	100 mg	230.00
	$C_{20}H_{39}NO_2$ Source: synthetic Appearance: solid Storage: -20°C Activity: acid ceramidase inhibitor	Mol. Wt.: 326 Purity: 98+% by TLC, GC Solubility: chloroform, ethanol, methanol, ethyl ether, DMSO Melting Point (°C): 63-66	CAS#: 111-58-0 Identity: confirmed by MS

Vitamin E

Tocopherols



1072 ***rac-alpha*-Tocopherol** **50 mg/ml, 1 ml** **145.00**
5,7,8-Trimethyltolcol

$C_{29}H_{50}O_2$

Source: synthetic

Appearance: liquid

Storage: $-20^{\circ}C$

Mol. Wt.: 431


Purity: 95% by TLC, 98% by GC, HPLC

Solubility: chloroform, ethanol, hexane, methanol

Solvent: hexane

CAS#: 10191-41-0

Identity: confirmed by MS



1071 ***rac-beta*-Tocopherol** **50 mg/ml, 1 ml** **210.00**
5,8-Dimethyltolcol

$C_{28}H_{48}O_2$

Source: synthetic

Appearance: liquid

Storage: $-20^{\circ}C$

Mol. Wt.: 417


Purity: 95% by TLC, 98% by GC, HPLC

Solubility: chloroform, ethanol, hexane, methanol

Solvent: hexane

CAS#: 148-03-8

Identity: confirmed by MS



1073 ***rac-gamma*-Tocopherol** **50 mg/ml, 1 ml** **210.00**
7,8-Dimethyltolcol

$C_{28}H_{48}O_2$

Source: synthetic

Appearance: liquid

Storage: $-20^{\circ}C$

Mol. Wt.: 417


Purity: 95% by TLC, 96% by GC, HPLC

Solubility: chloroform, ethanol, hexane, methanol

Solvent: hexane

CAS#: 73980-80-0

Identity: confirmed by MS



1790 ***(+)-delta*-Tocopherol** **50 mg/ml, 1 ml** **190.00**
8-Methyltolcol

$C_{27}H_{46}O_2$

Source: natural, plant

Appearance: liquid

Storage: $-20^{\circ}C$

Mol. Wt.: 403


Purity: 95% by TLC, 98% by GC

Solubility: chloroform, ethanol, hexane, methanol

Solvent: hexane


CAS#: 119-13-1

Identity: confirmed by MS



1074 **rac-5,7-Dimethyltocol** **50 mg/ml, 1 ml** **210.00**


$C_{28}H_{48}O_2$ **Mol. Wt.:** 417 **CAS#:** 493-35-6
Source: synthetic **Purity:** 95% by TLC, 98% by GC, HPLC **Identity:** confirmed by MS
Appearance: liquid **Solubility:** hexane, ethyl ether, chloroform, alcohols
Storage: $-20^{\circ}C$ **Solvent:** hexane



1797 **Tocol** **50 mg/ml, 1 ml** **200.00**
rac-Tocol

$C_{26}H_{44}O_2$ **Mol. Wt.:** 389 **CAS#:** 119-98-2
Source: synthetic **Purity:** 95% by TLC, 98% by GC, HPLC **Identity:** confirmed by MS
Appearance: liquid **Solubility:** hexane, methanol, ethanol
Storage: $-20^{\circ}C$ **Solvent:** hexane


Tocotrienols



2109 ***alpha*-Tocotrienol** **25 mg** **345.00**

3,4-Dihydro-2,5,7,8-tetramethyl-2R-[(3E,7E)-4,8,12-trimethyl-3,7,11-tridecatrienyl]-2H-1-benzopyran-6-ol


$C_{29}H_{44}O_2$ **Mol. Wt.:** 425 **CAS#:** 58864-81-6
Source: natural, plant **Purity:** 98+% by TLC, GC, HPLC **Identity:** confirmed by MS
Appearance: liquid **Solubility:** hexane, ethyl ether, ethanol
Storage: $-20^{\circ}C$



2110 ***beta*-Tocotrienol** **25 mg** **650.00**


[R-(E,E)]-3,4-Dihydro-2,5,8-trimethyl-2-(4,8,12-trimethyl-3,7,11-tridecatrienyl)-2H-1-benzopyran-6-ol

$C_{28}H_{42}O_2$ **Mol. Wt.:** 411 **CAS#:** 490-23-3
Source: semisynthetic, plant **Purity:** 98+% by TLC, GC, HPLC **Identity:** confirmed by MS
Appearance: liquid **Solubility:** chloroform, ethyl ether, hexane
Storage: $-20^{\circ}C$



2111 **gamma-Tocotrienol** **25 mg** **345.00**
 [R-(E,E)]-3,4-Dihydro-2,7,8-trimethyl-2-(4,8,12-trimethyl-3,7,11-tridecatrienyl)-2H-1-benzopyran-6-ol

$C_{28}H_{42}O_2$ **Mol. Wt.:** 411 **CAS#:** 14101-61-2
Source: natural, plant **Purity:** 98+% by TLC, GC, HPLC **Identity:** confirmed by MS
Appearance: liquid **Solubility:** chloroform, ethyl ether, hexane
Storage: -20°C




2112 **delta-Tocotrienol** **25 mg** **345.00**
 [R-(E,E)]-3,4-Dihydro-2,8-dimethyl-2-(4,8,12-trimethyl-3,7,11-tridecatrienyl)-2H-1-benzopyran-6-ol

$C_{27}H_{40}O_2$ **Mol. Wt.:** 397 **CAS#:** 25612-59-3
Source: natural, plant **Purity:** 98+% by TLC, GC, HPLC **Identity:** confirmed by MS
Appearance: liquid **Solubility:** chloroform, ethyl ether, hexane
Storage: -20°C

Sterols

Cholestane Derivatives



1006 **Cholesterol** **500 mg** **130.00**

$C_{27}H_{46}O$ **Mol. Wt.:** 387 **CAS#:** 57-88-5
Source: natural, ovine **Purity:** 98+% by TLC, GC
Appearance: solid **Solubility:** chloroform, ethanol
Storage: -20°C **Melting Point (°C):** 147-148

1115 **5-alpha-Cholestane** **100 mg** **140.00**

$C_{27}H_{48}$ **Mol. Wt.:** 373 **CAS#:** 481-21-0
Source: synthetic **Purity:** 98+% by GC
Appearance: solid **Solubility:** chloroform, ethyl ether, hexane
Storage: -20°C

1116	Coprostanol 5- <i>beta</i> -Cholestan-3- <i>beta</i> -ol	25 mg	240.00
	C₂₇H₄₈O Source: semisynthetic Appearance: solid Storage: -20°C	Mol. Wt.: 389 Purity: 98+% by GC Solubility: chloroform, ethyl ether, warm methanol Melting Point (°C): 101-103	CAS#: 360-68-9

Plant Sterols and Steryl Glucosides

1119	Plant Sterol Mixture Sterol mixture, qualitative	25 mg/ml, 1 ml	130.00
	Source: natural, plant Appearance: liquid Storage: -20°C Contains: Brassicasterol (13%), Campesterol (26%), Stigmasterol (7%), <i>beta</i> -Sitosterol (53%), in order of elution. Percentages are approximate.	Solubility: chloroform Solvent: chloroform	

1123	Plant Sterols Kit	1 kit	585.00
	Source: synthetic or plant Appearance: liquid/solid Storage: -20°C Contains in individual packages:	Solubility: chloroform Solvent: chloroform	
	Steryl Glucosides 25 mg	Lanosterol (55%) 100 mg	
	Esterified Steryl Glucosides 10 mg	Stigmasterol 25 mg	
	Plant Sterol Mixture 25 mg	Ergosterol 25 mg	
	<i>beta</i> -Sitosterol (55%) 100 mg	Coprostanol 5 mg	
	Desmosterol (85%) 2 mg	Cholestanol 100 mg	

1113	<i>beta</i>-Sitostanol Stigmastanol	50 mg	165.00
	C₂₉H₅₂O Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 417 Purity: 98+% by TLC, 97+% by GC Solubility: chloroform Melting Point (°C): 127-132	CAS#: 83-45-4

1120	Lanosterol	500 mg	105.00
	C₃₀H₅₀O Source: synthetic or plant Appearance: solid Storage: -20°C	Mol. Wt.: 427 Purity: 55% by TLC, GC Solubility: chloroform	CAS#: 79-63-0

1121	Stigmasterol 5,22-Cholestadien-24- <i>beta</i> -ethyl-3- <i>beta</i> -ol	100 mg	140.00
	C₂₉H₄₈O Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 413 Purity: 95% by TLC, GC Solubility: chloroform Melting Point (°C): 165-167	CAS#: 83-48-7

1122	Ergosterol	100 mg	140.00
	C₂₈H₄₄O Source: synthetic or plant Appearance: solid Storage: -20°C	Mol. Wt.: 397 Purity: 95% by TLC, GC Solubility: chloroform Melting Point (°C): 156-158	CAS#: 57-87-4

1117	Steryl Glucosides Sterolins	25 mg	145.00
	<p>$C_{35}H_{60}O_6$ Source: natural, plant Appearance: solid Storage: $-20^{\circ}C$ Sitosteryl (55.9%), Campesteryl (24.6%), Stigmasteryl (18.1%), <i>delta</i>-5-Avenasteryl (1.4%) Percentages are approximate. K. Phillips, J. of Food and Lipids, Vol. 12 pp.124-140 (2005)</p>	<p>Mol. Wt.: 577 (based on <i>beta</i>-sitosteryl glucoside) Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1 (warm) Melting Point ($^{\circ}C$): 283-287</p>	<p>Identity: confirmed by MS</p>

1118	Esterified Steryl Glucosides Esterified sterolins	10 mg	155.00
	<p>$C_{51}H_{90}O_7$ Source: natural, plant Appearance: solid Storage: $-20^{\circ}C$ Sterol, glucose and fatty acid in a molar ratio 1:1:1. Sitosterol (major), Campesterol, Stigmasteryl See Table III (pg. 99) for typical fatty acid content of products prepared from natural sources.</p>	<p>815 (based on <i>beta</i>-sitosteryl glucoside palmitate) Purity: 98+% by TLC Solubility: chloroform, ethyl ether, pyridine</p>	<p>Identity: confirmed by MS</p>

Standards and Reference Compounds

Food Industry Mixtures

Each methyl ester mixture is carefully prepared by weight.

4210	KEL-FIM-FAME-5 Mixture Methyl ester mixture	15.5 mg/ml, 1 ml	120.00
	<p>Source: synthetic or plant Appearance: liquid Storage: $-20^{\circ}C$</p>	<p>Solubility: heptane Solvent: heptane</p>	<p>Contains the methyl esters of the following fatty acids (mg/ml in brackets): C8:0 [0.3], C10:0 [0.5], C12:0 [1.0], C13:0 [0.5], C14:0 [0.5], C14:1 [0.3], C15:0 [0.3], C16:0 [2.0], C16:1 [1.0], C17:0 [0.5], C18:0 [1.0], C18:1tr [0.4], C18:1c [3.0], C18:2 [2.0], C20:0 [0.3], C18:3 [1.0], C20:1 [0.3], C22:0 [0.3], C22:1 [0.3] Listed in order of their elution using a SP-2330 30m x 0.25mm x 0.2μm capillary column.</p>
2009	FIM-FAME-6 Mixture Methyl ester mixture	33 mg/ml, 1 ml	145.00
	<p>Source: synthetic or plant Appearance: liquid Storage: $-20^{\circ}C$</p>	<p>Solvent: heptane</p>	<p>Contains the methyl esters of these fatty acids. Each methyl ester is 3.03% of the mixture except C16:0 which is 6.06%. C4:0, C6:0, C8:0, C10:0, C11:0, C12:0, C13:0, C14:0, C14:1(<i>cis</i>-9), C15:0, C15:1(<i>cis</i>-10), C16:0, C16:1(<i>cis</i>-9), C17:0, C17:1(<i>cis</i>-10), C18:0, C18:1(<i>trans</i>-9), C18:1(<i>cis</i>-9), C18:2(all <i>cis</i>-9,12), C20:0, C18:3(all <i>cis</i>-6,9,12), C20:1(<i>cis</i>-11), C18:3(all <i>cis</i>-9,12,15), C20:2(all <i>cis</i>-11,14), C22:0, C20:3(all <i>cis</i>-8,11,14), C22:1(<i>cis</i>-13), C20:3(all <i>cis</i>-11,14,17), C20:4(all <i>cis</i>-5,8,11,14), C22:2(all <i>cis</i>-13,16), C24:1(<i>cis</i>-15), C22:6(all <i>cis</i>-4,7,10,13,16,19) Listed in order of their elution using a SP-2560 100m x 0.25mm x 0.2μm capillary column.</p>

2010	FIM-FAME-7 Mixture Methyl ester mixture	30 mg/ml, 1 ml	145.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solubility: methylene chloride Solvent: methylene chloride	
	Contains the methyl esters of these fatty acids. (weight percent in [brackets]): C4:0 [4.0], C6:0 [4.0], C8:0 [4.0], C10:0 [4.0], C11:0 [2.0], C12:0 [4.0], C13:0 [2.0], C14:0 [4.0], C14:1(<i>cis</i> -9) [2.0], C15:0 [2.0], C15:1(<i>cis</i> -10) [2.0], C16:0 [6.0], C16:1(<i>cis</i> -9) [2.0], C17:0 [2.0], C17:1(<i>cis</i> -10) [2.0], C18:0 [4.0], C18:1(<i>trans</i> -9) [2.0], C18:1(<i>cis</i> -9) [4.0], C18:2(all <i>trans</i> -9,12) [2.0], C18:2(all <i>cis</i> -9,12) [2.0], C20:0 [4.0], C18:3(all <i>cis</i> -6,9,12) [2.0], C20:1(<i>cis</i> -11) [2.0], C18:3(all <i>cis</i> -9,12,15) [2.0], C21:0 [2.0], C20:2(all <i>cis</i> -11,14) [2.0], C22:0 [4.0], C20:3 (all <i>cis</i> -8,11,14) [2.0], C22:1(<i>cis</i> -13) [2.0], C20:3(all <i>cis</i> -11,14,17) [2.0], C20:4(all <i>cis</i> -5,8,11,14) [2.0], C23:0 [2.0], C22:2(all <i>cis</i> -13,16) [2.0], C24:0 [4.0], C20:5(all <i>cis</i> -5,8,11,14,17) [2.0], C24:1(<i>cis</i> -15) [2.0], C22:6(all <i>cis</i> -4,7,10,13,16,19) [2.0] Listed in order of their elution using a SP-2560 100m x 0.25mm x 0.2µm capillary column.		

2012	FIM-FAME-8 Mixture C18 Quantitative mixture	25 mg/ml, 1 ml	115.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solvent: methylene chloride	
	Contains the methyl esters of these fatty acids (weight percent in [brackets]): C18:0 [20.0], C18:1 [20.0], C18:2 [20.0], C18:3 [20.0], C18:4 [20.0] Listed in order of their elution using a SP-2330 30m x 0.25mm x 0.2µm capillary column.		

2013	FIM-FAME-9 Mixture C20 Quantitative mixture	25 mg/ml, 1 ml	115.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solvent: methylene chloride	
	Contains the methyl esters of these fatty acids (weight percent in [brackets]): C20:0 [14.3], C20:1 [14.3], C20:2 [14.3], C20:3 [14.3], C20:4 [14.3], C20:5 [14.3], C22:6 [14.3] Listed in order of their elution using a SP-2330 100m x 0.25mm x 0.2µm capillary column.		

Polyunsaturated Fatty Acid Methyl Ester Mixtures

These are complex qualitative standard mixtures of polyunsaturated fatty acid methyl esters. Because they are extracted from natural materials, relative peak sizes may vary from lot to lot.

1093	PUFA-1 Qualitative mixture	100 mg	155.00
	Source: natural, fish oil Appearance: liquid Storage: -20°C	Solubility: chloroform, ethanol, hexane, methanol	
	Contains: C14:0, C16:0, C16:1ω7, C18:1ω9, C18:1ω7, C18:2ω6, C20:1ω9, C18:4ω3, C22:1ω11, C22:1ω9, C20:5ω3, C22:5ω3, C22:6ω3		
1081	PUFA-2 Qualitative mixture	100 mg	165.00
	Source: natural, porcine Appearance: liquid Storage: -20°C	Solubility: alcohols, hexane, chloroform	
	Contains: C14:0, C16:0, C16:1ω7, C18:0, C18:1ω9, C18:1ω7, C18:2ω6, C18:3ω6, C18:3ω3, C20:1ω9, C20:2ω6, C20:3ω6, C20:4ω6, C20:5ω3, C22:4ω6, C22:5ω3, C22:6ω3		

1177	PUFA-3 Qualitative mixture	100 mg	165.00
	Source: natural, menhaden oil Appearance: liquid Solubility: alcohols, hexane, chloroform Storage: -20°C Contains: C14:0, C16:0, C16:1 ω 7, C16:2 ω 4, C16:3 ω 4, C16:4 ω 1, C18:0, C18:1 ω 9, C18:1 ω 7, C18:2 ω 6, C18:2 ω 4, C18:3 ω 4, C18:3 ω 3, C18:4 ω 3, C20:1 ω 9, C20:4 ω 6, C20:4 ω 3, C20:5 ω 3, C21:5 ω 3, C22:5 ω 3, C22:6 ω 3		

Carbohydrate Mixtures

1124	Alditol Acetate Mixture-1 Quantitative carbohydrate mixture	50 mg/ml, 1 ml	220.00
	Source: synthetic Appearance: liquid Solubility: chloroform Storage: -20°C Solvent: chloroform Contains: rhamnitol, fucitol, ribitol and arabinitol pentaacetates, 12.5 mg/ml each		

1125	Alditol Acetate Mixture-2 Quantitative carbohydrate mixture	50 mg/ml, 1 ml	220.00
	Source: synthetic Appearance: liquid Solubility: chloroform Storage: -20°C Solvent: chloroform Contains: mannitol, galactitol, glucitol and inositol hexaacetates, 12.5 mg/ml each		

Other Fatty Acid Methyl Ester Mixtures

1722	2-Hydroxy Methyl Ester Mixture Quantitative mixture	10 mg/ml, 1 ml	180.00
	Source: synthetic Appearance: liquid Solubility: chloroform Storage: -20°C Solvent: chloroform Contains: 2-OH C14:0, 20.0%; 2-OH C16:0, 20.0%; 2-OH C18:0, 15.0%; 2-OH C20:0, 15.0%; 2-OH C22:0, 10.0%; 2-OH C23:0, 10.0%; 2-OH C24:0, 10.0%		

1131	Cis-Trans FAME Isomer Standard Mixture	5.5 mg/ml, 5 ml	170.00
	Source: margarine Appearance: liquid Solubility: methylene chloride, chloroform Storage: -20°C Solvent: methylene chloride <p>Analysis of positional <i>cis-trans</i> fatty acid isomers is ever more important in light of the new food industry rules. These isomers can be resolved on Supelco SP-2560 or an equivalent capillary GC column. Use this specially formulated mixture to ensure proper operation of your column for this tricky separation. Mixture consists of <i>cis-trans</i> fatty acid isomers as methyl esters in methylene chloride.</p> <p>This is a qualitative mixture containing: C16:0, C18:0, C18:1 <i>trans</i> isomers (4 peaks), C18:1 <i>cis</i> & <i>trans</i> isomers (2 peaks), C18:1 <i>cis</i> isomers (4 peaks), C18:2, C20:0, C20:1 and C18:3 (same peak), C22:0 Listed in order of their elution using a SP-2560 100m x 0.25mm x 0.2μ capillary column.</p>		

2011	Long Chain Fatty Acid Methyl Ester Mixture C24:0, C26:0, C28:0, C30:0, C32:0 Fatty acid methyl ester mixture	25 mg/ml, 1 ml	140.00
	Source: synthetic Appearance: liquid Solubility: methylene chloride Storage: -20°C Solvent: methylene chloride Quantitative mixture contains: C24:0, 20.0%; C26:0, 20.0%; C28:0, 20.0%; C30:0, 20.0%; C32:0, 20.0%		

AOCS Animal and Vegetable Oil Reference Mixtures (RM Mixtures)

By studying problems with the quantitative analysis of animal and vegetable oils and fats, the American Oil Chemists' Society has found certain mixtures to be useful as reference standards. The composition of each mixture (see Table I below) is similar to the fatty acid distribution of certain oils. All mixtures are in methyl ester form and ready for GC analysis

Table I. AOCS Oil Reference Mixtures

Each methyl ester mixture is carefully prepared by weight and the composition verified by gas chromatography. The weight percentage of each component is indicated in the Table.

Mix No. Catalog No.	RM-1 1084	RM-2 1085	RM-3 1086	Rapeseed 1083	RM-4 1087	RM-5 1088	RM-6 1089
C8:0 Caprylate						7.0	
C10:0 Caprate						5.0	
C12:0 Laurate						48.0	
C14:0 Myristate			1.0	1.0		15.0	2.0
C16:0 Palmitate	6.0	7.0	4.0	4.0	11.0	7.0	30.0
C16:1 Palmitoleate (<i>cis</i> -9)							3.0
C18:0 Stearate	3.0	5.0	3.0	3.0	3.0	3.0	14.0
C18:1 Oleate (<i>cis</i> -9)	35.0	18.0	45.0	60.0	80.0	12.0	41.0
C18:2 Linoleate (all <i>cis</i> -9,12)	50.0	36.0	15.0	12.0	6.0	3.0	7.0
C18:3 Linolenate (all <i>cis</i> -9,12,15)	3.0	34.0	3.0	5.0			3.0
C20:0 Arachidate	3.0		3.0	3.0			
C20:1 Eicosenoate (<i>cis</i> -11)				1.0			
C22:0 Behenate			3.0	3.0			
C22:1 Erucate (<i>cis</i> -13)			20.0	5.0			
C24:0 Lignocerate			3.0	3.0			

1083 Rapeseed Oil Reference Mixture **25 mg/ml, 1 ml** **85.00**

Source: synthetic or plant
 Appearance: liquid Solubility: ethyl ether, methylene chloride
 Storage: -20°C Solvent: methylene chloride
 Suitable standard for low erucic acid oil

1084 RM-1 Mixture **50 mg** **85.00**

Source: synthetic or plant
 Appearance: liquid Solubility: chloroform, ethyl ether
 Storage: -20°C
 Suitable standard for corn, cottonseed, soybean, safflower, sesame, poppy seed, walnut kapok, and rice oils


1085 RM-2 Mixture **50 mg** **85.00**

Source: synthetic or plant
 Appearance: liquid Solubility: chloroform, ethanol, ethyl ether
 Storage: -20°C
 Suitable standard for linseed, perilla, hempseed, and rubberseed oils

1086	RM-3 Mixture	50 mg/ml, 1 ml	85.00
	Source: synthetic or plant Appearance: liquid Solubility: ethyl ether, methylene chloride Storage: -20°C Solvent: methylene chloride Suitable standards for peanut, rapeseed, and mustard seed oils		
1087	RM-4 Mixture	50 mg	85.00
	Source: synthetic or plant Appearance: liquid Solubility: chloroform, ethyl ether Storage: -20°C Suitable standard for olive, teaseed, and neatsfoot oils		
1088	RM-5 Mixture	50 mg	85.00
	Source: synthetic or plant Appearance: liquid Solubility: chloroform Storage: -20°C Suitable standard for coconut, palm kernel, babassu and ouri-ouri oils		
1089	RM-6 Mixture	50 mg	85.00
	Source: synthetic or plant Appearance: liquid Solubility: ethyl ether, methylene chloride Storage: -20°C Suitable standard for lard, beef tallow, mutton tallow, and palm oil		

Custom Mixtures

Custom fatty acid methyl ester mixtures can be prepared to your specification. Minimum quantity requirements apply to these orders.



Cat# 4210 spiked with 0.4 mg/ml C18:2t ester (methyl linoelaidate) and chromatographed on a Supelco SP 2330 fused silica column.

Peak number	FAME
1	C8:0
2	C10:0
3	C12:0
4	C13:0
5	C14:0
6	C14:1
7	C15:0
8	C16:0
9	C16:1
10	C17:0
11	C18:0
12	C18:1t-9
13	C18:1c-9
14	C18:2t,t-9,12
15	C18:2c,c-9,12
16	C20:0
17	C18:3
18	C20:1
19	C22:0
20	C22:1

Table II. Standards for GC Analysis**GLC Standard Mixtures**

GLC-10 through GLC-100 standards are **equal weight measures** of fatty acid methyl esters. They are **quantitative** standards, useful for determining relative **retention times and response factors**.

Each methyl ester mixture is carefully prepared by weight and the composition verified by gas chromatography. The weight percentage of each component is indicated below. All double bonds are *cis*.

Mixture Number Catalog Number	GLC-10 1095	GLC-30 1097	GLC-40 1098	GLC-50 1099	GLC-60 1100	GLC-70 1101	GLC-80 1102	GLC-90 1103	GLC-100 1104
C8:0 Methyl octanoate (caprylate)		20.0				20.0			
C9:0 Methyl nonanoate (pelargonate)						20.0			
C10:0 Methyl decanoate (caprate)		20.0				20.0			
C11:0 Methyl undecanoate (hendecanoate)						20.0			
C12:0 Methyl dodecanoate (laurate)		20.0				20.0			
C13:0 Methyl tridecanoate							20.0	20.0	
C14:0 Methyl tetradecanoate (myristate)		20.0					20.0		
C15:0 Methyl pentadecanoate							20.0	20.0	
C16:0 Methyl hexadecanoate (palmitate)	20.0	20.0	25.0				20.0		
C16:1 Methyl hexadecenoate (<i>cis</i> -9) (palmitoleate)				20.0					
C17:0 Methyl heptadecanoate (margarate)							20.0	20.0	
C18:0 Methyl octadecanoate (stearate)	20.0		25.0						20.0
C18:1 Methyl octadecenoate (<i>cis</i> -9) (oleate)	20.0			20.0					
C18:2 Methyl octadecadienoate (all <i>cis</i> -9,12) (linoleate)	20.0								
C18:3 Methyl octadecatrienoate (all <i>cis</i> -9,12,15) (linolenate)	20.0								
C19:0 Methyl nonadecanoate								20.0	20.0
C20:0 Methyl eicosanoate (arachidate)			25.0		25.0				20.0
C20:1 Methyl eicosenoate (<i>cis</i> -11)				20.0	25.0				
C20:2 Methyl eicosadienoate (all <i>cis</i> -11,14)					25.0				
C20:3 Methyl eicosatrienoate (all <i>cis</i> -11,14,17)					25.0				
C21:0 Methyl heneicosanoate								20.0	20.0
C22:0 Methyl docosanoate (behenate)			25.0						20.0
C22:1 Methyl docosenoate (<i>cis</i> -13) (erucate)				20.0					
C24:1 Methyl tetracosenoate (<i>cis</i> -15) (nervonate)				20.0					

1095 GLC-10 Mixture **50 mg** **85.00**

Source: synthetic or plant
Appearance: liquid **Solubility:** methylene chloride
Storage: -20°C

1097 GLC-30 Mixture **50 mg** **85.00**

Source: synthetic or plant
Appearance: liquid **Solubility:** methylene chloride
Storage: -20°C

1098 GLC-40 Mixture **50 mg/ml, 1 ml** **85.00**

Source: synthetic or plant
Appearance: liquid **Solubility:** methylene chloride
Storage: -20°C **Solvent:** methylene chloride

1099	GLC-50 Mixture	50 mg/ml, 1 ml	85.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solubility: methylene chloride Solvent: methylene chloride	
1100	GLC-60 Mixture	50 mg/ml, 1 ml	85.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solubility: methylene chloride Solvent: methylene chloride	
1101	GLC-70 Mixture	50 mg	85.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solubility: methylene chloride	
1102	GLC-80 Mixture	50 mg/ml, 1 ml	85.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solubility: methylene chloride Solvent: methylene chloride	
1103	GLC-90 Mixture	50 mg/ml, 1 ml	85.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solubility: methylene chloride Solvent: methylene chloride	
1104	GLC-100 Mixture	50 mg/ml, 1 ml	175.00
	Source: synthetic or plant Appearance: liquid Storage: -20°C	Solubility: methylene chloride Solvent: methylene chloride	

Water Soluble Fatty Acid Mixtures

1106	WSFA-2 Mixture	5 ml	165.00
	Water soluble fatty acid qualitative mixture		
	Appearance: liquid Storage: room temperature	Solubility: DI water Solvent: DI water	
	Contains: acetic, propionic, isobutyric, n-butyric, isovaleric and n-valeric acids 1mg/ml each		
1108	WSFA-4 Mixture	5 ml	165.00
	Water soluble fatty acid qualitative mixture		
	Appearance: liquid Storage: room temperature	Solubility: DI water Solvent: DI water	
	Contains: acetic, propionic, isobutyric, n-butyric, 2-methylbutyric, isovaleric and n-valeric acids 1mg/ml each		

Microbiology Standard Mixtures

1105	GLC-110 Mixture Bacterial lipid standard, qualitative mixture	10 mg/ml, 1 ml	175.00																																																				
	<p>Source: various Appearance: liquid Storage: -20°C</p> <p>Solubility: methylene chloride, chloroform Solvent: chloroform</p> <p>Contains:</p> <table border="0"> <tr> <td>Methyl 12-methyltridecanoate</td> <td>(iso-C14:0)</td> <td>Methyl 14-methylpentadecanoate</td> <td>(iso-C16:0)</td> </tr> <tr> <td>Methyl tetradecanoate (myristate)</td> <td>(C14:0)</td> <td>Methyl hexadecanoate (palmitate)</td> <td>(C16:0)</td> </tr> <tr> <td>Methyl 12-methyltetradecanoate</td> <td>(anteiso-C15:0)</td> <td>Methyl 14-methylhexadecanoate</td> <td>(anteiso-C17:0)</td> </tr> <tr> <td>Methyl pentadecanoate</td> <td>(C15:0)</td> <td></td> <td></td> </tr> </table>	Methyl 12-methyltridecanoate	(iso-C14:0)	Methyl 14-methylpentadecanoate	(iso-C16:0)	Methyl tetradecanoate (myristate)	(C14:0)	Methyl hexadecanoate (palmitate)	(C16:0)	Methyl 12-methyltetradecanoate	(anteiso-C15:0)	Methyl 14-methylhexadecanoate	(anteiso-C17:0)	Methyl pentadecanoate	(C15:0)																																								
Methyl 12-methyltridecanoate	(iso-C14:0)	Methyl 14-methylpentadecanoate	(iso-C16:0)																																																				
Methyl tetradecanoate (myristate)	(C14:0)	Methyl hexadecanoate (palmitate)	(C16:0)																																																				
Methyl 12-methyltetradecanoate	(anteiso-C15:0)	Methyl 14-methylhexadecanoate	(anteiso-C17:0)																																																				
Methyl pentadecanoate	(C15:0)																																																						
1114	Bacterial Acid Methyl Esters CP Mixture Quantitative mixture	10 mg/ml, 1 ml	225.00																																																				
	<p>Source: various Appearance: liquid Storage: -20°C</p> <p>Solubility: hexane, ethanol, methanol Solvent: methyl caproate</p> <p>A quantitative standard. Mixture consists of equal amounts of the compounds listed.</p> <table border="0"> <tr> <td>Methyl undecanoate</td> <td>C11:0</td> <td>Methyl hexadecanoate (<i>cis</i>-9), (palmitoleate)</td> <td>C16:1(<i>cis</i>-9)</td> </tr> <tr> <td>Methyl 2-hydroxydecanoate</td> <td>2-OH C10:0</td> <td>Methyl hexadecanoate, (palmitate)</td> <td>C16:0</td> </tr> <tr> <td>Methyl dodecanoate, (laurate)</td> <td>C12:0</td> <td>Methyl 15-methylhexadecanoate</td> <td>iso-C17:0</td> </tr> <tr> <td>Methyl tridecanoate</td> <td>C13:0</td> <td>Methyl <i>cis</i>-9,10-methylenehexadecanoate</td> <td>C17:0 (all <i>cis</i>-9,10)</td> </tr> <tr> <td>Methyl 2-hydroxydodecanoate</td> <td>2-OH C12:0</td> <td>Methyl heptadecanoate, (margarate)</td> <td>C17:0</td> </tr> <tr> <td>Methyl 3-hydroxydodecanoate</td> <td>3-OH C12:0</td> <td>Methyl 2-hydroxyhexadecanoate</td> <td>2-OH C16:0</td> </tr> <tr> <td>Methyl tetradecanoate, (myristate)</td> <td>C14:0</td> <td>Methyl octadecadienoate (all <i>cis</i>-9,12), (linoleate)</td> <td>C18:2 (all <i>cis</i>-9,12)</td> </tr> <tr> <td>Methyl 13-methyltetradecanoate</td> <td>iso-C15:0</td> <td>Methyl octadecenoate (<i>cis</i>-9), (oleate)</td> <td>C18:1(<i>cis</i>-9)</td> </tr> <tr> <td>Methyl 12-methyltetradecanoate</td> <td>anteiso-C15:0</td> <td>Methyl octadecenoate, (<i>trans</i>-9), (elaidate)</td> <td>C18:1 (<i>trans</i>-9)</td> </tr> <tr> <td>Methyl pentadecanoate</td> <td>C15:0</td> <td>Methyl octadecanoate, (stearate)</td> <td>C18:0</td> </tr> <tr> <td>Methyl 2-hydroxytetradecanoate</td> <td>2-OH C14:0</td> <td>Methyl <i>cis</i>-9,10-methyleneoctadecanoate</td> <td>C19:0 (all <i>cis</i>-9,10)</td> </tr> <tr> <td>Methyl 3-hydroxytetradecanoate</td> <td>3-OH C14:0</td> <td>Methyl nonadecanoate</td> <td>C19:0</td> </tr> <tr> <td>Methyl 14-methylpentadecanoate</td> <td>iso-C16:0</td> <td>Methyl eicosanoate, (arachidate)</td> <td>C20:0</td> </tr> </table>	Methyl undecanoate	C11:0	Methyl hexadecanoate (<i>cis</i> -9), (palmitoleate)	C16:1(<i>cis</i> -9)	Methyl 2-hydroxydecanoate	2-OH C10:0	Methyl hexadecanoate, (palmitate)	C16:0	Methyl dodecanoate, (laurate)	C12:0	Methyl 15-methylhexadecanoate	iso-C17:0	Methyl tridecanoate	C13:0	Methyl <i>cis</i> -9,10-methylenehexadecanoate	C17:0 (all <i>cis</i> -9,10)	Methyl 2-hydroxydodecanoate	2-OH C12:0	Methyl heptadecanoate, (margarate)	C17:0	Methyl 3-hydroxydodecanoate	3-OH C12:0	Methyl 2-hydroxyhexadecanoate	2-OH C16:0	Methyl tetradecanoate, (myristate)	C14:0	Methyl octadecadienoate (all <i>cis</i> -9,12), (linoleate)	C18:2 (all <i>cis</i> -9,12)	Methyl 13-methyltetradecanoate	iso-C15:0	Methyl octadecenoate (<i>cis</i> -9), (oleate)	C18:1(<i>cis</i> -9)	Methyl 12-methyltetradecanoate	anteiso-C15:0	Methyl octadecenoate, (<i>trans</i> -9), (elaidate)	C18:1 (<i>trans</i> -9)	Methyl pentadecanoate	C15:0	Methyl octadecanoate, (stearate)	C18:0	Methyl 2-hydroxytetradecanoate	2-OH C14:0	Methyl <i>cis</i> -9,10-methyleneoctadecanoate	C19:0 (all <i>cis</i> -9,10)	Methyl 3-hydroxytetradecanoate	3-OH C14:0	Methyl nonadecanoate	C19:0	Methyl 14-methylpentadecanoate	iso-C16:0	Methyl eicosanoate, (arachidate)	C20:0		
Methyl undecanoate	C11:0	Methyl hexadecanoate (<i>cis</i> -9), (palmitoleate)	C16:1(<i>cis</i> -9)																																																				
Methyl 2-hydroxydecanoate	2-OH C10:0	Methyl hexadecanoate, (palmitate)	C16:0																																																				
Methyl dodecanoate, (laurate)	C12:0	Methyl 15-methylhexadecanoate	iso-C17:0																																																				
Methyl tridecanoate	C13:0	Methyl <i>cis</i> -9,10-methylenehexadecanoate	C17:0 (all <i>cis</i> -9,10)																																																				
Methyl 2-hydroxydodecanoate	2-OH C12:0	Methyl heptadecanoate, (margarate)	C17:0																																																				
Methyl 3-hydroxydodecanoate	3-OH C12:0	Methyl 2-hydroxyhexadecanoate	2-OH C16:0																																																				
Methyl tetradecanoate, (myristate)	C14:0	Methyl octadecadienoate (all <i>cis</i> -9,12), (linoleate)	C18:2 (all <i>cis</i> -9,12)																																																				
Methyl 13-methyltetradecanoate	iso-C15:0	Methyl octadecenoate (<i>cis</i> -9), (oleate)	C18:1(<i>cis</i> -9)																																																				
Methyl 12-methyltetradecanoate	anteiso-C15:0	Methyl octadecenoate, (<i>trans</i> -9), (elaidate)	C18:1 (<i>trans</i> -9)																																																				
Methyl pentadecanoate	C15:0	Methyl octadecanoate, (stearate)	C18:0																																																				
Methyl 2-hydroxytetradecanoate	2-OH C14:0	Methyl <i>cis</i> -9,10-methyleneoctadecanoate	C19:0 (all <i>cis</i> -9,10)																																																				
Methyl 3-hydroxytetradecanoate	3-OH C14:0	Methyl nonadecanoate	C19:0																																																				
Methyl 14-methylpentadecanoate	iso-C16:0	Methyl eicosanoate, (arachidate)	C20:0																																																				
1075	Volatile Acid Mixture Qualitative mixture	100 ml	185.00																																																				
	<p>Appearance: liquid Storage: 4-8°C</p> <p>Solubility: DI water Solvent: DI water</p> <p>Contains: formic, acetic, propionic, isobutyric, n-butyric, isovaleric, n-valeric, isocaproic, n-caproic, and heptanoic acids</p>																																																						
1077	Non-Volatile Acid Mixture Qualitative mixture	100 ml	185.00																																																				
	<p>Appearance: liquid Storage: 4-8°C</p> <p>Solubility: DI water Solvent: DI water</p> <p>Contains: pyruvic, lactic, oxalacetic, oxalic, methylmalonic, malonic, fumaric and succinic acids.</p>																																																						

Biochemical Research Standard Mixtures

These mixtures are prepared by precise gravimetric technique. All mixtures contain equal amounts of listed components. A data sheet is supplied with each mixture.

1127	Polar Lipid Mixture TLC standards mixture	25 mg/ml, 1 ml	150.00
	<p>Source: natural, egg, ovine Appearance: liquid Storage: -20°C</p> <p>Solubility: chloroform/methanol, 2:1 Solvent: chloroform/methanol, 2:1</p> <p>Contains: cholesterol, phosphatidylethanolamine, lecithin, and lyso-lecithin.</p>		

1128	Sphingolipid Mixture TLC standards mixture	25 mg/ml, 1 ml	150.00
	Source: natural, bovine Appearance: liquid Storage: -20°C Contains: cerebrosides, sulfatides, and sphingomyelin.	Solubility: chloroform/methanol, 2:1 Solvent: chloroform/methanol, 2:1	
1129	Non-Polar Lipid Mixture A TLC standards mixture	25 mg/ml, 1 ml	135.00
	Source: natural, plant, ovine Appearance: liquid Storage: -20°C Contains: cholesteryl palmitate, tripalmitin, palmitic acid, and cholesterol.	Solubility: chloroform Solvent: chloroform	
1130	Non-Polar Lipid Mixture B TLC standards mixture	25 mg/ml, 1 ml	135.00
	Source: natural, plant, ovine Appearance: liquid Storage: -20°C Contains: cholesteryl oleate, methyl oleate, triolein, oleic acid, and cholesterol.	Solubility: chloroform Solvent: chloroform	

Glycosphingolipid Reference Mixtures for TLC

These mixtures are qualitative standards prepared from our purified glycosphingolipids.

1505	Neutral Glycosphingolipid Mixture Glycosylceramides, qualitative mixture	1 mg/ml, 1 ml	210.00
	Source: natural, bovine and porcine Appearance: liquid Storage: -20°C Contains: cerebrosides, lactosylceramides, ceramide trihexosides, globosides.	Solubility: chloroform/methanol, 2:1 Solvent: chloroform/methanol, 2:1	
1508	Monosialoganglioside Mixture	0.5 mg/ml, 1 ml	210.00
	Source: natural, bovine, human Appearance: liquid Storage: -20°C Contains: GM ₃ , GM ₂ , GM ₁	Solubility: chloroform/methanol/DI water, 2:1:0.1 Solvent: chloroform/methanol/DI water, 2:1:0.1	
1509	Disialoganglioside Mixture	0.5 mg/ml, 1 ml	210.00
	Source: natural, bovine Appearance: liquid Storage: -20°C Contains: GD ₃ , GD _{1a} , GD _{1b}	Solubility: chloroform/methanol/DI water, 2:1:0.1 Solvent: chloroform/methanol/DI water, 2:1:0.1	
1510	Lactosylceramide and Sialosyl Derivatives Mixture	0.5 mg/ml, 1 ml	225.00
	Source: natural, bovine Appearance: liquid Storage: -20°C Contains: LC, GM ₃ , GD ₃	Solubility: chloroform/methanol/DI water, 2:1:0.1 Solvent: chloroform/methanol/DI water, 2:1:0.1	
1511	Gangliotetraosylceramide and Sialosyl Derivatives Mixture	0.5 mg/ml, 1 ml	225.00
	Source: natural, bovine Appearance: liquid Storage: -20°C Contains: asialo-GM ₁ , GM ₁ , GD _{1a} , GD _{1b} , GT _{1b}	Solubility: chloroform/methanol/DI water, 2:1:0.1 Solvent: chloroform/methanol/DI water, 2:1:0.1	

Labeled Standards

Stable Isotope Labeled Standards

2079	D-erythro-Sphingosine, D9 15,15,16,16,17,17,18,18,18-D9-2-Amino-octadec-4-ene-1,3-diol	1 mg	385.00
	$C_{18}H_{28}D_9NO_2$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 309 Purity: 98+% by TLC, GC, HPLC Solubility: chloroform, ethanol, methanol, DMSO	Identity: confirmed by MS
2201	N-omega-CD₃-Octadecanoyl-D-erythro-sphingosine N-C18:0-CD ₃ -D-erythro-Ceramide; N-Stearoyl-CD ₃ -D-erythro-sphingosine	1 mg	350.00
	$C_{36}H_{68}NO_3D_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 569 Purity: 98+% by TLC, GC Solubility: chloroform, hot ethanol, DMF	Identity: confirmed by MS
2208 *NEW*	N-(32-Linoleoyloxy-dotriacontanoyl)-sphingosine-D9 EOS Ceramide, deuterated; O-acylceramide, deuterated	1 mg	550.00
	$C_{68}H_{120}D_9NO_5$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 1050 Purity: 98+% by TLC Solubility: chloroform, methanol, DMF	Identity: confirmed by MS
2202	N-omega-CD₃-Octadecanoyl-D-erythro-dihydrosphingosine N-C18:0-CD ₃ -D-erythro-Dihydroceramide; N-Stearoyl-CD ₃ -D-erythro-sphinganine	1 mg	325.00
	$C_{36}H_{70}D_3NO_3$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 571 Purity: 98% by TLC, GC, HPLC Solubility: hot ethanol, DMF, DMSO, chloroform/methanol, 2:1	Identity: confirmed by MS
2210 *NEW*	N-omega-CD₃-Octadecanoyl-phytosphingosine N-C18:0-CD ₃ -Phytoceramide; N-Stearoyl-CD ₃ -phytosphingosine	1 mg	275.00
	$C_{36}H_{70}D_3NO_4$ Source: semisynthetic, yeast (<i>Pichia ciferri</i>) Appearance: solid Storage: -20°C	Mol. Wt.: 587 Purity: 98+% by TLC, HPLC Solubility: chloroform, DMF, DMSO	Identity: confirmed by MS
2200	N-1-¹³C-Hexadecanoyl-D-erythro-sphingosylphosphorylcholine D-erythro-Sphingomyelin with 1- ¹³ C-palmitic acid; N-1- ¹³ C-Palmitoyl-sphingosylphosphorylcholine	1 mg	260.00
	$^{12}C_{38}^{13}CH_{79}N_2O_6P$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 703 Purity: 98+% by TLC Solubility: chloroform, ethanol, methanol	
2206	N-omega-CD₃-Octadecanoyl-D-erythro-sphingosine-1-phosphate C18:0-CD ₃ -Ceramide-1-phosphate; N-Stearoyl-CD ₃ -C1P	1 mg	365.00
	$C_{36}H_{69}D_3NO_6P$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 649 Purity: 98+% by TLC Solubility: chloroform/methanol/acetic acid, 60:15:25 chloroform/methanol/7.5M ammonium hydroxide 80:20:4	Identity: confirmed by MS


1914	N-Octadecanoyl-D₃₅-psychosine, (perdeuterated, C18:0 fatty acid) N-C18:0-D ₃₅ -Cerebrosides, perdeuterated; N-Stearoyl-D ₃₅ -psychosine, perdeuterated	5 mg	420.00
	$C_{42}H_{46}D_{35}NO_8$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 763 Purity: 98% by TLC Solubility: chloroform, hot ethanol, chloroform/methanol, 2:1 Deuterium labeled stearoyl sidechain	Identity: confirmed by MS
2209 *NEW*	¹³C₆-Glucosylsphingosine 1-(beta-D-Glucosyl-1,2,3,4,5,6- ¹³ C ₆)-sphingosine; ¹³ C ₆ - <i>lyso</i> -glucocerebroside	1 mg	550.00
	$C_{18}^{13}C_6H_{47}NO_7$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 468 Purity: 98+% by TLC, HPLC Solubility: ethanol, methanol, chloroform/methanol, 2:1	Identity: confirmed by MS
1533	N-omega-CD₃-Hexadecanoyl-glucopsychosine N-C16:0-CD ₃ -Glucopsychosine; N-C16:0-CD ₃ -Glucocerebroside; N-Palmitoyl-CD ₃ -glucopsychosine	1 mg	400.00
	$C_{40}H_{74}D_3NO_8$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 703 Purity: 98+% by TLC Solubility: chloroform/methanol, 2:1	Identity: confirmed by MS
1536	N-omega-CD₃-Octadecanoyl-sulfatide N-C18:0-CD ₃ -Sulfatide; N-Stearoyl-CD ₃ -sulfatide	1 mg	550.00
	$C_{42}H_{78}D_3NO_{11}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 811 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS
1534	N-omega-CD₃-Hexadecanoyl-lactosylceramide N-C16:0-CD ₃ -Lactosylceramide; N-Palmitoyl-CD ₃ -lactosylceramide	1 mg	460.00
	$C_{46}H_{84}D_3NO_{13}$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 865 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 5:1:0.1	Identity: confirmed by MS
1537	N-omega-CD₃-Octadecanoyl-ceramide trihexoside N-C18:0-D ₃ -CTH; N-C18:0-CD ₃ -Gb ₃ ; N-Octadecanoyl-CD ₃ -globotriaosylceramide; N-Stearoyl-CD ₃ -ceramide trihexoside	500 µg	495.00
	$C_{54}H_{98}D_3NO_{18}$ Source: semisynthetic, porcine RBC Appearance: solid Storage: -20°C	Mol. Wt.: 1055 Purity: 98+% by TLC, HPLC Solubility: DMSO, chloroform/methanol, 2:1	Identity: confirmed by MS
2050	N-omega-CD₃-Octadecanoyl monosialoganglioside GM₁ (NH₄⁺ salt) N-CD ₃ -Stearoyl GM ₁	500 µg	550.00
	$C_{73}H_{128}N_3O_{31}D_3 \cdot NH_3$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 1550 + NH ₃ Purity: 98% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	Identity: confirmed by MS

2051	N-<i>omega</i>-CD₃-Octadecanoyl monosialoganglioside GM₂ (NH₄⁺ salt) N-CD ₃ -Stearoyl GM ₂	250 µg	490.00
	C₆₇H₁₁₈D₃N₃O₂₆•NH₃ Source: semisynthetic, human Tay-Sachs	Mol. Wt.: 1388 + NH ₃ Purity: 98+% by TLC, MS	Identity: confirmed by MS
	Appearance: solid Storage: -20°C	Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	
2052	N-<i>omega</i>-CD₃-Octadecanoyl monosialoganglioside GM₃ (NH₄⁺ salt) N-CD ₃ -Stearoyl GM ₃	250 µg	410.00
	C₅₉H₁₀₅D₃N₂O₂₁•NH₃ Source: semisynthetic, bovine buttermilk	Mol. Wt.: 1185 + NH ₃ Purity: 98% by TLC	Identity: confirmed by MS
	Appearance: solid Storage: -20°C	Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water	
2054 *NEW*	N-<i>omega</i>-CD₃-Octadecanoyl disialoganglioside GD₃ N-CD ₃ -Stearoyl GD ₃	500 µg	550.00
	C₇₀H₁₂₂D₃N₃O₂₉ Source: semisynthetic bovine buttermilk	Mol. Wt.: 1476 Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid Storage: -20°C	Solubility: chloroform/methanol, 2:1, water	

Fluorescent Standards

Absorption: 460 nm Emission: 535 nm

1841	N-Hexanoyl-NBD-D-<i>erythro</i>-sphingosine	100 µg	155.00
1841-001	N-C6:0-NBD-Ceramide; N-C6:0-NBD-D- <i>erythro</i> -Sphingosine	1 mg	390.00
	C₃₀H₄₉N₅O₆ Source: synthetic	Mol. Wt.: 576 Purity: 98% by TLC	CAS#: 86701-10-2
	Appearance: solid Storage: -20°C	Solubility: chloroform, ethanol, methanol Melting Point (°C): 85-88	



1618	N-Dodecanoyl-NBD-D-<i>erythro</i>-sphingosine	100 µg	150.00
1618-001	N-C12:0-NBD-Ceramide; N-C12:0-NBD-D- <i>erythro</i> -Sphingosine	1 mg	375.00
	C₃₆H₆₁N₅O₆ Source: synthetic	Mol. Wt.: 660 Purity: 98% by TLC	Identity: confirmed by MS
	Appearance: solid Storage: -20°C	Solubility: methanol, chloroform/methanol, 2:1	

1857 1857-001	N-Hexanoyl-NBD-L-threo-sphingosine N-C6:0-NBD-Ceramide; N-C6:0-NBD-L-threo-Sphingosine	100 µg 1 mg	225.00 650.00
	$C_{30}H_{49}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 575 Purity: 98% by TLC Solubility: chloroform, ethanol, methanol	
1620	N-Dodecanoyl-NBD-L-threo-sphingosine N-C12:0-NBD-Ceramide; N-C12:0-NBD-L-threo-Sphingosine	100 µg	225.00
	$C_{36}H_{61}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 660 Purity: 98% by TLC Solubility: methanol, chloroform/methanol, 2:1	
1624	N-Hexanoyl-NBD-L-threo-dihydrosphingosine N-C6:0-NBD-Dihydroceramide; N-C6:0-NBD-L-threo-Dihydrosphingosine	100 µg	200.00
	$C_{30}H_{51}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 578 Purity: 98% by TLC Solubility: methanol, chloroform/methanol, 2:1	
1623	N-Dodecanoyl-NBD-L-threo-dihydrosphingosine N-C12:0-NBD-Dihydroceramide; N-C12:0-NBD-L-threo-Dihydrosphingosine	100 µg	200.00
	$C_{36}H_{63}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 662 Purity: 98% by TLC Solubility: methanol, chloroform/methanol, 2:1	
1626	N-Hexanoyl-NBD-D-erythro-dihydrosphingosine N-C6:0-NBD-Dihydroceramide; N-C6:0-NBD-D-erythro-Dihydrosphingosine	100 µg	200.00
	$C_{30}H_{51}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 578 Purity: 98% by TLC Solubility: methanol, chloroform/methanol, 2:1	Identity: confirmed by MS
1625	N-Dodecanoyl-NBD-D-erythro-dihydrosphingosine N-C12:0-NBD-Dihydroceramide; N-C12:0-NBD-D-erythro-Dihydrosphingosine	100 µg	200.00
	$C_{36}H_{63}N_5O_6$ Source: synthetic Appearance: solid Storage: -20°C	Mol. Wt.: 662 Purity: 98% by TLC Solubility: methanol, chloroform/methanol, 2:1	
1628	N-Hexanoyl-NBD-phytosphingosine N-C6:0-NBD-Phytoceramide; N-C6:0-NBD-Phytosphingosine	100 µg	200.00
	$C_{30}H_{51}N_5O_7$ Source: semisynthetic, bacteria Appearance: solid Storage: -20°C	Mol. Wt.: 594 Purity: 98% by TLC Solubility: methanol, chloroform/methanol, 2:1	CAS#: 477239-93-3
1627	N-Dodecanoyl-NBD-phytosphingosine N-C12:0-NBD-Phytoceramide; N-C12:0-NBD-Phytosphingosine	100 µg	185.00
	$C_{36}H_{63}N_5O_7$ Source: semisynthetic, bacteria Appearance: solid Storage: -20°C	Mol. Wt.: 678 Purity: 98% by TLC Solubility: methanol, chloroform/methanol, 2:1	CAS#: 388566-94-7

1912	N-Hexanoyl-NBD-sphingosylphosphorylcholine	100 µg	160.00
1912-001	N-C6:0-NBD-Sphingomyelin; N-C6:0-NBD-Sphingosylphosphorylcholine	1 mg	375.00
	$C_{35}H_{61}N_6O_9P$	Mol. Wt.: 740	CAS#: 94885-04-8
	Source: semisynthetic, bovine buttermilk	Purity: 98% by TLC	
	Appearance: solid	Solubility: chloroform, ethanol, methanol	
	Storage: -20°C	Mixture of <i>D-erythro</i> and <i>L-threo</i> isomers	
1619	N-Dodecanoyl-NBD-sphingosylphosphorylcholine	100 µg	200.00
1619-001	N-C12:0-NBD-Sphingomyelin; N-C12:0-NBD-Sphingosylphosphorylcholine	1 mg	395.00
	$C_{41}H_{73}N_6O_9P$	Mol. Wt.: 825	CAS#: 254117-01-6
	Source: semisynthetic, bovine buttermilk	Purity: 98% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: methanol, chloroform/methanol, 2:1	
	Storage: -20°C	Mixture of <i>D-erythro</i> and <i>L-threo</i> isomers	
1621	N-Hexanoyl-NBD-galactosylceramide	100 µg	250.00
	N-C6:0-NBD- <i>beta</i> -D-Galactosylsphingosine; N-C6:0-NBD-Cerebroside		
	$C_{36}H_{59}N_5O_{11}$	Mol. Wt.: 738	
	Source: semisynthetic, bovine	Purity: 98% by TLC	
	Appearance: solid	Solubility: methanol, chloroform/methanol, 5:1	
	Storage: -20°C		
1633	N-Dodecanoyl-NBD-galactosylceramide	100 µg	210.00
1633-001	N-C12:0-NBD- <i>beta</i> -D-Galactosylsphingosine; N-C12:0-NBD-Cerebroside	1 mg	450.00
	$C_{42}H_{71}N_5O_{11}$	Mol. Wt.: 822	CAS#: 474942-98-8
	Source: semisynthetic, bovine spinal cord	Purity: 98% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform, DMSO, chloroform/methanol, 2:1	
	Storage: -20°C		
2204	Lissamine-rhodamine B-dodecanoyl-galactosylceramide	500 µg	390.00
	Sulforhodamine B-C12:0 cerebroside		
	$C_{63}H_{99}N_4O_{14}S_2$	Mol. Wt.: 1201	
	Source: semisynthetic, bovine	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform/methanol 8:2, DMSO, DMF	
	Storage: -20°C		
	Absorption: 540 nm	Emission: 565 nm	
1622	N-Hexanoyl-NBD-glucosylceramide	100 µg	250.00
1622-001	N-C6:0-NBD- <i>beta</i> -D-Glucosylsphingosine; N-C6:0-NBD-Glucosylceramide, fluorescent	1 mg	400.00
	$C_{36}H_{59}N_5O_{11}$	Mol. Wt.: 738	
	Source: semisynthetic, bovine	Purity: 98% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: methanol, chloroform/methanol, 5:1	
	Storage: -20°C		
1632	N-Dodecanoyl-NBD-sulfatide	100 µg	170.00
1632-001	N-C12:0-NBD-Sulfatide; N-Dodecanoyl-NBD- <i>lyso</i> -sulfatide; N-Dodecanoyl-NBD-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	450.00
	$C_{42}H_{71}N_5O_{14}S$	Mol. Wt.: 901	
	Source: semisynthetic, bovine	Purity: 98+% by TLC	
	Appearance: solid	Solubility: chloroform/methanol, 2:1	
	Storage: -20°C		

1629	N-Hexanoyl-NBD-lactosylceramide	50 µg	265.00
1629-001	N-Hexanoyl-NBD- <i>beta</i> -D-lactosylsphingosine; N-C6:0-NBD- <i>beta</i> -D-Lactosylsphingosine; N-C6:0-NBD-Lactosylceramide	1 mg	500.00
	$C_{42}H_{69}N_5O_{16}$	Mol. Wt.: 900	
	Source: semisynthetic, bovine buttermilk	Purity: 98% by TLC	
	Appearance: solid	Solubility: chloroform/methanol, 2:1	
	Storage: -20°C		

1630	N-Dodecanoyl-NBD-lactosylceramide	50 µg	295.00
1630-001	N-Dodecanoyl-NBD- <i>beta</i> -D-lactosylsphingosine; N-C12:0-NBD- <i>beta</i> -D-Lactosylsphingosine; N-C12:0-NBD-Lactosylceramide	1 mg	500.00
	$C_{48}H_{81}N_5O_{16}$	Mol. Wt.: 984	
	Source: semisynthetic, bovine buttermilk	Purity: 98+% by TLC	
	Appearance: solid	Solubility: chloroform/methanol, 2:1	
	Storage: -20°C		

1631	N-Dodecanoyl-NBD-ceramide trihexoside	100 µg	265.00
1631-001	N-C12:0-NBD-CTH; N-C12:0-NBD-Globotriaosylceramide	1 mg	650.00
	$C_{54}H_{91}N_5O_{21}$	Mol. Wt.: 1145	
	Source: semisynthetic, porcine	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: DMSO; hot methanol, chloroform/methanol, 2:1	
	Storage: -20°C		

Biotin Labeled Standards

2081	N-Hexanoyl-biotin-D-erythro-sphingosine	5 mg	540.00
	N-C6:0-Biotin-D- <i>erythro</i> -ceramide		
	$C_{34}H_{62}N_4O_5S$	Mol. Wt.: 639	
	Source: synthetic	Purity: 98+% by TLC, HPLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform/methanol 2:1, DMF	
	Storage: -20°C		

2203	N-Hexanoyl-biotin-galactosylceramide	5 mg	500.00
	N-C6:0-Biotin- <i>beta</i> -D-galactosylsphingosine; N-C6:0-Biotin-cerebroside		
	$C_{40}H_{72}N_4O_{10}S$	Mol. Wt.: 801	
	Source: semisynthetic, bovine	Purity: 98+% by TLC, HPLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform/methanol 2:1, methanol, DMF	
	Storage: -20°C		

2085	N-Hexanoyl-biotin-glucosylceramide	5 mg	500.00
	N-C6:0-Biotin- <i>beta</i> -D-glucosylsphingosine; N-C6:0-Biotin-glucosylceramide		
	$C_{40}H_{72}N_4O_{10}S$	Mol. Wt.: 801	
	Source: semisynthetic, plant	Purity: 98+% by TLC	Identity: confirmed by MS
	Appearance: solid	Solubility: chloroform/methanol 2:1, methanol, DMF	
	Storage: -20°C		

2207	N-Hexanoyl-biotin-sulfatide N-C6:0-Biotin-sulfatide; N-Hexanoyl-biotin-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate	1 mg	450.00
	$C_{40}H_{72}N_4O_{13}S_2$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 881 Purity: 98+% by TLC Solubility: chloroform/methanol/water 2:1:0.1; methanol/water 9:1; DMF	Identity: confirmed by MS
2205	N-Hexanoyl-biotin-lactosylceramide N-C6:0-Biotin- <i>beta</i> -D-lactosylceramide	1 mg	280.00
	$C_{46}H_{82}N_4O_{15}S$ Source: semisynthetic, bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 963 Purity: 98+% by TLC Solubility: chloroform/methanol, 9:1, DMSO, DMF	Identity: confirmed by MS
2053	N-Hexanoyl-biotin-monosialoganglioside GM₁ Biotin-C6:0-GM ₁	500 µg	575.00
	$C_{71}H_{122}N_6O_{33}S$ Source: semisynthetic, bovine Appearance: solid Storage: -20°C	Mol. Wt.: 1620 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS
2055 *NEW*	N-Hexanoyl-biotin-disialoganglioside GD₃ Biotin-C6:0-GD ₃	500 µg	575.00
	$C_{68}H_{116}N_6O_{31}S$ Source: semisynthetic bovine buttermilk Appearance: solid Storage: -20°C	Mol. Wt.: 1546 Purity: 98+% by TLC Solubility: chloroform/methanol/DI water, 2:1:0.1	Identity: confirmed by MS

Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC
(actual composition may vary according to dietary history and growth condition of the source)

	Lecithin, egg	Phosphatidyl ethanolamine, egg	lyso-Lecithin, egg	Phosphatidylserine, bovine	Phosphatidylinositol, plant, wheat germ	Sulfatides, bovine	Cerebrosides, bovine	Sphingomyelin, bovine	Phosphatidic acid, egg	Ceramides (mixture)
Catalog Number	1044	1045	1046	1047	1048	1049	1050	1051	1053	1056
Fatty Acids										
C14:0										
C16:0	31	17	72	1	42	trace	trace	4	39	trace
C16:1										
C18:0	16	29	24	42		5	4	40	12	4
C18:1	31	17	3	27	6	trace			34	
C18:2	16	11			47				15	
C18:3					5					
C20:0				1		1	1	3		1
C20:1				4						
C20:4		12		4						
C21:0										
C22:0				1		7	4	13		4
C22:1				1		trace				
C22:6				7						
C23:0							2	2		2
C24:0						18	10	9		10
C24:1						29	15	22		15
C25:0						2	3			9
C25:1						2	1			1
C26:0						1	2			2
C26:1						3	1			1
C27:0						1	2			2
C27:1							2			2
C14:0 2-OH										
C16:0 2-OH										
C18:0 2-OH						5	15			15
C20:0 2-OH						trace	1			1
C22:0 2-OH						3	6			6
C23:0 2-OH							5			5
C24:0 2-OH						10	17			17
C24:1 2-OH						6	6			
C25:0 2-OH						2	3			3
C25:1 2-OH										
C26:0 2-OH										
C26:1 2-OH										
C16 cis 9,10 methylene										
C18 cis 9,10 methylene										
Others	6	14	1	12	0	5	0	7	0	0
Total	100	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Glucocerebrosides, Gaucher's spleen	Monogalactosyl di glyceride (hydrogenated), plant	Digalactosyl Diglyceride (hydrogenated), plant	Monosialo ganglioside GM ₁	Disialoganglioside GD _{1a}	Trisialoganglioside GT _{1b}	Gangliotetraosyl ceramide	Mixed Gangliosides, purified, bovine	Cerebroside; Kerasin (top spot)	Ceramide trihexosides
Catalog Number	1057	1058	1059	1061	1062	1063	1064	1065	1066	1067
Fatty Acids										
C14:0					1		trace	trace		
C16:0	26	23	9	2	1	1	1	1	trace	3
C16:1										
C18:0	9	77	91	90	89	87	86	86	5	2
C18:1						1	3	3		2
C18:2										
C18:3										
C20:0	5			3	2	4	4	4	1	2
C20:1										
C20:4										
C21:0										
C22:0	26			1	1	1	2	2	9	17
C22:1									trace	
C22:6										
C23:0	5					1	1	1	5	1
C24:0	22					1	1	1	25	29
C24:1	6			1		1	2	2	43	5
C25:0									3	
C25:1									3	
C26:0									2	
C26:1									4	
C27:0										
C27:1										
C14:0 2-OH										
C16:0 2-OH										
C18:0 2-OH										
C20:0 2-OH										
C22:0 2-OH										3
C23:0 2-OH										1
C24:0 2-OH										19
C24:1 2-OH										10
C25:0 2-OH										
C25:1 2-OH										
C26:0 2-OH										
C26:1 2-OH										
C16 cis 9,10 methylene										
C18 cis 9,10 methylene										
Others	1	0	0	3	6	3	0	0	0	6
Total	100	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Globosides, porcine	Lecithin, bovine	Esterified Steryl Glucosides	Cerebroside; Phrenosin (bottom spot)	Phosphatidyl ethanolamine, plant	Ceramides (non-hydroxy)	Ceramides (hydroxy)	Sphingomyelin, porcine RBC	Sphingomyelin, buttermilk
Catalog Number	1068	1070	1118	1138	1301	1322	1323	1328	1329
Fatty Acids									
C14:0		trace							1
C16:0	2	35	34		22			25	14
C16:1		1							
C18:0	2	14	8		3	11		7	3
C18:1		33	8		7				
C18:2			36		60				
C18:3			4		8				
C20:0	2		1			2		3	1
C20:1									
C20:4									
C21:0									
C22:0	20		4			10		9	26
C22:1									
C22:6									
C23:0	2		2			6		1	30
C24:0	33		2			24		22	21
C24:1	5					31		22	3
C25:0						3			
C25:1						3			
C26:0	2					2			
C26:1						3			
C27:0									
C27:1									
C14:0 2-OH									
C16:0 2-OH									
C18:0 2-OH				36			24		
C20:0 2-OH				1			1		
C22:0 2-OH	4			8			8		
C23:0 2-OH				6			6		
C24:0 2-OH	19			25			35		
C24:1 2-OH	9			9			17		
C25:0 2-OH				4			4		
C25:1 2-OH				2					
C26:0 2-OH				2					
C26:1 2-OH				2			2		
C16 cis 9,10 methylene									
C18 cis 9,10 methylene									
Others	0	17	1	5	0	5	3	11	1
Total	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Spingomyelin, egg	Phosphatidylinositol, plant, soy	Lactosylceramides	Disialoganglioside GD _{1b} , bovine	Monosialoganglioside GM ₂	Monosialoganglioside GM ₃	Disialoganglioside GD ₃ , buttermilk	Lactosylceramides buttermilk	Ceramide trihexosides (top spot)
Catalog Number	1332	1336	1500	1501	1502	1503	1504	1507	1513
Fatty Acids									
C14:0	trace			trace					
C16:0	72	32	14	1	2	6	8	12	1
C16:1									
C18:0	8	7	6	86	82	1	1	1	1
C18:1	3	7	4	3					
C18:2		47							
C18:3		6							
C20:0	2		1	4	7	1	1	1	2
C20:1									
C20:4									
C21:0						1	2		
C22:0	5		9	2	4	23	24	25	22
C22:1									
C22:6									
C23:0	1		1	1	trace	36	35	36	2
C24:0	2		15	1	1	22	21	21	58
C24:1	4		5	2	2	3	3		7
C25:0								1	1
C25:1									
C26:0									5
C26:1									
C27:0									
C27:1									
C14:0 2-OH									
C16:0 2-OH									
C18:0 2-OH			trace						
C20:0 2-OH									
C22:0 2-OH			8						
C23:0 2-OH									
C24:0 2-OH			24						
C24:1 2-OH			13						
C25:0 2-OH									
C25:1 2-OH									
C26:0 2-OH									
C26:1 2-OH									
C16 cis 9,10 methylene									
C18 cis 9,10 methylene									
Others	3	1	0	0	2	7	5	3	1
Total	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Ceramide trihexosides (bottom spot)	Tetraialoganglioside GQ _{1b}	Glucocerebrosides, buttermilk	Glucocerebrosides, plant	Mixed Gangliosides, purified, porcine	Fucosylated monosialoganglioside GM ₁	Disialoganglioside GD ₂	Monosialoganglioside GM ₄
Catalog Number	1514	1516	1521	1522	1525	1526	1527	1535
Fatty Acids								
C14:0								
C16:0	3	5	7		1	8	1	4
C16:1		1						
C18:0		80	2		87	2	89	2
C18:1		2						
C18:2		3						
C18:3								
C20:0		4	1		4	13	7	trace
C20:1								trace
C20:4								
C21:0			1					
C22:0	2	2	27		1	43	1	3
C22:1								4
C22:6								
C23:0			36		1	3	1	4
C24:0	3		23		1	26		6
C24:1					2	5	1	4
C25:0			1					
C25:1								
C26:0								
C26:1								
C27:0								
C27:1								
C14:0 2-OH				trace				
C16:0 2-OH				79				
C18:0 2-OH	1			trace				1
C20:0 2-OH	1							3
C22:0 2-OH	11			8				25
C23:0 2-OH	1			1				17
C24:0 2-OH	52			9				18
C24:1 2-OH	25							7
C25:0 2-OH								
C25:1 2-OH								
C26:0 2-OH								
C26:1 2-OH								
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	1	3	2	3	3			2
Total	100	100	100	100	100	100	100	100

Literature References

1. B.A. Fenderson, E.M. Eddy, S.Hakomori, *BioEssays* 12, 173, 1990
2. A. Gorio et al., *Exp. Brain Res.*, Suppl. 13, 283, 1984
3. F. Di Gregorio et al., *Neuropediatrics*, Suppl. 15, 93, 1984
4. J.S. Schneider et al., *Science*, 256, 843, 1992
5. R.W. Leeden, R.K. Yu, *Methods Enzymol*, 83, 139, 1982
6. Borman, S., *Chemical and Engineering News* 82:32, 31-35, 2004
7. M. Faucher et al., *J. Biol. Chem.* 263, 5319, 1988
8. B. M. Buehrer, R.M. Bell, *Adv. in Lipid Res.*, 26, 59, 1993
9. C. W. Sachs et al., *J. Biol. Chem.*, 270, 26639, 1995
10. Radin, N. S., *Biochem. J.* 371:243, 2003
11. Radin, N. S., *Biorg. Med. Chem.* 11:2123, 2003
12. Radin, N. S., *Cancer Investigation* 20:1, 2002
13. Sietsma H., Veldman R.J., Vander Kolk D., Ausema B., Nijhof W., Kamps W., Vellenga E., Kok J.W. *Clin. Cancer Res.* 6, 942-948, 2000
14. Shayman, J. A., *Kidney Int.* 58, 11-26, 2000
15. Kolesnick, R. N. et al. *J. Cell Physiol.* 184, 285-300, 2000
16. Smith, W. L. and Merrill, A. H., Jr. *J. Biol. Chem.* 277, 25841-25842, 2002
17. Pyne, S. and Pyne, N. J. *Biochem. J.* 349, 385-402, 2000
18. Merrill, A. H. Jr. *J. Biol. Chem.* 277, 25843-25846, 2002
19. Hannun, Y. A. and Obeid, L., *J. Biol. Chem.* 277, 25847-25850, 2002
20. van Meer, G. and Lisman, Q. J. *Biol. Chem.* 277, 25855-25858, 2002
21. Kolter, T., Proia, R. L. and Sandhoff, K. *J. Biol. Chem.* 277, 25859-25862, 2002
22. Radin, N. S., *Cancer Invest.* 20, 779-786, 2002
23. Tettamanti, G., Bassi R., Viani, P., Riboni, L., *Biochimie* 85(3-4), 423-437, 2003
24. Fredman, P., Hedberg, K., Brezicka, T., *Biodrugs* 17(3) 155-167, 2003
25. Hoeckstra, D., et al. *J. Lipid Res.* 44, 869-877, 2003
26. R.R, Vunnam, N.S. Radin, *Chem. Phys. Lipids*, 265, 1980
27. J. Inokuchi and N.S. Radin, *J. Lipid Res.*, 28, 565, 1987
28. N.S. Radin et al. *J. Biochem.*, 111, 191, 1992
29. J. Inokuchi et al., *Cancer letters*, 38, 23, 1987
30. Y. Lavie et al., *J. Biol. Chem.*, 271, 19530, 1996
31. Y. Lavie et al., *J. Biol. Chem.*, 271, in press, 1996
32. Y. Hannun et al., *Science*, 235, 670, 1987
33. S. Spiegel et al., *Proc. Intern. Conf. Biol. Function Glycosphingolipids*, Santa Barbara, CA, 1990
34. Jacqueline M. Kravcka, Li Li, Zdzislaw M. Szulc, Jacek Bielawski, Besium Ogretmen, Yusuf A. Hannun, Lina M. Obeid, and Alicja Bielawska. *J. Biol. Chem.*, 10, 1074/jbc. M700647200, February 5, 2007
35. M. Sugita et al. *Biochim. Biophys. Acta*, 398, 125, 1975
36. A. Bielawska et al. *J. Biol. Chem.*, 271, 12646, 1996
37. A. Bielawska et al. *ibid.* 267, 18493, 1992
38. J. M. L. Hauser et al., *J. Biol. Chem.* 269, 6803, 1994
39. Y. Hannun, *ibid.*, 269, 3125, 1994
40. R. Kolesnick, D.W. Golde, *Cell*, 77, 325, 1984
41. M.W. Pariza et al., *Cancer Res.* 43, 2444s, 1983
42. Y. L Ha et al., *J. Agr. Food Chem.*, 37, 75, 1989
43. P.W. Parodi, *Austral. J. Dairy Tech.*, 49, 93, 1994
44. S. Banni et al. *Abstr. 87th AOCS Mtg.*, 28, 1996
45. C. Ip et al., *Cancer Res.* 51, 6118, 1991
46. M. A. Belury, *Nutr. Rev.* 53 (4 Pt. 1), 83, 1995
47. C. Liew et al. *Carcinogenesis*, 16, 3037, 1995
48. T. D. Shultz et al. *Cancer Let.*, 63, 125, 1992
49. R.J. Nicolosi, *Abstr. 87th AOCS Mtg.*, 1996
50. K.N. Lee et al., *Atheroscl.*, 108(1), 19, 1994
51. B.F. Haumann, *Inform.*, 7(2), 152, 1996
52. M.W. Pariza et al., *Abstr.*, 87th AOCS Mtg., 1996
53. M.A. Belury et al., *Lipids*, 32, 199, 1997
54. Houseknecht, K.L. et al., *Biochem. Biophys. Res. Comm.*, 224, 678, 1998
55. W.W. Christie et al. *JAOCS*, 74, 1231, 1997
56. Sehat, N. et al., *Lipids*, 33, 217, 1998
57. Sébédio, J. L. et al., *Biochim. Biophys. Acta*, 1345,5, 1997

Note: We are pleased and honored to have the above list of scientists in the field of Lipid Research. In case your name and contribution is not listed above, we apologize. If you would like your publication listed in our next catalog, please send your name and publication to the attention of Marketing Department.

Cross Reference for Product Numbers and Catalog Pages

1006	81	1067-10	31	1154	61
1008	53	1068	33, 102	1155	64
1009	54	1069	47	1156	64
1010	54	1070	46, 102	1157	57
1011	54	1071	79	1161	54
1012	54	1072	79	1162	54
1013	54	1073	79	1163	53
1014	54	1074	80	1164	53
1015	55	1075	91	1165	53
1016	58	1077	91	1166	53
1017	58	1081	84	1167	63
1018	55	1083	86	1175	63
1019	55	1084	86	1177	85
1020	55	1085	86	1179	62
1021	55	1086	86, 87	1181	67
1022	59	1087	86, 87	1186	56
1023	59	1088	86, 87	1187	56
1024	60	1089	86, 87	1192	62
1025	60	1093	84	1193	62
1026	60	1095	89	1194	63
1027	61	1097	89	1195	77
1028	55	1098	89	1196	52
1029	55	1099	89, 90	1197	52
1030	55	1100	89, 90	1198	53
1031	56	1101	89, 90	1199	53
1032	62	1102	89, 90	1200	52
1033	62	1103	89, 90	1203	59
1034	63	1104	89, 90	1204	59
1035	56	1105	91	1205	62
1036	56	1106	90	1206	62
1037	56	1108	90	1207	77
1038	57	1113	82	1208	59
1040	58	1114	91	1209	58
1041	64	1115	81	1210	58
1042	63	1116	82	1233	61, 68
1044	45, 100	1117	83	1234	61, 68
1045	47, 100	1118	83, 102	1235	78
1046	46, 100	1119	82	1236	78
1047	46, 100	1120	82	1238	78
1048	46, 100	1121	82	1240	61, 67
1049	26, 100	1122	82	1241	56
1050	21, 100	1123	82	1242	56
1051	17, 100	1124	85	1243	58
1051-1	17	1125	85	1244	63
1053	46, 100	1127	91	1245	66
1056	13, 100	1128	92	1248	67
1057	24, 101	1129	92	1249	67
1057-25	24	1130	92	1251	57
1058	52, 101	1131	65, 85	1252	57
1059	52, 101	1136	64	1254	67
1061	36, 101	1138	21, 102	1255	66
1061-50	36	1147	58, 64	1256	67
1062	38, 101	1148	58, 64	1257	67
1063	39, 101	1149	59, 64	1261	53
1064	36, 101	1150	59, 65	1262	60, 65
1065	39, 101	1151	60, 65	1263	60, 65
1066	21, 101	1152	60, 65	1264	63
1067	31, 101	1153	61	1265	63

1266	59	1507-50	29	1631-001	32, 35, 98
1267	60	1508	39, 92	1632	28, 34, 97
1269	62	1509	40, 92	1632-001	28, 34, 97
1271	57	1510	40, 92	1633	23, 34, 97
1273	57	1511	40, 92	1633-001	23, 34, 97
1275	57	1512	36	1656	75
1276	61	1513	31, 103	1657	75
1277	61	1514	31, 104	1701	68
1278	66	1516	39, 104	1702	69
1301	47, 102	1517	29	1703	69
1303	51	1518	36	1704	69
1303-2	51	1520	31	1705	69
1305	22	1521	24, 104	1706	69
1306	25	1521-50	24	1707	69
1310	25	1522	24, 104	1708	69
1318	19	1522-100	24	1709	69
1319	19	1523	32	1710	70
1320	5, 45	1524	32	1711	70
1321	19	1525	39, 104	1712	70
1321-05	19	1526	37, 104	1713	70
1322	13, 102	1527	38, 104	1714	70
1322-05	13	1528	31	1715	70
1323	13, 102	1529	32	1716	70
1323-05	13	1530	31	1717	44
1324	4	1531	25	1718	44
1325	22	1532	29	1719	43
1326	4	1533	25, 33, 94	1720	43
1327	18	1534	30, 34, 94	1722	70, 85
1328	17, 102	1535	38, 104	1725	71
1329	17, 102	1536	28, 33, 94	1726	71
1330	5	1537	32, 34, 94	1727	72
1332	17, 103	1538	30	1728	72
1333	8	1539	25	1729	72
1334	22	1540	28, 30	1730	72
1335	22	1600	76	1731	72
1336	46, 103	1601	76	1732	72
1400	48	1602	76	1733	72
1425	48	1603	76	1734	72
1426	48	1605	76	1735	73
1427	49	1606	76	1736	73
1428	47	1612	77	1739	73
1429	48	1613	77	1740	73
1430	48	1614	77	1741	73
1431	49	1615	76	1742	73
1432	49	1616	77	1743	73
1433	50	1618	15, 95	1744	73
1434	50	1618-001	15, 95	1745	71
1435	50	1619	20, 97	1746	71
1436	50	1619-001	20, 97	1747	71
1437	49	1620	15, 96	1748	71
1438	50	1621	23, 34, 97	1749	43
1439	51	1622	26, 34, 97	1751	41, 78
1442	48	1622-001	26, 34, 97	1753	43
1443	49	1623	16, 96	1754	74
1444	50	1624	15, 96	1755	43
1445	49	1625	16, 96	1756	43
1500	29, 103	1626	16, 96	1758	68
1501	38, 103	1627	16, 96	1759	68
1502	37, 103	1628	16, 96	1760	74
1503	37, 103	1629	30, 35, 98	1761	74
1504	38, 103	1629-001	30, 35, 98	1773	51
1504-25	38	1630	30, 35, 98	1773-1	51
1505	39, 92	1630-001	30, 35, 98	1775	51
1507	29, 103	1631	32, 35, 98	1775-1	51

1790	79	1884	75	2035	14
1792	77	1888	28	2036	14
1797	80	1889	42	2037	8
1802	2	1890	18	2038	9
1803	19	1891	5	2039	10
1806	2	1892	5	2041	12
1807	3, 42	1893	5	2043	12
1807-025	3, 42	1894	14	2044	11
1809	8	1895	14	2045	13
1810	8	1896	6, 45	2046	20
1818	75	1897	14	2047	13
1819	75	1900	7	2048	10
1822	78	1901	7	2049	10
1823	78	1903	8	2050	37, 94
1826	2	1904	26	2051	37, 95
1828	7	1907	17	2052	37, 95
1829	7	1910	11	2053	37, 99
1830	8	1911	18	2054	38, 95
1831	3	1912	20, 97	2055	39, 99
1832	9	1912-001	20, 97	2076	27
1833	2	1913	19	2077	11
1834	11	1914	23, 33, 94	2078	12
1838	2	1915	9	2079	2, 93
1839	4	1916	10	2080	10
1840	3	1917	18	2081	11, 98
1841	15, 95	1918	18	2082	3
1841-001	15, 95	1930	10	2083	12
1842	11	1931	28	2084	10
1843	9	1932	27	2085	26, 98
1845	4	1933	27	2086	24
1846	3	1934	27	2087	22
1847	7	1935	27	2088	29
1848	7	1936	8	2089	25
1850	9	1937	22	2090	29
1851	4	1937-50	22	2091	22
1852	20	1938	27	2092	26
1854	12	1939	9	2093	12
1855	9	1939-25	9	2109	80
1857	15, 96	1945	26, 44	2110	80
1857-001	15, 96	1946	23, 45	2111	81
1858	44	1947	32	2112	81
1859	42	1950	40	2200	18, 93
1860	42	1951	40	2201	9, 93
1865	44	1954	40	2202	12, 93
1868	44	1960	41	2203	23, 98
1875	27	1964	41	2204	23, 35, 97
1876	4	1977	40	2205	30, 99
1877	74	2009	83	2206	20, 93
1878	74	2010	84	2207	28, 99
1880	75	2011	85	2208	10, 93
1881	74	2012	84	2209	24, 33, 94
1882	74	2013	84	2210	15, 93
1883	75	2034	14	4210	83

Product Name Index

(+)- <i>delta</i> -Tocopherol	79	Alditol Acetate Mixture-2	85
1,2-Diheptadecanoyl-sn-glycero-3-phosphorylcholine	48	<i>alpha</i> -Tocotrienol	80
1,2-Dilauroyl-sn-glycero-3-phosphorylcholine	48	Anti-ganglioside asialo GM ₁	40
1,2-Dilauroyl-sn-glycero-3-phosphorylethanolamine	50	Anti-ganglioside asialo GM ₂	40
1,2-Dilauroyl-sn-glycero-3-phosphorylglycerol	49	Anti-ganglioside GD _{1b}	41
1,2-Dimyrystoyl-sn-glycero-3-phosphatidic acid	47	Anti-ganglioside GD ₃	40
1,2-Dimyrystoyl-sn-glycero-3-phosphorylcholine	48	Anti-ganglioside GM ₁	40
1,2-Dimyrystoyl-sn-glycero-3-phosphorylethanolamine	50	Anti-globoside GL-4	41
1,2-Dimyrystoyl-sn-glycero-3-phosphorylglycerol	49	B	
1,2-Dipalmitoyl-sn-glycero-3-phosphatidic acid	48	Bacterial Acid Methyl Esters CP Mixture	91
1,2-Dipalmitoyl-sn-glycero-3-phosphorylcholine	48	<i>beta</i> -Sitostanol	82
1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine	50	<i>beta</i> -Tocotrienol	80
1,2-Dipalmitoyl-sn-glycero-3-phosphorylglycerol	49	C	
1,2-Distearoyl-phosphatidylethanolamine-methyl-polyethyleneglycol conjugate-2000	51	Ceramide trihexosides	31, 101
1,2-Distearoyl-sn-glycero-3-phosphatidic acid	48	Ceramide trihexosides (bottom spot)	31, 104
1,2-Distearoyl-sn-glycero-3-phosphorylcholine	49	Ceramide trihexosides (top spot)	31, 103
1,2-Distearoyl-sn-glycero-3-phosphorylethanolamine	50	Ceramides (hydroxy)	13, 102
1,2-Distearoyl-sn-glycero-3-phosphorylglycerol	50	Ceramides (mixture)	13, 100
10(E),12(Z)-Octadecadienoic acid	67	Ceramides (non-hydroxy)	13, 102
11-Hexadecenoic acid (92% <i>cis</i> , 8% <i>trans</i>)	59	Cerebroside; Kerasin (top spot)	21, 101
12-Methyltetradecanoic acid	76	Cerebroside; Phrenosin (bottom spot)	21, 102
¹³ C ₆ -Glucosylsphingosine	24, 33, 94	Cerebrosides, bovine	21, 100
13-Methyltetradecanoic acid	76	Cholesterol	81
14-Methylhexadecanoic acid	77	<i>cis</i> -9,10-Methyleneoctadecanoic acid (all <i>cis</i> -9,10)	78
15-Hydroxypentadecanoic acid	74	<i>Cis-Trans</i> FAME Isomer Standard Mixture	65, 85
15-Methylhexadecanoic acid	76	Conduritol B Epoxide	42
17-Hydroxyheptadecanoic acid	74	Coprostanol	82
1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylcholine	49	D	
1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylglycerol	50	D,L-2,6-Dimethylheptanoic acid	77
1-Palmitoyl-sn-glycero-3-phosphorylcholine	49	D,L-C16-Dihydrosphingosine (mixed isomers)	4
20-Hydroxyeicosanoic acid	74	D,L- <i>erythro</i> -C20-Dihydrosphingosine	4
22-Hydroxydocosanoic acid	75	D,L- <i>erythro</i> -Dihydrosphingosine	4
2-Acetyl-4-(1R, 2S, 3R, 4-tetrahydroxybutyl)-imidazole	44	D,L- <i>erythro</i> -PDMP	43
2-Fluoropalmitic acid	44	D,L- <i>erythro</i> -PPMP	43
2-Hydroxy Methyl Ester Mixture	70, 85	D,L- <i>threo</i> -PDMP	43
2-Hydroxydecanoic acid	68	D,L- <i>threo</i> -PPMP	43
2-Hydroxydocosanoic acid	70	<i>delta</i> -Tocotrienol	81
2-Hydroxydodecanoic acid	68	D- <i>erythro</i> -C12-Sphingosine	2
2-Hydroxyeicosanoic acid	69	D- <i>erythro</i> -C14-Sphingosine	2
2-Hydroxyhexadecanoic acid	69	D- <i>erythro</i> -C17-Sphingosine	3
2-Hydroxyoctadecanoic acid	69	D- <i>erythro</i> -C20-Dihydrosphingosine	4
2-Hydroxytetradecanoic acid	70	D- <i>erythro</i> -C20-Sphingosine	3
2-Hydroxytricosanoic acid	70	D- <i>erythro</i> -Dihydrosphingosine	3
2-Hydroxyundecanoic acid	72	D- <i>erythro</i> -Dihydrosphingosine-1-phosphate	20
3-Hydroxydodecanoic acid	72	D- <i>erythro</i> -Sphingosine	2
3-Hydroxyheptadecanoic acid	73	D- <i>erythro</i> -Sphingosine, D9	2, 93
3-hydroxyhexadecanoic acid	73	D- <i>erythro</i> -Sphingosine-1-phosphate	19
3-Hydroxyhexanoic acid	71	D- <i>erythro</i> -Sphingosylphosphorylcholine	19
3-Hydroxynonanoic acid	71	Digalactosyldiglyceride (hydrogenated), plant	52, 101
3-Hydroxyoctadecanoic acid	73	Disialoganglioside GD _{1a}	38, 101
3-Hydroxyoctanoic acid	71	Disialoganglioside GD _{1b} , bovine	38, 103
3-Hydroxytetradecanoic acid	73	Disialoganglioside GD ₂	38, 104
3-Hydroxytridecanoic acid	72	Disialoganglioside GD ₃ , buttermilk	38, 103
3-Hydroxyundecanoic acid	72	Disialoganglioside Mixture	40, 92
3-keto-C12-Dihydrosphingosine•HCl	5	D-MAPP	42
3-keto-C6-Dihydrosphingosine•HCl	5	Docosahexaenoic acid (all <i>cis</i> -4,7,10,13,16,19)	64
3-keto-C8-Dihydrosphingosine•HCl	5	Docosanoic acid	56
3-keto-Dihydrosphingosine•HCl	4	Docosapentaenoic acid (all <i>cis</i> -7,10,13,16,19)	63
5- <i>alpha</i> -Cholestane	81	Docosenoic acid (<i>cis</i> -13)	63
9(E),11(E)-Octadecadienoic acid	67	Dodecanoic acid	53
9(Z),11(E)-Octadecadienoic acid	66	D- <i>threo</i> -Dihydrosphingosine	4
9(Z),11(E)-Octadecadienoic acid (Na ⁺ salt)	66	D- <i>threo</i> -PDMP	43
9(Z),11(Z)-Octadecadienoic acid	67	D- <i>threo</i> -PPMP	44
A		E	
Alditol Acetate Mixture-1	85	Eicosadienoic acid (all <i>cis</i> -11,14)	62

Eicosanoic acid	55	<i>lyso</i> -Lactosylceramide, bovine buttermilk	29
Eicosapentaenoic acid (all <i>cis</i> -5,8,11,14,17)	63	<i>lyso</i> -Lactosylceramide, synthetic	29
Eicosatetraenoic acid (all <i>cis</i> -5,8,11,14)	63	<i>lyso</i> -Lecithin, egg	46, 100
Eicosenoic acid (<i>cis</i> -11)	62	<i>lyso</i> -Monosialoganglioside GM ₁	36
Ergosterol	82	<i>lyso</i> -Sulfatide	26
Esterified Steryl Glucosides	83, 102		
F			
FIM-FAME-6 Mixture	83		
FIM-FAME-7 Mixture	84		
FIM-FAME-8 Mixture	84		
FIM-FAME-9 Mixture	84		
Fucosylated monosialoganglioside GM ₁	37, 104		
G			
<i>gamma</i> -Tocotrienol	81		
Gangliotetraosylceramide	36, 101		
Gangliotetraosylceramide and Sialosyl Derivatives Mixture	40, 92		
Gangliotriaosylceramide	36		
GLC-10 Mixture	89		
GLC-100 Mixture	90		
GLC-110 Mixture	91		
GLC-30 Mixture	89		
GLC-40 Mixture	89		
GLC-50 Mixture	90		
GLC-60 Mixture	90		
GLC-70 Mixture	90		
GLC-80 Mixture	90		
GLC-90 Mixture	90		
Globosides	33, 102		
Glucocerebrosides, buttermilk	24, 104		
Glucocerebrosides, Gaucher's spleen	24, 101		
Glucocerebrosides, plant	24, 104		
Glucosylsphingosine, buttermilk	25		
Glucosylsphingosine, plant	25		
Glucosylsphingosine, synthetic	24		
H			
Heneicosanoic acid	56		
Heptadecanoic acid	55		
Heptadecenoic acid (<i>cis</i> -10)	59		
Heptanoic acid	52		
Hexacosanoic acid	57		
Hexadecanoic acid	54		
Hexadecenoic acid (<i>cis</i> -6)	58		
Hexadecenoic acid (<i>cis</i> -9)	58		
Hexadecenoic acid (<i>trans</i> -9)	58, 64		
K			
KEL-FIM-FAME-5 Mixture	83		
L			
Lactosylceramide and Sialosyl Derivatives Mixture	40, 92		
Lactosylceramides, bovine buttermilk	29, 103		
Lactosylceramides, porcine RBC	29, 103		
Lanosterol	82		
Lecithin, bovine	46, 102		
Lecithin, egg	45, 100		
<i>L-erythro</i> -Dihydrosphingosine	3		
<i>L-erythro</i> -Sphingosine	2		
Lissamine-rhodamine B-dodecanoyl-galactosylceramide	23, 35, 97		
L-MAPP	42		
Long Chain Fatty Acid Methyl Ester Mixture	85		
<i>L-threo</i> -Dihydrosphingosine (Safingol)	3, 42		
<i>L-threo</i> -PDMP	43		
<i>L-threo</i> -PPMP	44		
<i>L-threo</i> -Sphingosine	2		
<i>L-threo</i> -Sphingosylphosphorylcholine	19		
<i>lyso</i> -Ceramide trihexoside	31		
<i>lyso</i> -Dihydrosphingomyelin	19		
		Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i> , (>50% pure)	51
		Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i> , (>95% pure)	51
		Methyl 10(E),12(Z)-octadecadienoate	67
		Methyl 10-methylhexadecanoate	77
		Methyl 11-methyl dodecanoate	75
		Methyl 12-methyltetradecanoate	77
		Methyl 12-methyltridecanoate	75
		Methyl 13-methylpentadecanoate	77
		Methyl 13-methyltetradecanoate	76
		Methyl 14-methylhexadecanoate	77
		Methyl 14-methylpentadecanoate	76
		Methyl 15-hydroxypentadecanoate	74
		Methyl 15-methylhexadecanoate	76
		Methyl 17-hydroxyheptadecanoate	74
		Methyl 17-methyloctadecanoate	76
		Methyl 20-hydroxyeicosanoate	74
		Methyl 21-hydroxyheneicosanoate	75
		Methyl 22-hydroxydocosanoate	75
		Methyl 27-hydroxyheptacosanoate	75
		Methyl 2-fluoropalmitate	44
		Methyl 2-hydroxydecanoate	68
		Methyl 2-hydroxydocosanoate	70
		Methyl 2-hydroxydodecanoate	69
		Methyl 2-hydroxyeicosanoate	70
		Methyl 2-hydroxyhexadecanoate	69
		Methyl 2-hydroxyoctadecanoate	69
		Methyl 2-hydroxytetracosanoate	70
		Methyl 2-hydroxytetradecanoate	69
		Methyl 2-hydroxytricosanoate	70
		Methyl 30-hydroxytriacontanoate	75
		Methyl 3-hydroxydecanoate	72
		Methyl 3-hydroxydodecanoate	72
		Methyl 3-hydroxyheptadecanoate	73
		Methyl 3-hydroxyhexadecanoate	73
		Methyl 3-hydroxyhexanoate	71
		Methyl 3-hydroxynonanoate	71
		Methyl 3-hydroxyoctadecanoate	73
		Methyl 3-hydroxyoctanoate	71
		Methyl 3-hydroxytetradecanoate	73
		Methyl 3-hydroxytridecanoate	72
		Methyl 3-hydroxyundecanoate	72
		Methyl 9(E),11(E)-octadecadienoate	67
		Methyl 9(Z),11(E)-octadecadienoate	66
		Methyl 9(Z),11(Z)-octadecadienoate	67
		Methyl <i>alpha</i> -eleostearate	61, 68
		Methyl <i>cis</i> -9,10-methyleneoctadecanoate	78
		Methyl decanoate	53
		Methyl docosahexaenoate (all <i>cis</i> -4,7,10,13,16,19)	64
		Methyl docosanoate	56
		Methyl docosapentaenoate (all <i>cis</i> -7,10,13,16,19)	63
		Methyl docosenoate (<i>cis</i> -13)	63
		Methyl dodecanoate	54
		Methyl dotriacontanoate	57
		Methyl eicosadienoate (all <i>cis</i> -11,14)	62
		Methyl eicosanoate	56
		Methyl eicosapentaenoate (all <i>cis</i> -5,8,11,14,17)	63
		Methyl eicosatetraenoate (all <i>cis</i> -5,8,11,14)	63
		Methyl eicosatrienoate (all <i>cis</i> -5,8,11)	62
		Methyl eicosatrienoate (all <i>cis</i> -8,11,14)	62
		Methyl eicosenoate (<i>cis</i> -11)	62
		Methyl heneicosanoate	56
		Methyl heptadecanoate	55
		Methyl heptadecenoate (<i>cis</i> -10)	59

Methyl heptanoate	52	N-Decanoyl-D- <i>erythro</i> -sphingosine	8
Methyl hexacosanoate	57	N-Docosanoyl-D- <i>erythro</i> -sphingosylphosphorylcholine	18
Methyl hexadecanoate	55	N-Docosanoyl-glucopsychosine	25
Methyl hexadecenoate (<i>cis</i> -9)	58	N-Dodecanoyl- <i>beta</i> -D-galactosylceramide	22
Methyl hexadecenoate (<i>trans</i> -9)	58, 64	N-Dodecanoyl-D- <i>erythro</i> -sphingosine	8
Methyl hexanoate	52	N-Dodecanoyl-NBD-ceramide trihexoside	32, 35, 98
Methyl jacarate	61, 68	N-Dodecanoyl-NBD-D- <i>erythro</i> -dihydrosphingosine	16, 96
Methyl malvalate	78	N-Dodecanoyl-NBD-D- <i>erythro</i> -sphingosine	15, 95
Methyl nonadecanoate	55	N-Dodecanoyl-NBD-galactosylceramide	23, 34, 97
Methyl nonadecenoate (<i>cis</i> -10)	62	N-Dodecanoyl-NBD-lactosylceramide	30, 35, 98
Methyl nonanoate	53	N-Dodecanoyl-NBD-L- <i>threo</i> -dihydrosphingosine	16, 96
Methyl octacosanoate	57	N-Dodecanoyl-NBD-L- <i>threo</i> -sphingosine	15, 96
Methyl octadecadienoate (all <i>cis</i> -9,12)	60	N-Dodecanoyl-NBD-phytosphingosine	16, 96
Methyl octadecadienoate (all <i>trans</i> -9,12)	60, 65	N-Dodecanoyl-NBD-sphingosylphosphorylcholine	20, 97
Methyl octadecanoate	55	N-Dodecanoyl-NBD-sulfatide	28, 34, 97
Methyl octadecatrienoate (all <i>cis</i> -6,9,12)	61	N-Dodecanoyl-sulfatide	27
Methyl octadecatrienoate (all <i>cis</i> -9,12,15)	61	N-Dotriacontanoyl-D- <i>erythro</i> -sphingosine	10
Methyl octadecenoate (<i>cis</i> -11)	60	N-Eicosanoyl-D- <i>erythro</i> -sphingosylphosphorylcholine	18
Methyl octadecenoate (<i>cis</i> -9)	59	Neutral Glycosphingolipid Mixture	39, 92
Methyl octadecenoate (<i>trans</i> -11)	60, 65	N-Glycinated galactosylsphingosine	16, 96
Methyl octadecenoate (<i>trans</i> -9)	59, 65	N-Glycinated glucosylsphingosine	25
Methyl octanoate	53	N-Glycinated lactosylsphingosine	29
Methyl pentadecanoate	54	N-Glycinated <i>lyso</i> -ceramide trihexoside	31
Methyl pentadecenoate (<i>cis</i> -10)	58	N-Glycinated <i>lyso</i> -sulfatide	26
Methyl punicate	61, 67	N-Heptadecanoyl-ceramide trihexoside	32
Methyl stearidonate (all <i>cis</i> -6,9,12,15)	61	N-Heptadecanoyl-D- <i>erythro</i> -dihydrosphingosine	12
Methyl sterculate	78	N-Heptadecanoyl-D- <i>erythro</i> -sphingosine	9
Methyl tetracosanoate	57	N-Heptadecanoyl-lactosylceramide	30
Methyl tetracosenoate (<i>cis</i> -15)	64	N-Heptadecanoyl-sphingosylphosphorylcholine	18
Methyl tetradecanoate	54	N-Heptadecanoyl-sulfatide	27
Methyl tetradecenoate (<i>cis</i> -9)	58	N-Hexadecanoyl-ceramide trihexoside	31
Methyl triacontanoate	57	N-Hexadecanoyl-D- <i>erythro</i> -dihydrosphingosine	12
Methyl tricosanoate	56	N-Hexadecanoyl-D- <i>erythro</i> -sphingosine	9
Methyl tridecanoate	54	N-Hexadecanoyl-D- <i>erythro</i> -sphingosine (C16 sphingolipid base)	11
Methyl undecanoate	53	N-Hexadecanoyl-D- <i>erythro</i> -sphingosine-1-phosphate	20
Mixed Gangliosides, purified, bovine	39, 101	N-Hexadecanoyl-lactosylceramide	29
Mixed Gangliosides, purified, porcine	39, 104	N-Hexadecanoyl-phytosphingosine	14
Monogalactosyldiglyceride (hydrogenated), plant	52, 101	N-Hexadecanoyl-sulfatide	27
Monosialoganglioside GM ₁	36, 101	N-Hexanoyl-biotin-D- <i>erythro</i> -sphingosine	11, 98
Monosialoganglioside GM ₂	37, 103	N-Hexanoyl-biotin-disialoganglioside GD ₃	39, 99
Monosialoganglioside GM ₃	37, 103	N-Hexanoyl-biotin-galactosylceramide	23, 98
Monosialoganglioside GM ₄	38, 104	N-Hexanoyl-biotin-glucosylceramide	26, 98
Monosialoganglioside Mixture	39, 92	N-Hexanoyl-biotin-lactosylceramide	30, 99
		N-Hexanoyl-biotin-monosialoganglioside GM ₁	37, 99
N		N-Hexanoyl-biotin-sulfatide	28, 99
N-(1-Adamantaneacetyl)-ceramide trihexoside	32	N-Hexanoyl-D- <i>erythro</i> -dihydrosphingosine	11
N-(1-Adamantaneacetyl)-galactosylceramide	23, 45	N-Hexanoyl-D- <i>erythro</i> -sphingosine	7
N-(1-Adamantaneacetyl)-glucosylceramide	26, 44	N-Hexanoyl-D- <i>threo</i> -sphingosine	8
N-(30-Linoleoyloxy-triacontanoyl)-sphingosine	10	N-Hexanoyl-glucosylceramide	25
N-(32-Linoleoyloxy-dotriacontanoyl)-sphingosine-D9	10, 93	N-Hexanoyl-L- <i>erythro</i> -sphingosine	7
N-(R,S)- <i>alpha</i> -Hydroxydodecanoyl-D- <i>erythro</i> -dihydrosphingosine	12	N-Hexanoyl-L- <i>threo</i> -sphingosine	7
N-(R,S)- <i>alpha</i> -Hydroxyhexadecanoyl-D- <i>erythro</i> -dihydrosphingosine	13	N-Hexanoyl-NBD-D- <i>erythro</i> -dihydrosphingosine	16, 96
N-(R,S)- <i>alpha</i> -Hydroxyoctadecanoyl-D- <i>erythro</i> -dihydrosphingosine	13	N-Hexanoyl-NBD-D- <i>erythro</i> -sphingosine	15, 95
N-(R,S)- <i>alpha</i> -Hydroxyoctadecanoyl-D- <i>erythro</i> -sphingosine	11	N-Hexanoyl-NBD-galactosylceramide	23, 34, 97
N,N-Dihexyl-D- <i>erythro</i> -sphingosine	6, 45	N-Hexanoyl-NBD-glucosylceramide	26, 34, 97
N,N-Dimethyl-D- <i>erythro</i> -sphingosine	5, 45	N-Hexanoyl-NBD-lactosylceramide	30, 35, 98
N-1- ¹³ C-Hexadecanoyl-D- <i>erythro</i> -sphingosylphosphorylcholine	18, 93	N-Hexanoyl-NBD-L- <i>threo</i> -dihydrosphingosine	15, 96
N-Acetyl-D- <i>erythro</i> -dihydrosphingosine	11	N-Hexanoyl-NBD-L- <i>threo</i> -sphingosine	15, 96
N-Acetyl-D- <i>erythro</i> -sphingosine	7	N-Hexanoyl-NBD-phytosphingosine	16, 96
N-Acetyl-D- <i>erythro</i> -sphingosine (C14 sphingolipid base)	11	N-Hexanoyl-NBD-sphingosylphosphorylcholine	20, 97
N-Acetyl-L- <i>erythro</i> -sphingosine	7	N-Hexanoyl-phytosphingosine	14
N-Acetyl-L- <i>threo</i> -sphingosine	7	N-Nonadecanoyl-D- <i>erythro</i> -sphingosine	10
N-Acetyl-phytosphingosine	14	N-Nonadecanoyl-sulfatide	27
N-Acetyl-psychosine	22	N-Octadecanoyl-ceramide trihexoside	32
N-Acetyl-sphingosylphosphorylcholine	17	N-Octadecanoyl-D ₃₅ -psychosine	23, 33, 94
N-Acetyl-sulfatide	27	N-Octadecanoyl-D- <i>erythro</i> -dihydrosphingosine	12
N-Acyl-D- <i>erythro</i> -sphingosylphosphorylethanolamine	18	N-Octadecanoyl-D- <i>erythro</i> -sphingosine	9
		N-Octadecanoyl-D- <i>threo</i> -sphingosine	9
		N-Octadecanoyl-L- <i>erythro</i> -sphingosine	9
		N-Octadecanoyl-L- <i>threo</i> -sphingosine	9
		N-Octadecanoyl-phytosphingosine	14

N-Octadecanoyl-sphingosylphosphorylcholine	18	Phosphatidylethanolamine, plant	47, 102
N-Octadecanoyl-sulfated-lactosylceramide	28, 30	Phosphatidylinositol 3-phosphate, dipalmitoyl	51
N-Octadecanoyl-sulfatide	27	Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl	51
N-Octadecenoyl-(<i>cis</i> -9)-D- <i>erythro</i> -sphingosine	9	Phosphatidylinositol, plant, soy	46, 103
N-Octadecenoyl-(<i>cis</i> -9)-sulfatide	27	Phosphatidylinositol, plant, wheat germ	46, 100
N-Octanoyl- <i>beta</i> -D-galactosylceramide	22	Phosphatidylserine, bovine	46, 100
N-Octanoyl-D- <i>erythro</i> -dihydrosphingosine	12	Phytanic acid	77
N-Octanoyl-D- <i>erythro</i> -sphingosine	8	Phytosphingosine	5
N-Octanoyl-D- <i>threo</i> -sphingosine	8	Plant Sterol Mixture	82
N-Octanoyl-L- <i>threo</i> -sphingosine	8	Plant Sterols Kit	82
N-Octanoyl-phytosphingosine	14	Polar Lipid Mixture	91
N-Oleoylethanolamine	41, 78	Psychosine (free amine form)	22
N- <i>omega</i> -CD ₃ -Hexadecanoyl-glucopsychosine	25, 33, 94	Psychosine, synthetic	22
N- <i>omega</i> -CD ₃ -Hexadecanoyl-lactosylceramide	30, 34, 94	PUFA-1	84
N- <i>omega</i> -CD ₃ -Octadecanoyl disialoganglioside GD ₃	38, 95	PUFA-2	84
N- <i>omega</i> -CD ₃ -Octadecanoyl monosialoganglioside GM ₁	37, 94	PUFA-3	85
N- <i>omega</i> -CD ₃ -Octadecanoyl monosialoganglioside GM ₂	37, 95	R	
N- <i>omega</i> -CD ₃ -Octadecanoyl monosialoganglioside GM ₃	37, 95	<i>rac</i> -5,7-Dimethyltocol	80
N- <i>omega</i> -CD ₃ -Octadecanoyl-ceramide trihexoside	32, 34, 94	<i>rac-alpha</i> -Tocopherol	79
N- <i>omega</i> -CD ₃ -Octadecanoyl-D- <i>erythro</i> -dihydrosphingosine	12, 93	<i>rac-beta</i> -Tocopherol	79
N- <i>omega</i> -CD ₃ -Octadecanoyl-D- <i>erythro</i> -sphingosine	9, 93	<i>rac-gamma</i> -Tocopherol	79
N- <i>omega</i> -CD ₃ -Octadecanoyl-D- <i>erythro</i> -sphingosine-1-phosphate	20, 93	Rapeseed Oil Reference Mixture	86
N- <i>omega</i> -CD ₃ -Octadecanoyl-phytosphingosine	15, 93	RM-1 Mixture	86
N- <i>omega</i> -CD ₃ -Octadecanoyl-sulfatide	28, 33, 94	RM-2 Mixture	86
N- <i>omega</i> -Hydroxytriacontanoyl-D- <i>erythro</i> -sphingosine	10	RM-3 Mixture	87
Nonadecanoic acid	55	RM-4 Mixture	87
Nonadecenoic acid (<i>cis</i> -10)	62	RM-5 Mixture	87
Nonanoic acid	53	RM-6 Mixture	87
Non-Polar Lipid Mixture A	92	Royal Jelly acid	74
Non-Polar Lipid Mixture B	92	S	
Non-Volatile Acid Mixture	91	Sphingolipid Mixture	92
N-Pentadecanoyl-D- <i>erythro</i> -sphingosine	8	Sphingomyelin, bovine	17, 100
N-Pentadecanoyl-psychosine	22	Sphingomyelin, buttermilk	17, 102
N-Tetracosanoyl-D- <i>erythro</i> -dihydrosphingosine	12	Sphingomyelin, egg	17, 103
N-Tetracosanoyl-D- <i>erythro</i> -sphingosine	10	Sphingomyelin, porcine RBC	17, 102
N-Tetracosanoyl-phytosphingosine	14	Sphingosylphosphorylcholine	19
N-Tetracosanoyl-sulfatide	28	Stearidonic acid (all <i>cis</i> -6,9,12,15)	61
N-Tetracosenoyl-(<i>cis</i> -15)-D- <i>erythro</i> -sphingosine	10	Sterculic acid	78
N-Tetracosenoyl-(<i>cis</i> -15)-sulfatide	28	SteryI Glucosides	83
N-Triacontanoyl-D- <i>erythro</i> -sphingosine	10	Stigmasterol	82
N-Tricosanoyl-ceramide trihexoside	32	Sulfatides, bovine	26, 100
O		T	
Octadecadienoic acid (all <i>cis</i> -9,12)	60	Tetracosanoic acid	56
Octadecadienoic acid (all <i>trans</i> -9,12)	60, 65	Tetracosenoic acid (<i>cis</i> -15)	64
Octadecanoic acid	55	Tetradecanoic acid	54
Octadecatrienoic acid (all <i>cis</i> -6,9,12)	61	Tetradecenoic acid (<i>cis</i> -9)	57
Octadecatrienoic acid (all <i>cis</i> -9,12,15)	60	Tetrasialoganglioside GQ _{1b}	39, 104
Octadecenoic acid (<i>cis</i> -11)	59	Tocol	80
Octadecenoic acid (<i>cis</i> -9)	59	Tricosanoic acid	56
Octadecenoic acid (<i>trans</i> -11)	60, 65	Tridecanoic acid	54
Octadecenoic acid (<i>trans</i> -9)	59, 64	Trisialoganglioside GT _{1b}	39, 101
Octanoic acid	53	U	
P		Undecanoic acid	53
Pentadecanoic acid	54	V	
Pentadecenoic acid (<i>cis</i> -10)	58	Volatile Acid Mixture	91
Phosphatidic acid, egg	46, 100	W	
Phosphatidylethanolamine, bovine	47	WSFA-2 Mixture	90
Phosphatidylethanolamine, egg	47, 100	WSFA-4 Mixture	90

Category Index

2-Hydroxy Fatty Acids & Methyl Esters

2-Hydroxy Methyl Ester Mixture	70
2-Hydroxydecanoic acid	68
2-Hydroxydocosanoic acid	70
2-Hydroxydodecanoic acid	68
2-Hydroxyeicosanoic acid	69
2-Hydroxyhexadecanoic acid	69
2-Hydroxyoctadecanoic acid	69
2-Hydroxytetraacosanoic acid	70
2-Hydroxytetradecanoic acid	69
2-Hydroxytricosanoic acid	70
Methyl 2-hydroxydecanoate	68
Methyl 2-hydroxydocosanoate	70
Methyl 2-hydroxydodecanoate	69
Methyl 2-hydroxyeicosanoate	70
Methyl 2-hydroxyhexadecanoate	69
Methyl 2-hydroxyoctadecanoate	69
Methyl 2-hydroxytetraacosanoate	70
Methyl 2-hydroxytetradecanoate	69
Methyl 2-hydroxytricosanoate	70

2-Hydroxy-Ceramides

N-(R,S)- <i>alpha</i> -Hydroxyoctadecanoyl-D- <i>erythro</i> -sphingosine	11
---------------------------------------------------------------------------	----

2-Hydroxy-Dihydroceramides

N-(R,S)- <i>alpha</i> -Hydroxydodecanoyl-D- <i>erythro</i> -dihydrosphingosine	12
N-(R,S)- <i>alpha</i> -Hydroxyhexadecanoyl-D- <i>erythro</i> -dihydrosphingosine	13
N-(R,S)- <i>alpha</i> -Hydroxyoctadecanoyl-D- <i>erythro</i> -dihydrosphingosine	13

3-Hydroxy Fatty Acids & Methyl Esters

3-Hydroxydecanoic acid	72
3-Hydroxydodecanoic acid	72
3-Hydroxyheptadecanoic acid	73
3-Hydroxyhexadecanoic acid	73
3-Hydroxyhexanoic acid	71
3-Hydroxynonanoic acid	71
3-Hydroxyoctadecanoic acid	73
3-Hydroxyoctanoic acid	71
3-Hydroxytetradecanoic acid	73
3-Hydroxytridecanoic acid	72
3-Hydroxyundecanoic acid	72
Methyl 3-hydroxydecanoate	72
Methyl 3-hydroxydodecanoate	72
Methyl 3-hydroxyheptadecanoate	73
Methyl 3-hydroxyhexadecanoate	73
Methyl 3-hydroxyhexanoate	71
Methyl 3-hydroxynonanoate	71
Methyl 3-hydroxyoctadecanoate	73
Methyl 3-hydroxyoctanoate	71
Methyl 3-hydroxytetradecanoate	73
Methyl 3-hydroxytridecanoate	72
Methyl 3-hydroxyundecanoate	72

3-keto-Dihydrosphingosines

3-keto-C12-Dihydrosphingosine•HCl	5
3-keto-C6-Dihydrosphingosine•HCl	5
3-keto-C8-Dihydrosphingosine•HCl	5
3-keto-Dihydrosphingosine•HCl	4

Anteiso-Fatty Acids & Methyl Esters

12-Methyltetradecanoic acid	76
14-Methylhexadecanoic acid	77

Methyl 12-methyltetradecanoate	77
Methyl 13-methylpentadecanoate	77
Methyl 14-methylhexadecanoate	77

Antibodies Directed Against Glycolipids

Anti-ganglioside asialo GM ₁	40
Anti-ganglioside asialo GM ₂	40
Anti-ganglioside GD _{1b}	41
Anti-ganglioside GD ₃	40
Anti-ganglioside GM ₁	40
Anti-globoside GL-4	41

AOCS Animal & Vegetable Oil Reference Mixtures

Rapeseed Oil Reference Mixture	86
RM-1 Mixture	86
RM-2 Mixture	86
RM-3 Mixture	87
RM-4 Mixture	87
RM-5 Mixture	87
RM-6 Mixture	87

Bacterial Tetraethers

Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i> , (>50% pure)	51
Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i> , (>95% pure)	51

Biochemical Research Standard Mixtures

Non-Polar Lipid Mixture A	92
Non-Polar Lipid Mixture B	92
Polar Lipid Mixture	91
Sphingolipid Mixture	92

Biotin Labeled Standards

N-Hexanoyl-biotin-D- <i>erythro</i> -sphingosine	98
N-Hexanoyl-biotin-galactosylceramide	98
N-Hexanoyl-biotin-glucosylceramide	98
N-Hexanoyl-biotin-lactosylceramide	99
N-Hexanoyl-biotin-monosialoganglioside GM ₁	99
N-Hexanoyl-biotin-sulfatide	99

Branched & Cyclic Fatty Acids

13-Methyltetradecanoic acid	76
15-Methylhexadecanoic acid	76
Methyl 11-methyldodecanoate	75
Methyl 12-methyltridecanoate	75
Methyl 13-methyltetradecanoate	76
Methyl 14-methylpentadecanoate	76
Methyl 15-methylhexadecanoate	76
Methyl 17-methyloctadecanoate	76

Carbohydrate Mixtures

Alditol Acetate Mixture-1	85
Alditol Acetate Mixture-2	85

Ceramide Trihexosides

Ceramide trihexosides	31
Ceramide trihexosides (bottom spot)	31
Ceramide trihexosides (top spot)	31
<i>lyso</i> -Ceramide trihexoside	31
N-(1-Adamantaneacetyl)-ceramide trihexoside	32
N-Dodecanoyl-NBD-ceramide trihexoside	32
N-Glycinated <i>lyso</i> -ceramide trihexoside	31
N-Heptadecanoyl-ceramide trihexoside	32
N-Hexadecanoyl-ceramide trihexoside	31

N-Octadecanoyl-ceramide trihexoside	32	N-Heptadecanoyl-D- <i>erythro</i> -dihydrosphingosine	12
N- <i>omega</i> -CD ₃ -Octadecanoyl-ceramide trihexoside	32	N-Hexadecanoyl-D- <i>erythro</i> -dihydrosphingosine	12
N-Tricosanoyl-ceramide trihexoside	32	N-Hexanoyl-D- <i>erythro</i> -dihydrosphingosine	11
		N-Octadecanoyl-D- <i>erythro</i> -dihydrosphingosine	12
		N-Octanoyl-D- <i>erythro</i> -dihydrosphingosine	12
		N- <i>omega</i> -CD ₃ -Octadecanoyl-D- <i>erythro</i> -dihydrosphingosine	12
Ceramides		N-Tetracosanoyl-D- <i>erythro</i> -dihydrosphingosine	12
Ceramides (hydroxy)	13		
Ceramides (mixture)	13	Dihydrosphingosines	
Ceramides (non-hydroxy)	13	D,L-C16-Dihydrosphingosine (mixed isomers)	4
N-(30-Linoleoyloxy-triacontanoyl)-sphingosine	10	D,L- <i>erythro</i> -C20-Dihydrosphingosine	4
N-(32-Linoleoyloxy-dotriacontanoyl)-sphingosine-D9	10	D,L- <i>erythro</i> -Dihydrosphingosine	4
N-Acetyl-D- <i>erythro</i> -sphingosine	7	D- <i>erythro</i> -C20-Dihydrosphingosine	4
N-Acetyl-D- <i>erythro</i> -sphingosine (C14 sphingolipid base)	11	D- <i>erythro</i> -Dihydrosphingosine	3
N-Acetyl-L- <i>erythro</i> -sphingosine	7	D- <i>threo</i> -Dihydrosphingosine	4
N-Acetyl-L- <i>threo</i> -sphingosine	7	L- <i>erythro</i> -Dihydrosphingosine	3
N-Decanoyl-D- <i>erythro</i> -sphingosine	8	L- <i>threo</i> -Dihydrosphingosine (Safingol)	3
N-Dodecanoyl-D- <i>erythro</i> -sphingosine	8		
N-Dotriacontanoyl-D- <i>erythro</i> -sphingosine	10	Enzyme Inhibitors	
N-Heptadecanoyl-D- <i>erythro</i> -sphingosine	9	2-Acetyl-4-(1R, 2S, 3R, 4-tetrahydroxybutyl)-imidazole	44
N-Hexadecanoyl-D- <i>erythro</i> -sphingosine	9	2-Fluoropalmitic acid	44
N-Hexadecanoyl-D- <i>erythro</i> -sphingosine (C16 sphingolipid base)	11	Conduritol B Epoxide	42
N-Hexanoyl-biotin-D- <i>erythro</i> -sphingosine	11	D,L- <i>erythro</i> -PDMP	43
N-Hexanoyl-D- <i>erythro</i> -sphingosine	7	D,L- <i>erythro</i> -PPMP	43
N-Hexanoyl-D- <i>threo</i> -sphingosine	8	D,L- <i>threo</i> -PDMP	43
N-Hexanoyl-L- <i>erythro</i> -sphingosine	7	D,L- <i>threo</i> -PPMP	43
N-Hexanoyl-L- <i>threo</i> -sphingosine	7	D-MAPP	42
N-Nonadecanoyl-D- <i>erythro</i> -sphingosine	10	D- <i>threo</i> -PDMP	43
N-Octadecanoyl-D- <i>erythro</i> -sphingosine	9	D- <i>threo</i> -PPMP	44
N-Octadecanoyl-D- <i>threo</i> -sphingosine	9	L-MAPP	42
N-Octadecanoyl-L- <i>erythro</i> -sphingosine	9	L- <i>threo</i> -Dihydrosphingosine (Safingol)	42
N-Octadecanoyl-L- <i>threo</i> -sphingosine	9	L- <i>threo</i> -PDMP	43
N-Octadecenoyl-(<i>cis</i> -9)-D- <i>erythro</i> -sphingosine	9	L- <i>threo</i> -PPMP	44
N-Octanoyl-D- <i>erythro</i> -sphingosine	8	Methyl 2-fluoropalmitate	44
N-Octanoyl-D- <i>threo</i> -sphingosine	8	N-(1-Adamantaneacetyl)-galactosylceramide	45
N-Octanoyl-L- <i>threo</i> -sphingosine	8	N-(1-Adamantaneacetyl)-glucosylceramide	44
N- <i>omega</i> -CD ₃ -Octadecanoyl-D- <i>erythro</i> -sphingosine	9	N,N-Dihexyl-D- <i>erythro</i> -sphingosine	45
N- <i>omega</i> -Hydroxytriacontanoyl-D- <i>erythro</i> -sphingosine	10	N,N-Dimethyl-D- <i>erythro</i> -sphingosine	45
N-Pentadecanoyl-D- <i>erythro</i> -sphingosine	8	N-Oleylethanolamine	41
N-Tetracosanoyl-D- <i>erythro</i> -sphingosine	10		
N-Tetracosenoyl-(<i>cis</i> -15)-D- <i>erythro</i> -sphingosine	10	Fluorescent Glycolipids	
N-Triacontanoyl-D- <i>erythro</i> -sphingosine	10	Lissamine-rhodamine B-dodecanoyl-galactosylceramide	35
		N-Dodecanoyl-NBD-ceramide trihexoside	35
Cholestane Derivatives		N-Dodecanoyl-NBD-D- <i>erythro</i> -dihydrosphingosine	16
5- <i>alpha</i> -Cholestane	81	N-Dodecanoyl-NBD-D- <i>erythro</i> -sphingosine	15
Cholesterol	81	N-Dodecanoyl-NBD-galactosylceramide	34
Coprostanol	82	N-Dodecanoyl-NBD-lactosylceramide	35
		N-Dodecanoyl-NBD-L- <i>threo</i> -dihydrosphingosine	16
Conjugated Linoleic Acid Isomers (CLA)		N-Dodecanoyl-NBD-L- <i>threo</i> -sphingosine	15
10(E),12(Z)-Octadecadienoic acid	67	N-Dodecanoyl-NBD-phytosphingosine	16
9(E),11(E)-Octadecadienoic acid	67	N-Dodecanoyl-NBD-sphingosylphosphorylcholine	20
9(Z),11(E)-Octadecadienoic acid	66	N-Dodecanoyl-NBD-sulfatide	34
9(Z),11(E)-Octadecadienoic acid (Na+ salt)	66	N-Hexanoyl-NBD-D- <i>erythro</i> -dihydrosphingosine	16
9(Z),11(Z)-Octadecadienoic acid	67	N-Hexanoyl-NBD-D- <i>erythro</i> -sphingosine	15
Methyl 10(E),12(Z)-octadecadienoate	67	N-Hexanoyl-NBD-galactosylceramide	34
Methyl 9(E),11(E)-octadecadienoate	67	N-Hexanoyl-NBD-glucosylceramide	34
Methyl 9(Z),11(E)-octadecadienoate	66	N-Hexanoyl-NBD-lactosylceramide	35
Methyl 9(Z),11(Z)-octadecadienoate	67	N-Hexanoyl-NBD-L- <i>threo</i> -dihydrosphingosine	15
Methyl <i>alpha</i> -eleostearate	68	N-Hexanoyl-NBD-L- <i>threo</i> -sphingosine	15
Methyl jacarate	68	N-Hexanoyl-NBD-phytosphingosine	16
Methyl punicate	67	N-Hexanoyl-NBD-sphingosylphosphorylcholine	20
		Fluorescent Standards	
Cyclopropyl Fatty Acids & Methyl Esters		Lissamine-rhodamine B-dodecanoyl-galactosylceramide	97
<i>cis</i> -9,10-Methyleneoctadecanoic acid (all <i>cis</i> -9,10)	78	N-Dodecanoyl-NBD-ceramide trihexoside	98
Methyl <i>cis</i> -9,10-methyleneoctadecanoate	78	N-Dodecanoyl-NBD-D- <i>erythro</i> -dihydrosphingosine	96
Methyl malvalate	78	N-Dodecanoyl-NBD-D- <i>erythro</i> -sphingosine	95
Methyl sterculate	78	N-Dodecanoyl-NBD-galactosylceramide	97
Sterculic acid	78	N-Dodecanoyl-NBD-lactosylceramide	98
Dihydroceramides			
N-Acetyl-D- <i>erythro</i> -dihydrosphingosine	11		

Phosphatidylethanolamine, bovine	47	Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl	51
Phosphatidylethanolamine, egg	47		
Phosphatidylethanolamine, plant	47	Phosphosphingolipids	
Phosphatidylinositol, plant, soy	46	N-1-13C-Hexadecanoyl-D-erythro-sphingosylphosphorylcholine	18
Phosphatidylinositol, plant, wheat germ	46	N-Acetyl-sphingosylphosphorylcholine	17
Phosphatidylserine, bovine	46	N-Acyl-D-erythro-sphingosylphosphorylethanolamine	18
		N-Docosanoyl-D-erythro-sphingosylphosphorylcholine	18
		N-Eicosanoyl-D-erythro-sphingosylphosphorylcholine	18
		N-Heptadecanoyl-sphingosylphosphorylcholine	18
		N-Octadecanoyl-sphingosylphosphorylcholine	18
		Sphingomyelin, bovine	17
		Sphingomyelin, buttermilk	17
		Sphingomyelin, egg	17
		Sphingomyelin, porcine RBC	17
Omega Hydroxy Fatty Acids		Phytoceramides	
15-Hydroxypentadecanoic acid	74	N-Acetyl-phytosphingosine	14
17-Hydroxyheptadecanoic acid	74	N-Hexadecanoyl-phytosphingosine	14
20-Hydroxyeicosanoic acid	74	N-Hexanoyl-phytosphingosine	14
22-Hydroxydocosanoic acid	75	N-Octadecanoyl-phytosphingosine	14
Methyl 15-hydroxypentadecanoate	74	N-Octanoyl-phytosphingosine	14
Methyl 17-hydroxyheptadecanoate	74	N-omega-CD ₃ -Octadecanoyl-phytosphingosine	15, 93
Methyl 20-hydroxyeicosanoate	74	N-Tetracosanoyl-phytosphingosine	14
Methyl 21-hydroxyheneicosanoate	75		
Methyl 22-hydroxydocosanoate	75	Phytosphingosines	
Methyl 27-hydroxyheptacosanoate	75	Phytosphingosine	5
Methyl 30-hydroxytriacontanoate	75		
Royal Jelly acid	74	Plant Sterols & Steryl Glucosides	
		beta-Sitosterol	82
Other Branched Fatty Acids		Ergosterol	82
D,L-2,6-Dimethylheptanoic acid	77	Esterified Steryl Glucosides	83
Methyl 10-methylhexadecanoate	77	Lanosterol	82
Phytanic acid	77	Plant Sterol Mixture	82
		Plant Sterols Kit	82
		Steryl Glucosides	83
		Stigmasterol	82
Other Fatty Acid Methyl Ester Mixtures		Polyunsaturated Fatty Acid Methyl Ester Mixtures	
2-Hydroxy Methyl Ester Mixture	85	PUFA-1	84
Cis-Trans FAME Isomer Standard Mixture	85	PUFA-2	84
Long Chain Fatty Acid Methyl Ester Mixture	85	PUFA-3	85
Phosphates		Saturated Fatty Acids & Methyl Esters	
D-erythro-Dihydrosphingosine-1-phosphate	20	Docosanoic acid	56
D-erythro-Sphingosine-1-phosphate	19	Dodecanoic acid	53
N-Hexadecanoyl-D-erythro-sphingosine-1-phosphate	20	Eicosanoic acid	55
N-omega-CD ₃ -Octadecanoyl-D-erythro-sphingosine-1-phosphate	20	Heneicosanoic acid	56
		Heptadecanoic acid	55
		Heptanoic acid	52
		Hexacosanoic acid	57
		Hexadecanoic acid	54
		Methyl decanoate	53
		Methyl docosanoate	56
		Methyl dodecanoate	54
		Methyl dotriacontanoate	57
		Methyl eicosanoate	56
		Methyl heneicosanoate	56
		Methyl heptadecanoate	55
		Methyl heptanoate	52
		Methyl hexacosanoate	57
		Methyl hexadecanoate	55
		Methyl hexanoate	52
		Methyl nonadecanoate	55
		Methyl nonanoate	53
		Methyl octacosanoate	57
		Methyl octadecanoate	55
		Methyl octanoate	53
		Methyl pentadecanoate	54
		Methyl tetracosanoate	57
Phosphatidic Acid Derivatives			
1,2-Dimyristoyl-sn-glycero-3-phosphatidic acid	47		
1,2-Dipalmitoyl-sn-glycero-3-phosphatidic acid	48		
1,2-Distearoyl-sn-glycero-3-phosphatidic acid	48		
Phosphatidylcholines			
1,2-Diheptadecanoyl-sn-glycero-3-phosphorylcholine	48		
1,2-Dilauroyl-sn-glycero-3-phosphorylcholine	48		
1,2-Dimyristoyl-sn-glycero-3-phosphorylcholine	48		
1,2-Dipalmitoyl-sn-glycero-3-phosphorylcholine	48		
1,2-Distearoyl-sn-glycero-3-phosphorylcholine	49		
1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylcholine	49		
1-Palmitoyl-sn-glycero-3-phosphorylcholine	49		
Phosphatidylethanolamines			
1,2-Dilauroyl-sn-glycero-3-phosphorylethanolamine	50		
1,2-Dimyristoyl-sn-glycero-3-phosphorylethanolamine	50		
1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine	50		
1,2-Distearoyl-phosphatidylethanolamine-methyl-polyethyleneglycol conjugate-2000	51		
1,2-Distearoyl-sn-glycero-3-phosphorylethanolamine	50		
Phosphatidylglycerols			
1,2-Dilauroyl-sn-glycero-3-phosphorylglycerol	49		
1,2-Dimyristoyl-sn-glycero-3-phosphorylglycerol	49		
1,2-Dipalmitoyl-sn-glycero-3-phosphorylglycerol	49		
1,2-Distearoyl-sn-glycero-3-phosphorylglycerol	50		
1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylglycerol	50		
Phosphatidylinositol Phosphates			
Phosphatidylinositol 3-phosphate, dipalmitoyl	51		

Methyl tetradecanoate	54	N-Tetracosenoyl-(<i>cis</i> -15)-sulfatide	28
Methyl triacontanoate	57	Sulfatides, bovine	26
Methyl tricosanoate	56		
Methyl tridecanoate	54	Tocopherols	
Methyl undecanoate	53	(+)- <i>delta</i> -Tocopherol	79
Nonadecanoic acid	55	<i>rac</i> -5,7-Dimethyltolcol	80
Nonanoic acid	53	<i>rac-alpha</i> -Tocopherol	79
Octadecanoic acid	55	<i>rac-beta</i> -Tocopherol	79
Octanoic acid	53	<i>rac-gamma</i> -Tocopherol	79
Pentadecanoic acid	54	Tocol	80
Tetracosanoic acid	56		
Tetradecanoic acid	54	Tocotrienols	
Tricosanoic acid	56	<i>alpha</i> -Tocotrienol	80
Tridecanoic acid	54	<i>beta</i> -Tocotrienol	80
Undecanoic acid	53	<i>delta</i> -Tocotrienol	81
		<i>gamma</i> -Tocotrienol	81
Sphingosines		Trans Fatty Acids & Methyl Esters	
D- <i>erythro</i> -C12-Sphingosine	2	<i>Cis-Trans</i> FAME Isomer Standard Mixture	65
D- <i>erythro</i> -C14-Sphingosine	2	Hexadecenoic acid (<i>trans</i> -9)	64
D- <i>erythro</i> -C17-Sphingosine	3	Methyl hexadecenoate (<i>trans</i> -9)	64
D- <i>erythro</i> -C20-Sphingosine	3	Methyl octadecadienoate (all <i>trans</i> -9,12)	65
D- <i>erythro</i> -Sphingosine	2	Methyl octadecenoate (<i>trans</i> -11)	65
D- <i>erythro</i> -Sphingosine, D9	2	Methyl octadecenoate (<i>trans</i> -9)	65
L- <i>erythro</i> -Sphingosine	2	Octadecadienoic acid (all <i>trans</i> -9,12)	65
L- <i>threo</i> -Sphingosine	2	Octadecenoic acid (<i>trans</i> -11)	65
N,N-Dihexyl-D- <i>erythro</i> -sphingosine	6	Octadecenoic acid (<i>trans</i> -9)	64
N,N-Dimethyl-D- <i>erythro</i> -sphingosine	5		
		Unsaturated Fatty Acids & Methyl Esters	
Sphingosylphosphorylcholines		11-Hexadecenoic acid (92% <i>cis</i> , 8% <i>trans</i>)	59
D- <i>erythro</i> -Sphingosylphosphorylcholine	19	Docosahexaenoic acid (all <i>cis</i> -4,7,10,13,16,19)	64
L- <i>threo</i> -Sphingosylphosphorylcholine	19	Docosapentaenoic acid (all <i>cis</i> -7,10,13,16,19)	63
<i>lyso</i> -Dihydrosphingomyelin	19	Docosenoic acid (<i>cis</i> -13)	63
Sphingosylphosphorylcholine	19	Eicosadienoic acid (all <i>cis</i> -11,14)	62
		Eicosapentaenoic acid (all <i>cis</i> -5,8,11,14,17)	63
Stable Isotope Labeled Standards		Eicosatetraenoic acid (all <i>cis</i> -5,8,11,14)	63
¹³ C ₆ -Glucosylsphingosine	94	Eicosenoic acid (<i>cis</i> -11)	62
D- <i>erythro</i> -Sphingosine, D9	93	Heptadecenoic acid (<i>cis</i> -10)	59
N-(32-Linoleoyloxy-dotriacontanoyl)-sphingosine-D9	93	Hexadecenoic acid (<i>cis</i> -6)	58
N-1- ¹³ C-Hexadecanoyl-D- <i>erythro</i> -		Hexadecenoic acid (<i>cis</i> -9)	58
sphingosylphosphorylcholine	93	Hexadecenoic acid (<i>trans</i> -9)	58
N-Octadecanoyl-D ₃₅ -psychosine	94	Methyl <i>alpha</i> -eleostearate	61
N- <i>omega</i> -CD ₃ -Hexadecanoyl-glucopsychosine	94	Methyl docosahexaenoate (all <i>cis</i> -4,7,10,13,16,19)	64
N- <i>omega</i> -CD ₃ -Hexadecanoyl-lactosylceramide	94	Methyl docosapentaenoate (all <i>cis</i> -7,10,13,16,19)	63
N- <i>omega</i> -CD ₃ -Octadecanoyl disialoganglioside GD ₃	95	Methyl docosenoate (<i>cis</i> -13)	63
N- <i>omega</i> -CD ₃ -Octadecanoyl monosialoganglioside GM ₁	94	Methyl eicosadienoate (all <i>cis</i> -11,14)	62
N- <i>omega</i> -CD ₃ -Octadecanoyl monosialoganglioside GM ₂	95	Methyl eicosapentaenoate (all <i>cis</i> -5,8,11,14,17)	63
N- <i>omega</i> -CD ₃ -Octadecanoyl monosialoganglioside GM ₃	95	Methyl eicosatetraenoate (all <i>cis</i> -5,8,11,14)	63
N- <i>omega</i> -CD ₃ -Octadecanoyl-ceramide trihexoside	94	Methyl eicosatrienoate (all <i>cis</i> -5,8,11)	62
N- <i>omega</i> -CD ₃ -Octadecanoyl-D- <i>erythro</i> -dihydrosphingosine	93	Methyl eicosatrienoate (all <i>cis</i> -8,11,14)	62
N- <i>omega</i> -CD ₃ -Octadecanoyl-D- <i>erythro</i> -sphingosine	93	Methyl eicosenoate (<i>cis</i> -11)	62
N- <i>omega</i> -CD ₃ -Octadecanoyl-D- <i>erythro</i> -sphingosine-1-		Methyl heptadecenoate (<i>cis</i> -10)	59
phosphate	93	Methyl hexadecenoate (<i>cis</i> -9)	58
N- <i>omega</i> -CD ₃ -Octadecanoyl-sulfatide	94	Methyl hexadecenoate (<i>trans</i> -9)	58
		Methyl jacarate	61
Sulfatides		Methyl nonadecenoate (<i>cis</i> -10)	62
<i>lyso</i> -Sulfatide	26	Methyl octadecadienoate (all <i>cis</i> -9,12)	60
N-Acetyl-sulfatide	27	Methyl octadecadienoate (all <i>trans</i> -9,12)	60
N-Dodecanoyl-NBD-sulfatide	28	Methyl octadecatrienoate (all <i>cis</i> -6,9,12)	61
N-Dodecanoyl-sulfatide	27	Methyl octadecatrienoate (all <i>cis</i> -9,12,15)	61
N-Glycinated <i>lyso</i> -sulfatide	26	Methyl octadecenoate (<i>cis</i> -11)	60
N-Heptadecanoyl-sulfatide	27	Methyl octadecenoate (<i>cis</i> -9)	59
N-Hexadecanoyl-sulfatide	27	Methyl octadecenoate (<i>trans</i> -11)	60
N-Hexanoyl-biotin-sulfatide	28	Methyl octadecenoate (<i>trans</i> -9)	59
N-Nonadecanoyl-sulfatide	27	Methyl pentadecenoate (<i>cis</i> -10)	58
N-Octadecanoyl-sulfated-lactosylceramide	28	Methyl punicate	61
N-Octadecanoyl-sulfatide	27	Methyl stearidonate (all <i>cis</i> -6,9,12,15)	61
N-Octadecenoyl-(<i>cis</i> -9)-sulfatide	27	Methyl tetracosenoate (<i>cis</i> -15)	64
N- <i>omega</i> -CD ₃ -Octadecanoyl-sulfatide	28	Methyl tetradecenoate (<i>cis</i> -9)	58
N-Tetracosanoyl-sulfatide	28	Nonadecenoic acid (<i>cis</i> -10)	62

Octadecadienoic acid (all <i>cis</i> -9,12)	60
Octadecadienoic acid (all <i>trans</i> -9,12)	60
Octadecatrienoic acid (all <i>cis</i> -6,9,12)	61
Octadecatrienoic acid (all <i>cis</i> -9,12,15)	60
Octadecenoic acid (<i>cis</i> -11)	59
Octadecenoic acid (<i>cis</i> -9)	59
Octadecenoic acid (<i>trans</i> -11)	60
Octadecenoic acid (<i>trans</i> -9)	59
Pentadecenoic acid (<i>cis</i> -10)	58
Stearidonic acid (all <i>cis</i> -6,9,12,15)	61
Tetracosenoic acid (<i>cis</i> -15)	64
Tetradecenoic acid (<i>cis</i> -9)	57

Unusual Fatty Acids & Derivatives

N-Oleylethanolamine	78
---------------------	----

Water Soluble Fatty Acid Mixtures

WSFA-2 Mixture	90
WSFA-4 Mixture	90

MATREYA

Matreya LLC Ordering Information

Office Hours: Monday - Friday, 8:00am - 4:00pm, Eastern Time EST/EDT

Ordering and Customer Service

Telephone (Toll-Free)	(800) 342 3595
Telephone (world-wide)	(814) 355 1030
Fax (24/7 world-wide)	(814) 355 1031
e-mail	customerservice@matreya.com
e-commerce website	www.matreya.com

Technical Service

Telephone (worldwide)	(814) 355 1030
Fax (worldwide)	(814) 355 1031
e-mail	techservice@matreya.com

Terms

Prices and discounts are subject to change without notice. Freight charges are prepaid and added as a separate charge on the invoice. Orders placed are F.O.B. origin State College, PA

Net 30 Days. (upon credit approval)

Shipping

Standard orders:	FedEx 2 nd -Day
Rush Orders:	FedEx Priority or Standard overnight
International Orders:	FedEx International Priority

Items requiring dry ice for shipping will be charged a \$40 fee.

NO MINIMUM ORDER REQUIRED!

FREE 2-Day shipping on continental US orders of \$500.00 or more.

Return Policy

All returns must have prior authorization. Items incorrectly ordered by the customer and returned to Matreya are subject to a 25% (\$25 minimum) restocking charge. The following items may not be returned: frozen products, custom synthesized products and accommodation orders. If a product has been incorrectly sent to you due to an error on our part, a credit will be issued to your account immediately. Please inspect and verify your order upon receipt. No products will be returned after 30 days (assuming proper storage and handling by customer).

Disclaimer

The information and data included in this catalog are correct and reliable to the best of our knowledge. However, we offer no guarantees and assume no responsibility for this information. No license or immunity under any patent is granted or implied through our sale of any material. Any information sent to us by the customer, relative to his or her interest in Matreya's products, must be submitted voluntarily and without obligation on the part of Matreya.

Matreya LLC, 2178 High Tech Road, State College, PA 16803-1734 USA Federal ID No. 20-1237500

www.matreya.com

MATREYA

International Dealers & Representatives

Australia (South Australia)

Adelab Scientific
36 Holland Street
Thebarton 5031
Tel: 61-88-234-7955
Fax: 61-88-234-7897
Email: info@adelab.com.au
Web: www.adelab.com.au

France

Biovalley
183, Avenue Georges Chemenceau
Nanterre, 92000
Tel: 33-16-007-2020
Fax: 33-16-007-5051
Email: info@biovalley.fr
Web: www.biovalley.fr

Italy

Superchrom S.r.l.
Via G. Miglioli 2/A
Cernusco Sul Naviglio, 20063
Tel: 39-02-738-6315
Fax: 39-027-010-0100
Email: superchrom@superchrom.it
Web: www.superchrom.it

Netherlands

DistriLab BV
Leusderend 26
Leusden, 3832RC
Tel: 31-33-494-7834
Fax: 31-33-432-1441
Email: info@distriLab.nl
Web: www.distriLab.nl

Spain

Teknokroma Analítica, SA
Camí de Can Calders, 14
Apartado de Correos, 147
Sant Cugat del Vallès
Barcelona ES-08173
Tel: 34-93-674-8800
Fax: 34-93-675-2405
Email: export@teknokroma.es
Web: www.teknokroma.es

Taiwan

Super Chroma Enterprise Ltd
5F, 226 Roosevelt Rd, Section 5
Taipei ROC, 116
Tel: 88-622-930-7110
Fax: 88-622-930-7112
Email: sc@superchroma.com.tw
Web: www.superchroma.com.tw

Canada

MJS Biolynx
PO Bag 1150
300 Laurier Blvd
Brockville, Ontario K6V 5W1
Tel: 613-498-2126
Toll free: 888-593-5969
Fax: 613-342-1341
Email: sales@biolynx.ca
Web: www.biolynx.ca

Germany

BIOTREND Chemikalien GmbH
Eupenerstr. 157
Köln, 50933
Tel: 49-221-949-8320
Fax: 49-221-949-8325
Email: info@biotrend.com
Web: www.biotrend.com

Japan

Techno Chemical Corp
27-9, Honkomagome 1 Chome
Bunkyo-ku, Tokyo 113
Tel: 81-33-947-7310
Fax: 81-33-947-7306
Email: info@technochemical.com
Web: www.technochemical.com

New Zealand

Phenomenex NZ Ltd.
PO Box 31-6-1, Milford
North Shore City 0629
Tel: 649-478-0951
Fax: 649-478-0952
Email: info@phenomenex.co.za
Web: www.phenomenex.com

Sweden

Larodan AB
Karolinska Institutet Science Park
Retzius vag 8
Solna 17165
Tel: 464-016-2200
Fax: 464-015-2201
Email: info@larodan.se
Web: www.larodan.se

United Kingdom (England)

Universal Biologicals Ltd.
Passhouse Farmhouse
Papworth St. Agnes
Cambridge CB23 3QU
Tel: 44-148-083-9015
Fax: 44-148-083-1912
Email: info@universalsbiologicals.com
Web: www.universalsbiologicals.com

China

Shenzhen Minn Bolin Bio-Tech Co., LTD
Rm 8A, No. 2, Yihe Bldg, Maoye Cheng
Dafen Village, Buji, Longgang District
Shenzhen, 518112
Tel: 867-556-165-1418 (1428)
Fax: 867-558-209-6552
Email: peacerli@mbolin-lkllabs.com
Web: www.mbolin-lkllabs.com

India

Clementia Biotech
140, First Floor,
RG Mall, Sector - 9 Rohini
New Delhi, 110085
Tel: 91-11-27553034
Email: info@clementiabiotech.com
Web: www.clementiabiotech.com

Korea

Kim & Friends, Inc.
SK Twintech Tower B-304
345-9 Gasan-dong, Geumcheon-gu
Seoul 153-773
Tel: 82-2-2647-6611
Fax: 82-2-2647-6687
Email: kslee@kimnfriends.co.kr
Web: www.kimnfriends.co.kr

Singapore/Malaysia/Indonesia

Precision Technologies Pte Ltd
211 Henderson Rd
13-02 Henderson Industrial Park
Singapore 159552
Tel: 65-6273-4573
Fax: 65-6273-8898
Email: precision@pretech.com.sg
Web: www.pretech.com.sg

Switzerland

VWR International AG-Life Sciences
Lerzenstrasse 16/18
Dietikon, CH-8953
Tel: 41-44-745-1313
Fax: 41-44-745-1410
Email: info@ch.vwr.com
Web: www.vwr.com

2017-2018

www.matreya.com